

# INDEX

- Absorbance, of diarylethene, 263
- Absorbance change, pump intensity versus, 262
- Absorption. *See also* Excited state absorption (ESA); Femtosecond transient absorption measurements; Linear absorption coefficient; Multiphoton absorption; Nonlinear absorption coefficient ( $\beta$ ); Nonlinear absorption data; One-photon absorption (OPA); Steady state absorption spectra; Three-photon absorption; Transient absorption; Two-photon absorption (TPA)
- linear, 205
- nonlinear, 190–191
- Absorption cross sections, 118. *See also* OPA cross section; TPA cross section ( $\delta$ )
- Absorption spectra, 82, 83, 86, 88–89, 97, 101, 102, 103, 181. *See also* TPA spectra
- of anthracene, 160
- of conjugated polymers, 170
- of excited-state xanthone ketyl radicals, 90
- of irradiated stilbene, 58–61
- of 1-methoxynaphthalene, 76–77
- of methyl-substituted ladder-type poly(*p*-phenylene), 163–165
- of 1-NpCH<sub>2</sub>-OBP, 73
- of oligothiophenes, 84
- steady state, 14–15, 38
- transient, 37–45
- Acceptor groups, in symmetric donor-acceptor-donor chromophores, 226
- (Acceptor- $\pi$ )<sub>3</sub>-donor chromophores, 246–247, 248
- Acceptor- $\pi$ -acceptor (A- $\pi$ -A) compounds, symmetric, 215–225
- Acetonitrile, 58
- Acetylene mesomer, of polydiacetylene, 168–169
- Acousto-optic measurements, 145
- Acrylamide, TP initiated polymerization of, 274, 278
- Acrylate monomer, in three-dimensional microfabrication, 289–290
- Acrylates, TP initiated polymerization of, 274, 276, 278, 286, 289
- Acrylonitrile, in 3D data storage, 302
- Addressing, in 3D data storage, 300–301
- Alkenes, infrared multiphoton excitation isomerization of, 260
- Alkoxy groups, in nonlinear absorbing chromophores, 252, 253, 254
- Alkylene groups, in octupolar chromophores, 247, 248
- Alkylhalides, pulse radiolysis in, 58
- Alkylsulfonyl substituent, in symmetric donor-acceptor-donor chromophores, 226
- AM1 approach, 125
- Amino acids, TPEF and, 255

- Amino nitrogen, in octupolar chromophores, 243–244, 245
- N*-(4-Aminophenyl)fulgimide, as 3D data storage material, 299
- 4,4'-Amino-stilbenes, 197, 202
- Amino-substituted distyrylbenzene, 289, 290
- Amplitude, chromophore number and, 18
- AN152, medical applications of, 260, 318
- Angle dependence, between monomer units, 180
- Anisole (ANS), 58–60, 61
- Anisotropy decay times, 18, 19, 20–21
- Anisotropy decay traces, 19, 21–22
- Annihilation processes, 44, 47. *See also* Singlet–singlet annihilation
- Anthracene  
energy gap law and, 70–71  
TPA spectra of, 160
- Anthracene moieties  
in octupolar chromophores, 239  
in symmetric donor-acceptor-donor chromophores, 231  
in symmetric donor- $\pi$ -donor chromophores, 209–210, 210
- Anthraquinones, 64
- Anti-Stokes–Raman scattering, 121
- Apertureless NSOM, in three-dimensional microfabrication, 285, 289
- Aromatic hydrocarbons (AHs)  
energy gap law and, 70–71  
multiphoton absorption by, 157
- Aromatic resonance, 182
- Aromatic rings, TPA data for, 158, 159
- Arylene-ethylene pattern, in octupolar chromophores, 240
- ASPT lasing dye, in upconverted lasing, 321, 322
- Autofluorescence of tissue, 317
- Average initiation rate, of radical polymerization, 279
- Average molecular weight, 178
- Azaxanthone (AX), excited-state ketyl radicals of, 88–89
- Azobenzenes  
donor–acceptor substituted, 187  
in nonlinear absorbing chromophores, 251
- B3LYP geometries, for two-photon absorption, 126–128
- Band formation, 179–180. *See also* Optical band-gap materials
- Barbituric acid moiety, in symmetric acceptor- $\pi$ -acceptor chromophores, 220–221, 222
- Bathochromic shifts, 126–128  
of nonlinear absorbing chromophores, 254  
of octupolar chromophores, 240  
of symmetric donor-acceptor-donor chromophores, 230
- Bay-functionalized perylene bisimide, 7–8
- Beam pulse shapes, 297  
for quantifying TPA cross section, 145–146, 147, 149–150, 151, 152  
in three-dimensional microfabrication, 284, 287
- Beam–sample interaction area, 283
- Benzene(s)  
TPA data for, 158, 159  
TP transition in, 139
- Benzene moiety, in octupolar chromophores, 242, 243
- Benzene ring, in nonlinear absorbing chromophores, 254–255
- Benzocyclobutene, 94–95
- Benzophenone (BP), 4–5  
bond dissociation from  $T_n$  states in, 72–74  
excited-state ketyl radicals of, 85–88, 89–90  
naphthalene  $T_n$  states and, 66–67  
substituent effects on  $T_n$  state lifetimes of, 71–72
- Benzophenone derivatives (BPD), excited-state ketyl radicals of, 85–88, 89
- Benzophenoxazin dye, as 3D data storage material, 299
- 6-Benzothiazol-2-yl-(2-naphthyl) diphenylamine, in optical circuits, 309
- 7-Benzothiazol-2-yl-9,9-diethyl-fluoren-2-yl-diphenylamine, in 3D data storage, 294
- Benzothiazole groups, in symmetric acceptor- $\pi$ -acceptor chromophores, 223–224
- Benzothiazolyl group, 295
- 1-[(4-Benzoylphenoxy)methyl]naphthalene (1-NpCH<sub>2</sub>-OBP), bond dissociation from  $T_n$  states in, 72–74
- 2-[(4-Benzoylphenoxy)methyl]naphthalene (2-NpCH<sub>2</sub>-OBP), 100  
bond dissociation from  $T_n$  states in, 73–74
- Biexcitonic states, 171
- Bifluorenyl model dimer, 174
- Bi mechanistic optical power limiter, 324
- Biological structures, TPEF of, 255

- Biomedical applications, of two-photon absorption, 315–321
- Bioprobes, fluorescent, 255, 256, 259–260
- Biphenyl, 63, 64, 204, 205
- Biphenyl hinge, 203, 204
- Biphenyl moiety, in symmetric acceptor- $\pi$ -acceptor chromophores, 224
- [(1,8-Bis[(4-benzoylphenoxy)methyl]naphthalene (1,8-(BPO-CH<sub>2</sub>)<sub>2</sub>Np), stepwise bond photocleavage in, 100–103
- (*E,E,E,E,E,E*)-1,13-Bis-[4-(diethylamino)phenyl]-tri-deca-1,3,5,6,8,10,12-hexaen-7-one, 274, 275, 276
- 2,7-Bis[[4-(dimethylamino)phenyl]methylene]-cycloheptanone, 277, 278
- 2,5-Bis(alkylsulfonyl)benzene unit, in symmetric donor-acceptor-donor chromophores, 226
- Bis-dioxaborine units, in symmetric acceptor- $\pi$ -acceptor chromophores, 221, 223
- Bis(diphenylamino)diphenyl hexatriene in holographic recording, 306–307 in 3D optical data storage, 303–304
- 4,4'-Bis(*N,N*-dimethylamino)benzophenone, 280
- 1,1'-2,2'-Bis(*o*-chlorophenyl)-4,4',5,5'-tetraphenyl-bisimidazole, 278
- p*-Bis(*o*-methylstyryl)benzene, as TPA measurement standard, 159
- 1,2-Bis-[(phenylthio)methylbenzene], laser photolysis of, 265, 266
- Bis(pyridylstyryl)-substituted diacetylene unit in symmetric acceptor- $\pi$ -acceptor chromophores, 221, 222
- Bit erasing, in 3D data storage, 299, 303
- B(mesityl)<sub>2</sub> group, in neutral donor- $\pi$ -acceptor chromophores, 191–192
- Bond dissociation, from T<sub>n</sub> states, 72–75
- Bond length alternation (BLA), 174–175, 182 in symmetric donor- $\pi$ -donor chromophores, 210 two-photon absorption and, 130–132
- Born–Oppenheimer approximation, two-photon excitation and, 139–140
- Boron in ionic donor- $\pi$ -acceptor chromophores, 196 in neutral donor- $\pi$ -acceptor chromophores, 191 in symmetric acceptor- $\pi$ -acceptor chromophores, 220, 223
- Bragg diffraction in optical band-gap materials, 305 in upconverted lasing, 322
- Brilliant Green, 237–238
- Bromine, 269
- Butadienes, infrared multiphoton excitation isomerization of, 260
- Butatriene mesomer, of polydiacetylene, 168–169
- t*-Butyl methacrylate, 290
- C625, medical applications of, 259–260
- Cancer photodynamic therapy for, 91–94, 318–321 TP applications in treating, 315, 316, 318
- Carbazole, 202
- Carbazole moieties, in symmetric acceptor- $\pi$ -acceptor chromophores, 216, 224–225
- Carbon nanotubes, in limiting laser optical power, 324
- Carbon–silicon (C–Si) bond, bond dissociation of, 74–75
- Carbon tetrachloride (CCl<sub>4</sub>) electron transfer to, 76–80 naphthalene T<sub>n</sub> states and, 67–70
- Cationic polymerization, two-photon initiated, 281–283
- Cavity lasing, 321
- C–C cleavage, in infrared multiphoton excitation isomerization, 260
- Cell imaging, 315, 317–318
- Centrosymmetric chromophores, three-level TPA model for, 136–139
- CEO-INDO/S procedure, 13. *See also* ZINDO/S package
- Change of state dipole moment ( $\Delta\mu_{01}$ ) in ionic donor- $\pi$ -acceptor chromophores, 194–195, 196–197 in neutral donor- $\pi$ -acceptor chromophores, 183–185, 186 in octupolar chromophores, 233, 237 in symmetric donor- $\pi$ -donor chromophores, 198–201
- Change of transmission. *See* Nonlinear transmission methods
- Channel structures, 290, 291
- Charge separated resonance structure, bond length alternation and, 132
- Charge-shift probes, 193

- Charge transfer (CT), 3  
 in octupolar chromophores, 233, 236
- Chemical reactions, multibeam-irradiation-induced, 55–57
- Chemotherapy, 255, 259–260  
 TP fluorescence in, 318
- Chinoid pattern, 196
- Chirp, 152
- Chromophore conjugation, 134
- Chromophore design, two-photon absorption optimization and, 156–255
- Chromophore-labeled dendrimers, 5–6
- Chromophores. *See also* Chromophore design; Multichromophoric systems; TP chromophores  
 bond length alternation in, 130  
 calculating distances between, 22  
 diphenylaminofluorene-based, 207  
 dipolar, 182–197  
 dithienothiophene-based, 207–208  
 future applications of, 329, 330  
 with large  $\pi$ -systems, 157–182  
 in nanoclinics, 315  
 neutral donor- $\pi$ -acceptor, 182–193  
 in nonlinear TPA measurements, 154–155  
 in numerical TPA simulations, 156  
 octupolar, 233–250  
 in photodynamic therapy, 319  
 photo-orientation of, 262  
 polymers with nonlinear absorbing, 250–255  
 “propeller shaped,” 233–250  
 quadrupolar, 136, 237  
 quantifying TPA cross section in, 144, 149–150, 152–154  
 singlet oxygen and, 266–267, 268  
 spectra and, 14–15  
 in structural isomers, 12–13  
 surface plasmon effects in, 140–143  
 symmetric acceptor- $\pi$ -acceptor, 215–225  
 symmetric donor-acceptor-donor, 225–233  
 symmetric donor- $\pi$ -donor, 197–215  
 in two-level TPA model, 134  
 two-photon absorbing, 6  
 two-photon excitation of, 116, 118, 122, 123–125  
 two-photon excited fluorescence of, 255–260  
 in two-photon initiated cationic polymerization, 281–283  
 ultrafast energy hopping between, 21  
 in upconverted lasing, 321–323
- Chromophoric backbones, 2, 3
- Chrysene (CHR)  
 direct observation of  $T_n$  states in, 85  
 energy gap law and, 70–71
- Circular grating structure, in upconverted lasing, 322, 323
- cis-trans* isomerization, of stilbene, 58–61, 64.  
*See also trans-cis* photoisomerization
- Coinitiators, 272–274, 278
- Collection efficiency ( $\eta_{col}$ ), 152
- Collimated light, in nonlinear TPA measurements, 154–155
- Complex metal ions, fluorescent chromophores and, 258
- Complex refractive index, 120
- Computer applications. *See* Three-dimensional data storage
- Concentration of propagating radicals, in radical polymerization, 279–280
- Concerted mechanism, of dissociative ELT, 80–81
- Condensed aromatic rings, TPA data for, 158
- Conjugated oligomers, two-photon absorption by, 160–182
- Conjugated polymers  
 bearing PPV pattern, 169–170  
 TPA data from, 181–182  
 two-photon absorption by, 160–182
- Conjugating rods, in symmetric donor- $\pi$ -donor chromophores, 204–205, 205–206. *See also* Connecting rods; Rod shapes
- Conjugation chain, in symmetric donor- $\pi$ -donor chromophores, 211–212
- Conjugation length, 169  
 effective, 175–177, 178  
 of phenyl-substituted ladder-type poly(*p*-phenylene), 166–167
- Connecting rods. *See also* Conjugating rods  
 in (donor- $\pi$ )<sub>3</sub>-donor chromophores, 244  
 in symmetric donor-acceptor-donor chromophores, 231  
 in symmetric donor- $\pi$ -donor chromophores, 204–205, 205–206
- Continuum beam, in nonlinear TPA measurements, 154–155
- Coulomb interactions, 11–12
- Coumarin-2 dyes, 5–6
- Coumarin-305, in fabricating optical band-gap materials, 305–306
- Coumarin-343 dye, 6

- Coupling energy ( $E_{\text{coup}}$ ), in octupolar chromophores, 236
- Coupling mechanism, of octupolar chromophores, 240
- Covalent bonding, two-photon absorption and, 173–174
- Crosslinked polymers, in 3D data storage, 301
- Crosslinked TP initiated polymerization, 279
- Crosslinking
- simultaneous two-photon initiated, 272–283
  - in two-photon initiated cationic polymerization, 281
- Crotonic esters, infrared multiphoton excitation isomerization of, 260
- Crotonitrile, infrared multiphoton excitation isomerization of, 260
- Crystallization, fabricating optical band-gap materials via, 305
- Crystals. *See also* Liquid crystals (LCs); Photonic crystals
- as optical band-gap materials, 305
  - organic molecular, 328
- Crystal Violet, 237–238
- Cubic trenches, 291
- Cyano groups, 269–270
- in neutral donor- $\pi$ -acceptor chromophores, 190–191
  - in symmetric donor-acceptor-donor chromophores, 229, 230, 231
- Cycloaddition
- of *o*-quinodimethane, 94–96
  - TP-excitation initiated, 115
  - two-photon induced, 265–266, 267
- Cyclobutenes, infrared multiphoton excitation isomerization of, 260
- Cyclophane derivatives, of symmetric donor- $\pi$ -donor chromophores, 212–214
- Damping factors, 123, 126
- Data storage
- holographic, 302–303
  - three-dimensional, 115, 293–304
- Davydov splitting, 215
- Decay times, 34
- Degenerate four-wave mixing, 145
- Degenerate TPA, 121
- $\delta$  values. *See* TPA cross section ( $\delta$ )
- Dendrimers, 2
- chromophore-labeled, 5–6
  - energy transfer among, 2–10
  - first generation, 10
  - metal-containing, 3–4
  - naphthalene, 4–5
  - phenylacetylene, 2–3
  - phenylacetylene-based, 9–10
  - polyacetylene, 3
  - polyphenylene, 2, 8–9
  - poly(propylene-imine), 7
  - in radical polymerization, 280
  - second generation, 10, 11
  - spectra of first generation, 14–15
- Dendrimer systems
- distyrylbenzene, 7
  - triarylamine, 7
- Dendritic structures, ensemble photophysics of
- rigid polyphenylene-based, 1–51
- Density functional theory (DFT), for two-photon absorption, 126–128
- Density of radicals ( $\rho$ ), in radical polymerization, 281
- Depolarization, fast, 20, 21
- Depolarization time constant, 19
- De Schryver, F. C., 1
- Dexter interactions, 3
- Dialdehyde unit, in nonlinear absorbing chromophores, 252, 253
- Dialkylamino group
- in neutral donor- $\pi$ -acceptor chromophores, 190–191
  - in octupolar chromophores, 237
- Dialkylaniline pattern, of symmetric donor-acceptor-donor chromophores, 229
- Diarylyl(methanato)boron difluorides, 223
- Diarylethene, absorbance of, 263
- Dibenzanthracene (DBA), energy gap law and, 70–71
- Dichlorobenzene (DCB), naphthalene  $T_n$  states and, 68–70
- 1,2-Dichloroethane, 58, 59
- 1,2-Dicyano-1,2-bis-(2,4,5-trimethyl-3-thienyl)ethane, two-photon induced isomerization of, 260–261
- cis*-1,2-Dicyano-1,2-bis(2,4,5-trimethyl-3-thienyl)ethane, 302
- Dicyanoanthracene, 64
- Dicyanobenzene (DCNB), naphthalene  $T_n$  states and, 68–70
- 4-(Dicyanomethylene)-2-methyl-6-(4-dimethylaminostyryl)4H-pyrene, 280

- Dicyanomethylidenedihydrothioene moiety, in symmetric acceptor- $\pi$ -acceptor chromophores, 221, 222
- 1,4-Dicyanonaphthalene (DCN), 91
- Diels–Alder cycloaddition, 12, 45
- Diethyl vinylbenzylphosphonate, in 3D data storage, 294
- Differential interference contrast (DIC) optics, in 3D data storage, 294
- Difuranonaphthalenes, singlet oxygen and, 268–269, 270
- Diisocyanate moiety, in nonlinear absorbing chromophores, 252
- Dimesitylboryl group, in symmetric acceptor- $\pi$ -acceptor chromophores, 220
- 1,2-Dimethoxybenzene, in symmetric donor-acceptor-donor chromophores, 229
- 4,4'-Dimethoxybenzophenone, excited-state ketyl radicals of, 87–88, 89
- 5,7-Dimethoxycoumarin, cycloaddition of, 265, 267
- 3-[1-(1,2-Dimethyl-1H-indol-3-yl)-ethylidene]-4-isopropylidene-dihydrofuran-2,5-dione in 3D data storage, 302
- two-photon induced isomerization of, 262–263
- 2,2'-Dimethyl-distyrylbenzene, 219
- N,N*-Dimethylformamide, 64
- Dimethylsulfoxide (DMSO), 146, 147
- Dioxaborine unit, in symmetric acceptor- $\pi$ -acceptor chromophores, 221, 223
- Dipentaerythritol pentaacrylate, in 3D optical data storage, 303–304
- Diphenylaminofluorene-based chromophores, 207
- Diphenylamino group
- in neutral donor- $\pi$ -acceptor chromophores, 189
- for symmetric donor- $\pi$ -donor chromophores, 208–209
- Diphenylanthracene, 169–170, 171
- Diphenylbutadiene, *trans-cis* photoisomerization of, 264–265
- 1,2-Diphenylcyclobutene (CB), 61
- Diphenylpolyenes
- with more than two ethylene moieties, 171
- TP excitation spectra of, 171
- trans*-Diphenylpolyenes, TPA cross section in, 157–159
- Di(*p*-methoxyphenyl)methyl chloride (An<sub>2</sub>CHCl), three-color three-laser photochemistry of, 99–100
- Dipolar chromophores, 182–197
- OPA and TPA of, 186
- Dipolar chromophore structure
- bond length alternation and, 132
- in two-level TPA model, 133
- Dipole, transition, 178, 180, 186
- Dipole moment (*M*), 123. *See also* Transition dipole moment (*M<sub>nm</sub>*)
- induced, 120
- of ionic donor- $\pi$ -acceptor compounds, 193
- transition, 175–177, 180–181, 190
- Dipole moment changes ( $\Delta\mu$ ), 123
- in two-level TPA model, 133
- Direct excitation, of retinal, 265
- Disperse Red, 251
- Distyrylbenzene, amino-substituted, 290
- Distyrylbenzene dendrimer systems, 7
- Distyrylbenzene derivatives
- in singlet oxygen generation, 269–270
- as symmetric donor-acceptor-donor chromophores, 229–230
- as symmetric donor- $\pi$ -donor chromophores, 202–203
- TPA data for, 159
- Distyrylbenzene moiety
- in nonlinear absorbing chromophores, 252, 253, 254
- in symmetric acceptor- $\pi$ -acceptor chromophores, 220
- in symmetric donor- $\pi$ -donor chromophores, 213
- in two-photon initiated cationic polymerization, 282
- Dithienothiophene-based chromophores, 207–208
- DNA cleavage method, 98
- DNA damaging, two-color two-laser, 91–94
- Donor–acceptor distances, 21
- Donor-acceptor-donor (**D-A-D**) compounds, symmetric, 225–233
- (Donor- $\pi$ )<sub>3</sub>-donor chromophores, 243–244, 245
- (Donor- $\pi$ )<sub>3</sub>- $\pi$  chromophores, 242–243
- (Donor- $\pi$ -acceptor)<sub>3</sub>-donor chromophores, 247, 249
- Donor- $\pi$ -acceptor (**D- $\pi$ -A**) compounds
- ionic, 193–197
- neutral, 182–193
- Donor- $\pi$ -donor (**D- $\pi$ -D**) compounds
- symmetric, 197–215
- in three-dimensional microfabrication, 286
- as TP initiators, 274

- Doracure photoinitiators, 272, 273
- Double-conjugated-segment molecules, 225
- Doublet states, excited, 55
- Driving force ( $-\Delta G_{\text{ELT}}$ ), in electron transfer, 77–78
- Drug delivery screening, 315
- Effective conjugation length ( $n_e$ ), 175–177, 178
- Effective emitter concentration, 178
- Effective molecular weight, 177–178
- Efficiency ( $E$ ), 20
- Electron affinity, of symmetric donor-acceptor-donor chromophores, 230
- Electron beams, from linear accelerator, 54
- Electron density, in symmetric donor-acceptor-donor chromophores, 229, 230
- Electron ejection, from NDI, 93
- Electronic coupling, 124, 169
- in polyacetylene dendrimers, 3
- singlet oxygen and, 266–267
- Electronic excitation transfer, 10–14
- Electronic state parameters, 175
- Electron-rich moieties, in symmetric donor- $\pi$ -donor chromophores, 208–209
- Electrons, in surface plasmon effects, 141
- Electron transfer (ELT), 68. *See also* ELT entries;
- Marcus ELT theory; Photoinduced ELT;
- Selective ELT quenching
- in dendritic macromolecules, 2–10
- photoinduced, 272–274
- secondary, 58
- “sticky” dissociative, 79
- in  $\text{TiO}_2$  photocatalytic reactions, 96–98
- from  $T_n$  states, 75–81, 82
- in two-photon initiated cationic polymerization, 281–282
- Electron-withdrawing groups, 219, 225–226
- in octupolar chromophores, 246–247, 248
- ELT pathways, of excited-state benzophenone ketyl radicals, 88–89. *See also* Electron transfer (ELT)
- ELT quenching, 68
- Emission maxima, 295–296
- Emission spectra, steady state, 14–15
- Emission wavelength, chromophore number and, 18
- Energy exchange rate, 121
- Energy gap law,  $T_n$  states and, 70–71
- Energy hopping, 23–24
- singlet–singlet annihilation versus, 44–45, 46
- Energy relaxation processes, 285
- Energy transfer (ENT) process, 66, 68–70, 75.
- See also* ENT quenching
- in dendritic macromolecules, 2–10
- Ensemble photophysics, of rigid polyphenylene-based dendritic structures, 1–51
- ENT quenching, 68–70, 91. *See also* Energy transfer (ENT) process
- Epoxy resins
- in holographic recording, 307
- in optical circuits, 309
- TP initiated polymerization of, 286
- Erasing, in 3D data storage, 299, 303
- 1-Ethylanthralene, electron transfer from  $T_n$  states in, 77, 79
- Ethynyl-anthryl moiety, in octupolar chromophores, 239
- Ethynyl-phenylene-ethynyl connector, in symmetric donor- $\pi$ -donor chromophores, 211–212
- “Excimer-like” entity, 18
- Excitation, multistep, 54. *See also* TPE entries;
- TP excitation maxima; TP excited states;
- Two-photon excitation (TPE)
- Excitation energy, 123, 126, 127, 169
- in multichromophoric compounds, 25–37
- Excitation intensity, TPEF and, 165, 166
- Excited ketyl radicals, studied via two-color two-flash photolysis, 85–91
- Excited radical anions, pulse radiolysis of, 64–65
- Excited radical cations
- fluorescence from, 61–64
- pulse radiolysis of, 58–61
- studied via two-color two-flash photolysis, 91
- Excited state absorption (ESA), 38
- in two-photon absorption, 117, 123–124
- Excited state processes, in multichromophoric systems, 2–10
- Excited states, 55. *See also* TP excited states
- Exciton coupling energy, in symmetric donor- $\pi$ -donor chromophores, 213
- Excitonic states, 171
- Exciton model of linear aggregates, 175
- Extinction coefficient ( $\epsilon$ ), 134
- in ionic donor- $\pi$ -acceptor chromophores, 194–195
- in neutral donor- $\pi$ -acceptor chromophores, 183–185
- of nonlinear absorbing chromophores, 253
- in octupolar chromophores, 234–235

- Extinction coefficient ( $\epsilon$ ) (*Continued*)  
 in symmetric acceptor- $\pi$ -acceptor chromophores, 217–218  
 in symmetric donor-acceptor-donor chromophores, 227–228  
 in symmetric donor- $\pi$ -donor chromophores, 198–201, 202
- Fabrication techniques, for optical band-gap materials, 305–306, 307
- Face-centered cubic (fcc) crystals, in optical band-gap materials, 305
- Fast depolarization, 20, 21
- Femtosecond fluorescence upconversion measurements, 25–37, 44
- Femtosecond (fs) lasers, 54, 98  
 multiphoton excitation via, 113–114  
 quantifying TPA cross section via, 144, 147–148  
 singlet oxygen production and, 268  
 for 3D optical memory systems, 294  
 TPA data taken with, 156–157  
 waveguides and, 309
- Femtosecond transient absorption measurements of, 37–45  
 by methyl-substituted ladder-type poly(*p*-phenylene), 163
- Fermi Golden Rule, 10–11
- Field intensity/strength, with two-photon absorption, 120–121
- First generation dendrimers, 10
- Fluorene(s), 174, 175, 204, 205–206  
 donor-acceptor substituted, 187–188  
 TPA spectra of, 176
- Fluorene moiety, 240–241  
 in symmetric acceptor- $\pi$ -acceptor chromophores, 224
- Fluorescein, 255  
 as TPA measurement standard, 152–154, 155
- Fluorescence, 123, 124. *See also* Two-photon excited fluorescence (TPEF)  
 of excited di(*p*-methoxyphenyl)methyl radical, 99–100  
 of excited radical cations, 61–64  
 by ionic donor- $\pi$ -acceptor compounds, 193  
 of liquid crystals, 303  
 in neutral donor- $\pi$ -acceptor chromophores, 188–189  
 of octupolar chromophores, 240–241  
 quantifying TPA cross section via, 144  
 solvent-dependent, 188–189  
 surface-plasmon-enhanced, 143  
 in symmetric donor- $\pi$ -donor chromophores, 205, 206  
 in 3D data storage, 298–299, 300
- Fluorescence cross-section action, in fluorescence imaging applications, 257
- Fluorescence decay times, 177  
*m*-C1P<sub>x</sub>, 17
- Fluorescence excitation spectra, 214
- Fluorescence imaging, 311–315  
 in 3D data storage, 295–296
- Fluorescence imaging applications, fluorescence cross-section action in, 257
- Fluorescence quantum yield ( $\Phi_f$ ), 177  
 from ionic donor- $\pi$ -acceptor chromophores, 194–195, 196–197  
 from neutral donor- $\pi$ -acceptor chromophores, 183–185, 186  
 from nonlinear absorbing chromophores, 253  
 from octupolar chromophores, 234–235  
 from symmetric acceptor- $\pi$ -acceptor chromophores, 217–218  
 from symmetric donor-acceptor-donor chromophores, 227–228  
 from symmetric donor- $\pi$ -donor chromophores, 198–201
- Fluorescence resonance energy transfer (FRET), 271
- Fluorescence spectra, 172  
 of excited radical cations, 91  
 of excited-state azaxanthone ketyl radicals, 88–89  
 of excited-state xanthone ketyl radicals, 88, 90
- femtosecond fluorescence upconversion measurements and, 25–37  
 femtosecond transient absorption measurements and, 39–42  
 magic angle polarization and, 15–18  
 by methyl-substituted ladder-type poly(*p*-phenylene), 165  
 steady state, 14–15  
 in time-resolved fluorescence polarization measurements, 18–25
- Fluorescent bioprobes, 255, 256  
 medical applications of, 259–260
- Fluorescent dye, in 3D data storage, 294–295
- Fluorinated moieties, 186–187
- Fluorinated phenyl rings, 186–187
- Fluorinated polyenes, 134

- Forbidden transitions, 168  
Förster interactions, 3  
Förster theory, 10–11, 14, 20, 21–22, 23, 24, 37, 46, 47  
  excited radical cations in, 91  
Fragmentation, in infrared multiphoton excitation isomerization, 260  
Franck–Condon factors, 140  
Fresnel reflectance, 147  
Fujitsuka, Mamoru, 53  
Fulgide, in holographic data storage, 302
- Gaussian beam propagation  
  limiting laser optical power and, 326, 328  
  in 3D data storage, 300–301  
Gaussian pulse, 145–146, 147, 150, 284  
Generation number, femtosecond transient absorption measurements and, 44  
Gold nanoparticles (AuNPs), 89–91  
  surface plasmon effects and, 142–143  
Gold tips, in three-dimensional microfabrication, 289–290  
Göppert-Mayer, Maria, 113, 118  
Göppert-Mayer (GM), 118  
Green band, 139  
Group delay dispersion (GDD), 152  
Guanine (G), in two-color two-laser DNA damaging studies, 93–94
- Hammett constants, 71–72, 188, 191–192, 216  
Hartree–Fock geometries, 128  
Heisenberg relationship, 118  
Heptathiophene dye, 5–6  
Herzberg–Teller expansion, 139, 254  
Heteroaromatics, TPA data for, 159  
Heterocyclic moieties, in neutral donor- $\pi$ -acceptor chromophores, 187–188  
Heterodyned Kerr effect, 145  
Hexadienes, infrared multiphoton excitation isomerization of, 260  
Higher triplet excited ( $T_n$ ) states, energy transfer from, 66–70  
Highest occupied molecular orbital (HOMO), 182  
  symmetry and, 129  
Hofkens, J., 1  
Hole transfer, 63  
Holographic data storage systems, 302–303  
Holographic recording, 302–303  
  optical band-gap materials for, 306–307  
Holography, 115
- Hopping, rate constant of, 20–21, 22. *See also* Energy hopping  
Hybrid organic/inorganic ordered nanostructures, in holographic recording, 307  
Hydrogen ions ( $H^+$ ), in two-photon initiated cationic polymerization, 282  
Hydrogen loss, in infrared multiphoton excitation isomerization, 260  
Hyperbolic secant. *See*  $\text{sech}^2$ -shaped pulse  
Hyperpolarizability ( $\gamma$ ), 160  
  bond length alternation and, 130–132  
  nonlinear, 190–191, 250, 251  
  second, 182  
Hyperthermal effect, 316  
Hypsochromic shift of TP excitation maximum, by symmetric donor- $\pi$ -donor chromophores, 210
- Image potential states, surface plasmon effects and, 141  
Index of refraction  
  in 3D data storage, 294, 298  
  in waveguide materials, 310  
Indirect population, 124  
INDO-MRDCI/SOS method, 238. *See also* ZINDO/S package  
Induced dipole moment, 120  
Induced polarization, 120  
Infrared multiphoton excitation, isomerization via, 260. *See also* Near infrared (NIR) light  
Initiation, of radical polymerization, 276  
Intensity of light, in radical polymerization, 280–281  
Interchromophoric distances, 22  
Interdisciplinary cooperation, in TPA research, 329  
Intermediates, three-laser control of, 99–100  
Intermolecular charge transfer, TP-excitation initiated, 115  
Intermolecular ELT, 75  
Internal conversion (IC), 116  
Intersystem crossing (ISC), 66, 116  
  quantum yield for, 241  
  singlet oxygen and, 266  
Intramolecular charge transfer (ICT)  
  in neutral donor- $\pi$ -acceptor chromophores, 182, 186  
  in octupolar chromophores, 240  
  of symmetric donor-acceptor-donor chromophores, 230

- Intramolecular ELT, 75
- Ionic donor- $\pi$ -acceptor (**D- $\pi$ -A**) compounds, 193–197
- Irgacure photoinitiators, 272, 273
- Iron oxide ( $\text{Fe}_2\text{O}_3$ ), in nanoclinics, 315, 316
- Isomerization. *See also* Photoisomerization  
in 3D data storage, 294, 301  
TP-excitation initiated, 115  
two-photon-induced, 260–265
- J aggregates, 31
- Keggin-type polyoxometalates (POMs), in  $\text{TiO}_2$   
photocatalytic reactions, 96–98
- Ketyl radicals, studied via two-color two-flash  
photolysis, 85–91
- Kinetic analysis, of TP induced radical  
polymerization, 279
- Lactone, rhodamine B, 296
- Ladder-type polymers, TPA cross section for,  
163
- Lambert–Beer law, 117, 177
- Laser calorimetry, 144, 148
- Laser flash photolysis. *See also* Two-color  
two-laser flash photolysis  
combined with pulse radiolysis, 57, 58  
cycloaddition and, 265, 266, 267  
of stilbene, 58–61
- Laser pulse, ultrashort, 298
- Lasers, 54  
in efficient TPE, 278  
femtosecond, 54, 98, 113–114  
in fluorescence imaging, 313–314  
in radical polymerization, 280–281  
limiting optical power of, 323–328  
nanoclinics and, 317–318  
nanosecond, 57, 85  
picosecond, 81, 85  
power dependence of, 165  
pulsed, 65–66  
studying  $T_n$  states via, 66  
in three-dimensional microfabrication, 284,  
285, 286, 287, 288, 291, 293  
TPEF with, 150–154  
upconverting, 321–323
- Laser scanning confocal microscope (LSCM),  
311–312
- Lattice constant, in optical band-gap materials,  
305
- Lattice structures, in TPA microfabrication, 283
- Layer-by-layer structure formation, fabricating  
optical band-gap materials via, 305–306, 307
- Lewis acids, in two-photon initiated cationic  
polymerization, 282
- Light. *See also* Near infrared (NIR) light;  
Ultraviolet entries  
in near-field fluorescence imaging, 312, 313  
in nonlinear TPA measurements, 154–155  
optical band-gap materials and, 305  
optical power limiters and, 324  
in radical polymerization, 280–281  
in 3D data storage, 295  
in 3D microfabrication, 284, 285, 286, 287,  
288  
two-photon absorption and, 118, 120–121  
in waveguide materials, 308–309
- Light intensity isophotes, laser beam, 297
- Limiting anisotropy, 18, 19–20
- Linear absorption, in symmetric donor- $\pi$ -donor  
chromophores, 205
- Linear absorption coefficient, 144
- Linear accelerator, electron beam from, 54
- Linear coupling model, vibrational contributions  
in, 140
- Linker groups, 319
- Lippert–Mataga equations, 133, 191, 240
- Liquid crystals (LCs), in 3D optical data storage,  
303–304
- Lor, M., 1
- Lorenz field correction, 122
- Lowest unoccupied molecular orbital (LUMO),  
182
- Magic angle polarization, time-resolved  
fluorescence measurements under, 15–18
- Magnesium ions, fluorescent chromophores and,  
258, 259
- Magnetic fields, nanoclinics and, 316
- Magnetic iron oxide ( $\text{Fe}_2\text{O}_3$ ), in nanoclinics,  
315, 316
- “Magnetic nanoclinic,” 114–115
- Magnetocytolysis, 316, 318
- Majima, Tetsuro, 53
- Marcus ELT theory, 80–81
- Materials, two-photon-absorbing, 144–155  
*m*-C1P<sub>1</sub>, femtosecond transient absorption  
measurements of, 37–39, 42  
*m*-C1P<sub>3</sub>, femtosecond transient absorption  
measurements of, 39–42

- m*-C1P<sub>x</sub>. *See also* Meta-substituted first generation carbon core dendrimers  
femtosecond fluorescence upconversion  
measurements of, 27, 28, 33  
fit parameters of, 19  
fluorescence decay times of, 17
- MCF-7 cells, fluorescence imaging of, 317–318
- Mechanism of action, 255
- Mechanistic TPA models, 132–143
- Medical applications  
of fluorescent bioprobes, 259–260  
of two-photon absorption, 315–321
- Memory, optical, 293–294
- 3-Mercapto-4-methyl-4H-1,2,4-triazole, 278
- Mesomers, of polydiacetylene, 168–169
- Metal-containing dendrimers, 3–4
- Metal ions, fluorescent chromophores and, 257–258
- Metal particles, surface plasmon effects in, 141–142
- Meta-substituted compounds, spectra of, 14–15
- Meta-substituted first generation carbon core dendrimers. *See also m*-C1P<sub>x</sub>  
para-substituted carbon core dendrimers versus, 18  
single-photon timing measurements of, 17–18  
time-resolved fluorescence polarization measurements of, 18–19
- Methacrylates, TP initiated polymerization of, 274, 275, 286
- p*-Methoxymethylbenzophenone (BPCH<sub>2</sub>OCH<sub>3</sub>), bond dissociation from T<sub>n</sub> states in, 74
- 1-Methoxynaphthalene, electron transfer from T<sub>n</sub> states in, 76–77
- Methyl groups, in nonlinear absorbing chromophores, 252, 253
- Methylmethacrylate, in 3D data storage, 294, 302
- 1-Methylnaphthalene, electron transfer from T<sub>n</sub> states in, 77
- Methyl-substituted ladder-type poly(*p*-phenylene), 163–165
- Microbull, 292, 293
- Microcapsules, 292, 293
- Microchain, 292, 293
- Microchannel structures, 290, 291
- Microfabrication  
three-dimensional, 284–293  
three-dimensional lithographic, 285–293  
two-photon initiated crosslinking polymerization in, 272, 283
- Microgearwheel, 292, 293
- Microscopes, in three-dimensional microfabrication, 284, 285
- Microscopy, fluorescence imaging and, 311
- Microtweezers, 289
- Mirror-scanner system, in three-dimensional microfabrication, 286–287
- Mode-locked Ti:sapphire laser, 150, 151, 152  
in fluorescence imaging, 313–314  
in nonlinear TPA measurements, 154–155  
in three-dimensional microfabrication, 286–288
- Mode-locked Nd:YAG laser, 304
- Molar extinction coefficient, 134
- Molecular engineering, 128  
of TP fluorophores, 204–205
- Molecular memory, 56
- Molecular orbital (MO) methods, 125
- Molecular orbitals, symmetry and, 129
- Molecular parameters, with two-photon absorption, 119–120, 123, 125
- Molecular switching, 57
- Molecular weight  
average, 178  
effective, 177–178
- Monomer units, angle dependence between, 180
- Monomode waveguides, 309
- Morse potential curve, 78
- Multibeam irradiation  
applications of, 55–57  
photochemistry of short-lived species using, 53–109
- Multichromophoric systems. *See also* Chromophores  
excited state processes in, 2–10  
femtosecond fluorescence upconversion measurements of, 25–37, 46  
femtosecond transient absorption measurements of, 37–45  
time-resolved fluorescence polarization measurements of, 20–25
- Multiphoton absorption  
by aromatic hydrocarbons, 157  
surface plasmon effects and, 142
- Multiphoton excitation  
isomerization via, 260  
in photochemistry, 113
- Multiple phenyl rings, TPA data for, 158

- Multireference double excitation configuration interaction (MRDCI) scheme, 125, 128  
with three-level TPA model, 136, 138
- Multistep excitation, 54
- Multivalence-bond state approach, for two-photon absorption, 128
- Nanoclinics, 315–318
- Nanoengineered particles, surface plasmon effects in, 141–142
- Nanofabrication  
three-dimensional, 284–293  
two-photon initiated crosslinking polymerization in, 272, 283
- Nanometer-sized metal particles, surface plasmon effects in, 141–142
- Nanoparticles, in fabricating optical band-gap materials, 305. *See also* Nanoclinics
- Nanoscope IIIA controller, 314
- Nanosecond (ns) lasers, 57  
direct measurement of  $T_n$  state lifetimes via, 85  
ketyl radicals in the excited state studied via, 85  
singlet oxygen production and, 268  
studying  $T_n$  states via, 66  
TPA data taken with, 156–157
- Nanosecond optical parametric oscillator, 174
- Naphthacenepyrindone, in 3D data storage, 302
- Naphthalidiimide (NDI), in two-color two-laser DNA damaging studies, 92–94
- Naphthalene (Np)  
energy gap law and, 70–71  
 $T_n$  states of, 66–70
- Naphthalene dendrimers, 4–5
- Naphthalene derivatives (NpD)  
electron transfer from  $T_n$  states in, 75–81, 82  
substituent effects on  $T_n$  state lifetimes of, 71–72
- Naphthylmethylhalides, 72
- Naphthylmethyl radical, bond dissociation from  $T_n$  states in, 72–74
- NDI-conjugated oligodeoxynucleotide (NDI-ODN), in two-color two-laser DNA damaging studies, 92–94
- Nd:YAG laser, 304  
in three-dimensional microfabrication, 293
- Nd:YAG pumped optical parametric oscillator, singlet oxygen production and, 268
- Near-field fluorescence imaging, 312–313
- Near-field optical microscopy (NFOM), 283
- Near-field scanning optical microscopy (NSOM), in three-dimensional microfabrication, 285, 289
- Near infrared (NIR) light, in photodynamic therapy, 318–319. *See also* Infrared multiphoton excitation; NIR pulsed lasers
- Neutral donor- $\pi$ -acceptor (D- $\pi$ -A) compounds, 182–193
- Neutral resonance structure, bond length alternation and, 132
- NIR pulsed lasers, nanoclinics and, 317. *See also* Near infrared (NIR) light
- o*-Nitrobenzaldehyde, TP photochemical rearrangement of, 297
- Nitrobenzene, in neutral donor- $\pi$ -acceptor chromophores, 190
- Nitrogen  
in octupolar chromophores, 243–244, 245  
in 3D data storage materials, 295
- Nonlinear absorbing chromophores, polymers with, 250–255
- Nonlinear absorption, 190–191
- Nonlinear absorption coefficient ( $\beta$ ), 144, 146, 147, 181  
limiting laser optical power and, 326
- Nonlinear absorption data, of conjugated polymers and oligomers, 160–163, 181–182
- Nonlinear hyperpolarizability, 190–191  
in nonlinear absorbing chromophores, 250, 251
- Nonlinear optical parameters, 182  
two-photon absorption and, 116–122
- Nonlinear optical responses, surface plasmon effects and, 142
- Nonlinear refraction, in limiting laser optical power, 324
- Nonlinear refractive index, 181
- Nonlinear scattering, in limiting laser optical power, 324
- Nonlinear TPA  
in absolute evaluation of TPA materials, 154–155  
experimental techniques in, 144–148
- Nonlinear transmission methods, 188  
for quantifying TPA cross section, 144, 145–147
- Nonprotonated species, in 3D data storage, 295–296
- Normalized TP excitation spectra, 170–171

- Normalized transmittance ( $T_N$ ), 146–147
- Numerical TPA simulations, 156
- Octupolar (acceptor- $\pi$ )<sub>3</sub>-donor chromophores, 246–247, 248
- Octupolar chromophores, with large two-photon absorptivity, 233–250
- Octupolar (donor- $\pi$ -acceptor)<sub>3</sub>-donor chromophores, 247, 249
- Oligomer length ( $OL$ )  
TPA cross section and, 173
- Oligomers, 177  
in TP fluorophore molecular engineering, 204–205  
two-photon absorption by conjugated, 160–182  
urethane acrylate, 309
- Oligo(*p*-phenylenevinylene)s (OPVn), direct observation of  $T_n$  states in, 84
- Oligothiophenes, 173  
direct observation of  $T_n$  states in, 81–84
- One-photon absorption (OPA), 134. *See also* OPA entries  
fluorescence imaging and, 311  
with ionic donor- $\pi$ -acceptor compounds, 193  
in nonlinear absorbing chromophores, 252, 253  
by phenyl-substituted ladder-type poly(*p*-phenylene), 165–167  
Strickler–Berg relationship and, 177  
in three-dimensional microfabrication, 286  
two-photon absorption and, 117–118  
in two-photon initiated cationic polymerization, 282
- One-photon (OP) excitation, 113, 115, 134–135. *See also* OP excitation entries  
of conjugated polymers, 163  
of rhodamine B, 256–257  
symmetry and, 128–129  
in three-level TPA model, 138–139
- One-photon resonance, in symmetric donor- $\pi$ -donor chromophores, 211
- Onium compounds, as cointiators, 272
- Onsager TPA model, 133, 188–189
- OPA cross section, in octupolar chromophores, 246
- OPA maxima  
in neutral donor- $\pi$ -acceptor chromophores, 183–185  
in symmetric acceptor- $\pi$ -acceptor chromophores, 217–218, 223  
in symmetric donor-acceptor-donor chromophores, 227–228, 232  
in symmetric donor- $\pi$ -donor chromophores, 198–201, 202, 212  
in TP initiators, 275
- OPA spectra, 181  
of conjugated polymers, 170  
of nonlinear absorbing chromophores, 254  
of octupolar chromophores, 239, 241  
of symmetric donor-acceptor-donor chromophores, 232  
of symmetric donor- $\pi$ -donor chromophores, 210
- OP broadband UV excitation, cycloaddition via, 265. *See also* One-photon entries
- OP excitation maxima  
in ionic donor- $\pi$ -acceptor chromophores, 194–195  
of nonlinear absorbing chromophores, 253  
in octupolar chromophores, 234–235, 236
- OP excitation spectra, 181
- OP excited fluorescence (OPEF), 151, 152. *See also* Single-photon excited fluorescence (SPEF) spectra
- OP excited states  
energetic ordering of, 157  
of symmetric donor- $\pi$ -donor chromophores, 212
- OP initiated polymerization, 279  
in three-dimensional microfabrication, 284–285
- Optical band-gap materials, 305–308. *See also* Photonic band-gap materials
- Optical circuits, 309
- Optical elements, in absolute evaluation of TPA materials, 149
- Optical feedback, in upconverted lasing, 322
- Optical memory, three-dimensional, 293–294, 304
- Optical parameters, nonlinear, 116–122, 182
- Optical parametric oscillator (OPO), singlet oxygen production and, 268
- Optical power limiters, 114, 323–328
- Optical probes, in nanoclinics, 315
- Optical susceptibility, 181–182
- Organic boron compounds, 191
- Organic chromophores, surface plasmon effects and, 142–143
- Organic compounds, UV absorption spectra of, 325

- Organic molecular crystals, limiting laser optical power and, 328
- Orthogonal addressing, in 3D data storage, 300–301
- Oxidation potential, of TP initiators, 275
- Oxygen. *See* Singlet oxygen
- “Page-by-page” writing, in 3D data storage, 300–301
- Paracyclophane, in symmetric donor- $\pi$ -donor chromophores, 214–215
- Para-substituted carbon core dendrimers. *See also p-C* entries
- meta-substituted carbon core dendrimers versus, 18
  - single-photon timing measurements of, 15–17
- Para-substituted compounds, spectra of, 14–15
- Para-substituted first generation carbon core dendrimers, time-resolved fluorescence polarization measurements of, 19–25
- Parity, in two-photon absorption, 128–129
- Parity forbidden transitions, 113
- p*-C1P<sub>1</sub>
- femtosecond fluorescence upconversion measurements of, 25–30, 31–33
  - femtosecond transient absorption measurements of, 37–39, 40
  - molecular structure of, 12–13
- p*-C1P<sub>2</sub>, molecular structure of, 12–13
- p*-C1P<sub>3</sub>, femtosecond transient absorption measurements of, 39–42
- p*-C1P<sub>4</sub>
- femtosecond fluorescence upconversion measurements of, 26–37
  - molecular structure of, 12–13
  - time-resolved fluorescence decays of, 16
- p*-C1P<sub>4</sub>A, molecular structure of, 13–14
- p*-C1P<sub>4</sub>B, molecular structure of, 13–14
- p*-C1P<sub>x</sub>, femtosecond fluorescence upconversion measurements of, 27, 28, 29, 30
- p*-C2P<sub>1</sub>, femtosecond transient absorption measurements of, 42–45
- p*-C2P<sub>4</sub>
- femtosecond transient absorption measurements of, 42–45
  - isomeric structures of, 25
  - molecular structure of, 10, 11
- p*-C2P<sub>x</sub>
- energy hopping in, 23–24
  - femtosecond fluorescence upconversion measurements of, 26, 27, 34–37
  - fit parameters of, 17
- Pentadienes, infrared multiphoton excitation isomerization of, 260
- Pentafluorobenzene moiety, 186
- in ionic donor- $\pi$ -acceptor chromophores, 196
- Perturbation theory, two-photon absorption and, 118–119
- Peryleneimide (PI), 29, 36–37, 42, 43
- Peryleneimide chromophores, 8–9, 10, 14, 45–47
- Perylene units, 7–8
- Phenazine, 64
- 2-Phenoxyethyl acrylate, polymerization of, 278
- p*-Phenoxyethylbenzophenone (BPCh<sub>2</sub>Oph), bond dissociation from T<sub>n</sub> states in, 74
- Phenylacetylene-based dendrimers, 9–10
- Phenylacetylene dendrimers, energy transfer in, 2–3
- Phenylene-ethynylene oligomers, 204–205
- Phenylene-vinylene hinge, in symmetric donor- $\pi$ -donor chromophores, 208–209
- Phenylene-vinylene oligomers, 204–205
- Phenyl moiety
- in neutral donor- $\pi$ -acceptor chromophores, 192
  - in octupolar chromophores, 237
- Phenyl rings
- fluorinated, 186–187
  - in symmetric acceptor- $\pi$ -acceptor chromophores, 220
  - in symmetric donor- $\pi$ -donor chromophores, 213
  - TPA data for, 158, 159
- Phenyl-substituted ladder-type poly(*p*-phenylene), 165–167
- Phosphorescence, of singlet oxygen, 268
- Photobiology, singlet oxygen studies in, 266
- Photobleaching, 142, 148
- Photocatalytic reactions, two-color laser control of, 96–98
- Photochemical reactions, TP-excitation initiated, 115
- Photochemistry
- of di(*p*-methoxyphenyl)methyl chloride, 99–100
  - of short-lived species, 53–109
  - two-photon absorption in, 113–115
- Photochromic matrices, in 3D data storage, 301–302
- Photocrosslinked polymers, in 3D data storage, 301

- Photocycloaddition, TP-excitation initiated, 115
- Photodynamic therapy (PDT), 56, 91–94, 315, 318–321
- multiphoton excitation in, 114, 115
  - singlet oxygen in, 268
- Photoinduced acid generator, 295
- Photoinduced electron transfer (ELT), 75–81, 82, 272–274
- DNA damage and, 92
- Photoinduced processes, 55–56
- Photoinitiators, 272, 273. *See also* TP initiators
- Photoisomerization, of ionic donor- $\pi$ -acceptor chromophores, 196. *See also* Isomerization; *trans-cis* photoisomerization
- Photoluminescence intensity, 166
- Photoluminescence spectra, by methyl-substituted ladder-type poly(*p*-phenylene), 165
- Photolysis, cycloaddition and, 265, 266, 267. *See also* Pulse radiolysis–laser flash photolysis; Three-color three-laser flash photolysis; Two-color two-laser flash photolysis
- Photon flux (*F*)
- in radical polymerization, 281
  - temporal fluctuations in, 149–150
- “Photonic atoms,” 305
- Photonic band-gap materials, 283. *See also*
- Optical band-gap materials
  - three-dimensional microfabrication of, 288–289, 290, 293
- Photonic crystals, 283
- as optical band-gap materials, 305, 306, 308
- Photonic properties, of symmetric acceptor- $\pi$ -acceptor chromophores, 219
- Photon number noise, in nonlinear absorbing chromophores, 250
- Photo-orientation, of chromophores, 262
- Photophysics, of rigid polyphenylene-based dendritic structures, 1–51
- Photopolymerization
- in 3D data storage, 294
  - TPA processes in, 283
  - two-photon initiated cationic, 281–283
- Photorefractive effect, in 3D data storage, 299–300
- Photosynthesis, 97
- Picosecond lasers, 81
- direct measurement of  $T_n$  state lifetimes via, 85
  - ketyl radicals in the excited state studied via, 85
- Picosecond relaxation process, 39
- $\pi$ -electron density
- in polyenes, 130–132
  - surface plasmon effects and, 141
  - in two-level TPA model, 133
- Pigment–protein complexes, 31
- Pinhole aperture, 312
- $\pi$ -structures, 157
- $\pi$ -systems ( $\Pi$ -systems)
- chromophores with large, 157–182
  - polarizable, 215, 216, 219
- Planarity, 186
- Point dipole–point dipole approximation, for symmetric donor- $\pi$ -donor chromophores, 213
- Polarity, solvent, 206
- Polarizability, second-order nonlinear, 120
- Polarizable  $\Pi$ -systems, 215, 216, 219
- Polarization, magic angle, 15–18
- Polarization ratio, two-photon excited fluorescence and, 148
- Poly[1,4-phenylene-1,2-di(phenoxyphenyl)vinylene], in waveguide materials, 309–310
- Poly[1,4-phenylene-1,2-diphenylvinylene-*co*-2,7-fluorenylene-1,2-di-phenylvinylene], in waveguide materials, 309–310
- Poly[1,6-bis(3,6-dihexadecyl-*N*-carbazolyl)2,4-hexadiyne], TPA cross section of, 167–168
- Poly(2,2'-bipyridine-5,5'-diethynylene[2,5-di(2-ethylhexyl)oxy-1,4-phenylene]ethynylene), 178
- Polyacetylene dendrimers, electronic coupling in, 3
- Polyaddition, of nonlinear absorbing chromophores, 252
- Poly(arylenevinylene)s, 177
- Polychlorobenzenes, electron transfer to, 80–81, 82
- Poly(diacetylenes), 171
- mesomers of, 168–169
- Polyene chain, 203
- Polyenes, 197
- bond length alternation in, 130
  - donor–acceptor substituted, 186–187
  - fluorinated, 134
  - TPA data for, 159

- Poly(ethylene terephthalate), in 3D data storage, 302
- Polyfluorene(s), 175  
TP excitation of, 167
- Poly-HEMA (polyhydroxyethylmethacrylate) rod, as lasing material, 322
- Polymer dispersed liquid crystals, in 3D optical data storage, 303–304
- Polymerization  
simultaneous two-photon initiated, 272–283  
TPA processes in, 283  
two-photon initiated cationic, 281–283
- Polymer photochromic matrices, in 3D data storage, 301–302
- Polymers  
as lasing materials, 321–322  
with nonlinear absorbing chromophores, 250–255  
in 3D data storage, 301–304  
two-photon absorption by conjugated, 160–182
- Polymer science, singlet oxygen studies in, 266
- Poly(methylmethacrylate)  
in 3D data storage, 294, 302  
in waveguide materials, 309
- Polyoxometalates (POMs), in TiO<sub>2</sub> photocatalytic reactions, 96–98
- Polyphenylene-based dendritic structures, ensemble photophysics of, 1–51
- Polyphenylene dendrimers, 8–9  
as chromophoric backbones, 2–10
- Poly(phenylene vinylene) (PPV), 163, 169–170
- Polypyridine ligands, 3–4
- Poly(*p*-phenylene)  
methyl-substituted ladder-type, 163–165  
phenyl-substituted ladder-type, 165–167
- Poly(*p*-phenylene acetylene), 163
- Poly(propylene-imine) dendrimers, 7
- Poly(styrene-*co*-acrylonitrile), 289, 290  
in fabricating optical band-gap materials, 306, 308
- Polystyrene nanoparticles, in fabricating optical band-gap materials, 305
- Poly(thiophene), 163
- Poly(vinyl alcohol) (PVA) film, 89
- Porphyrin arrays, 7
- Porphyrins, in photodynamic therapy, 319
- Porphyrin unit, singlet oxygen generation and, 271
- Positive-resist structures, 291
- Positive tone resins, in three-dimensional microfabrication, 290
- Positive TP microlithography, 290
- Primary initiator particle density ( $\rho_0$ ), in radical polymerization, 281
- Prismatic cavities, 291–293
- Prismatic trenches, 291
- Product analysis, rate constant determination from, 94–96
- Propagation, of radical polymerization, 276
- “Propeller shaped” chromophores, with large two-photon absorptivity, 233–250
- Proteins, TPEF and, 255
- Protonation, in 3D data storage, 295–296
- Proton diffusion, in three-dimensional microfabrication, 291
- Pulse collision addressing, in 3D data storage, 300–301
- Pulsed femtosecond lasers, for 3D optical memory systems, 294
- Pulsed infrared multiphoton excitation, isomerization via, 260
- Pulsed lasers, 65–66
- Pulsed laser systems, fluorescence imaging and, 311
- Pulse propagation, in TPA equipment design, 156
- Pulse radiolysis, 54
- Pulse radiolysis–laser flash photolysis, 57–65  
of excited radical anions, 64–65  
of excited radical cations, 58–61  
of NDI-ODN, 92–94
- Pulse shapes, 145–146, 147, 149–150, 151, 152  
in three-dimensional microfabrication, 284
- Pulse width, 152
- Pump and probe method, 54, 81, 98
- Pumped optical parametric oscillator, singlet oxygen production and, 268
- Pump intensity, absorbance change versus, 262
- Pump power, 166
- Pyrazoline moiety, in neutral donor- $\pi$ -acceptor chromophores, 190
- Pyrene–perylene bisimide system, 7–8
- Pyrene units, 7–8
- Pyridinium moiety, in symmetric acceptor- $\pi$ -acceptor chromophores, 224
- Pyridinium nitrogen, in symmetric acceptor- $\pi$ -acceptor chromophores, 221, 222
- Pyridinium ring, in ionic donor- $\pi$ -acceptor chromophores, 196

- Pyrrrole group, in symmetric acceptor- $\pi$ -acceptor chromophores, 224
- Pyryllium chromophores, 247, 249
- Quadrupolar chromophores, 136
- octupolar chromophores versus, 237
- Quadrupolar pattern
- in fluorescent chromophore/metal ion complexes, 259
- in symmetric acceptor- $\pi$ -acceptor chromophores, 216
- in symmetric donor-acceptor-donor chromophores, 226, 231, 232
- Quantum yield, 262
- fluorescence, 177, 183–185, 186, 194–195, 196–197, 198–201, 217–218, 227–228, 234–235, 253
- for singlet oxygen generation, 269, 270
- in 3D data storage, 298
- two-photon, 178
- Quantum yield for intersystem crossing ( $\Phi_{isc}$ ), 241
- Quenching, 63, 64, 65, 68–70
- intermolecular ELT, 75
- by singlet oxygen, 266
- o*-Quinodimethane, cycloaddition of, 94–96
- Quinolinium group, in ionic donor- $\pi$ -acceptor chromophores, 196
- Radical cations
- fluorescence from excited, 61–64
- pulse radiolysis of excited, 58–61
- studied via two-color two-flash photolysis, 91
- Radical concentration, in radical polymerization, 279–280
- Radical formation, in radical polymerization, 276
- Radical ions, 58. *See also* Excited radical entries
- Radical polymerization/copolymerization
- of nonlinear absorbing chromophores, 252
- two-photon initiated, 274–281
- Raman spectroscopy, of phenyl-substituted ladder-type poly(*p*-phenylene), 166
- Random hopping, 24
- Rate constants
- laser photolysis in determination of, 94–96
- two-photon absorption and, 119
- Reactive intermediates, studied via two-color two-flash photolysis, 91–98
- Read only memory (ROM) storage system, in 3D data storage, 296
- Refractive index
- complex, 120
- nonlinear, 181
- in 3D data storage, 294, 298
- in waveguide materials, 310
- Refractive-index changes, in limiting laser optical power, 324
- Relaxation process, 39
- Research, in TP applications, 328–330
- Resolution
- improving, 328
- in near-field fluorescence imaging, 312
- Resonance(s). *See also* Fluorescence resonance energy transfer (FRET)
- aromatic, 182
- bond length alternation and, 130–132
- one-photon, 211
- two-photon absorption and, 119, 123
- Resonant two-photon ionization (RTPI), 99–100
- Response theory, for two-photon absorption, 125–126, 139
- Retinal, direct excitation of, 265
- Reverse molecular motifs, 215
- Reverse saturable absorption, in limiting laser optical power, 324
- Rewritable 3D optical data storage
- materials for, 299
- photorefractive effect in, 299–300
- Rewritable technique, for 3D optical memory systems, 294
- Rhodamine B, 256–257
- Rhodamine B lactone, in 3D data storage, 296
- Rhodamine B molecules, fluorescence imaging of, 314–315
- Rigid polyphenylene-based dendritic structures, ensemble photophysics of, 1–51
- Rod shapes, in waveguide materials, 309. *See also* Conjugating rods; Connecting rods
- Ruthenium complexes, 3–4
- $S_0$  state
- in symmetric donor- $\pi$ -donor chromophores, 211, 212
- in three-level TPA model, 137–138
- $S_1$  state, 116, 124
- in symmetric donor- $\pi$ -donor chromophores, 211
- in three-level TPA model, 137–139

- $S_2$  state, 124  
in symmetric donor- $\pi$ -donor chromophores, 211  
in three-level TPA model, 137–139
- Scanning electron microscopy (SEM), 278–279  
of photonic band-gap crystal, 308  
in three-dimensional microfabrication, 293
- Schweitzer, G., 1
- Screening metal ions, fluorescent chromophores and, 257–258
- sech<sup>2</sup>-shaped pulse, 145–146, 147, 150
- Secondary electron transfer (ELT), 58
- Second generation dendrimers, 10, 11
- Second harmonic generation (SHG), 294
- Second hyperpolarizability, 182
- Selective ELT quenching, 64
- Selective protonation, in 3D data storage, 295–296
- Semiclassical perturbation theory, two-photon absorption and, 118–119
- Sensitizers, coinitiators and, 272–274, 278
- Short-lived species, multibeam irradiation photochemistry of, 53–109
- $\sigma$ -bonds, in symmetric donor- $\pi$ -donor chromophores, 213
- Silaffin-1 protein, in holographic recording, 307
- Silica nanospheres, in holographic recording, 307
- Silica shell, in nanoclinics, 315, 316
- Silver nanoparticles, surface plasmon effects and, 142, 143
- Simultaneous TPA. *See* Two-photon absorption (TPA)
- Single-photon excited fluorescence (SPEF) spectra, of symmetric acceptor- $\pi$ -acceptor chromophores, 221, 222. *See also* OP excited fluorescence (OPEF)
- Single-photon timing (SPT) measurements, 15–25, 37
- Singlet oxygen, 114  
applications of, 267–268  
in photodynamic therapy, 319  
photogeneration of, 115, 266–271
- Singlet–singlet annihilation, 31, 32–34, 40–42, 43  
energy hopping versus, 44–45, 46
- Singlet–singlet excitation hopping, 23
- “Site-selective” processes, 56
- Solvation shell, 29
- Solvatochromism, 123  
with ionic donor- $\pi$ -acceptor compounds, 193  
in neutral donor- $\pi$ -acceptor chromophores, 182, 186–187  
in octupolar chromophores, 244  
in symmetric donor-acceptor-donor chromophores, 226, 229  
in symmetric donor- $\pi$ -donor chromophores, 206
- Solvent-dependent fluorescence, 188–189
- Solvent polarity  
in oligothiophenes, 84  
symmetric donor- $\pi$ -donor chromophores and, 206
- Spherical porphyrin arrays, 7
- “Stack-of-logs” structure, in fabricating optical band-gap materials, 306
- Steady-state absorbance, 262
- Steady-state absorption spectra, 38
- Steady-state radical concentration, in radical polymerization, 280
- Steady-state spectra, 14–15
- Stepwise bond cleavage, 100–103
- Stepwise mechanism, of dissociative ELT, 80–81
- Stern–Volmer plots, 69
- “Sticky” dissociative ELT model, 79
- Stilbene (St), 197, 202  
*cis*–*trans* isomerization of, 58–61, 64  
cycloaddition of, 265  
medical applications of chromophores with, 259–260  
as TPA measurement standard, 159  
*trans*–*cis* photoisomerization of, 263–264
- Stilbene derivatives  
in three-level TPA model, 136–139  
TPA cross sections of, 174–175
- Stilbenoid pattern, 196
- Stokes–Raman scattering, 121
- Stokes shifts, 86, 87, 88, 219
- Storage density, of 3D optical memory, 294
- Streak camera detection, 98
- Strehmel, Bernd, 111
- Strehmel, Veronika, 111
- Strickler–Berg relationship, 177
- Structural isomers, chromophores in, 12–13
- Styrene, in 3D data storage, 302
- Substituent effects, on  $T_n$  state lifetimes, 71–72
- Substituted benzenes, TPA data for, 158
- Substituted benzophenone (BPD), substituent effects on  $T_n$  state lifetimes of, 71–72
- Substituted butadienes, infrared multiphoton excitation isomerization of, 260

- Substituted cyclobutenes, infrared multiphoton excitation isomerization of, 260
- Sulfonyl substituents, in octupolar chromophores, 246–247, 248
- Sum over states (SOS) method, 122–123, 125–126
- Surface plasmons, in two-photon absorption, 140–143
- Susceptibility, third-order nonlinear, 122
- Switching, in waveguide materials, 310
- Symmetric acceptor- $\pi$ -acceptor (**A- $\pi$ -A**) compounds, 215–225
- Symmetric chromophores, 197–233  
singlet oxygen and, 266–267
- Symmetric donor-acceptor-donor (**D-A-D**) compounds, 225–233
- Symmetric donor- $\pi$ -donor (**D- $\pi$ -D**) compounds, 197–215
- Symmetry. *See also* Centrosymmetric chromophores  
of octupolar chromophores, 233, 236  
two-photon absorption and, 128–130
- $T_1$  state, 116. *See also*  $T_n$  states  
singlet oxygen and, 267
- TDDFT calculations, 203. *See also* Time-dependent density functional theory (TDDFT)
- Temperature, in quantifying TPA cross section, 148. *See also* Thermal entries
- Temporal coherence ( $g_c$ ,  $g_p$ ), 149–150, 151
- Termination, of radical polymerization, 276, 279
- Terrylenediimide core, 8–9
- Tetrabutylammonium chloride, 63
- 1,2,4,5-Tetrafluorobenzene, 134
- Tetrafluorobenzene moiety, in symmetric donor-acceptor-donor chromophores, 226
- cis*-1,2,3,4-Tetrahydro-2,3-naphthalenedicarboxylic anhydride, 94–95
- Tetrahydrofuran solution, transmissivity of, 327
- Thermal blooming, 148
- Thermal lensing, 144
- Thermally crosslinked polymers, in 3D data storage, 301
- Thick samples, nonlinear absorption measurements of, 145–146, 149, 152
- Thienyl rings, in symmetric acceptor- $\pi$ -acceptor chromophores, 220
- Thin samples, nonlinear absorption measurements of, 146, 182
- Thiophene moieties  
in neutral donor- $\pi$ -acceptor chromophores, 189, 192–193  
in symmetric acceptor- $\pi$ -acceptor chromophores, 216, 224
- Thiophene-vinylene-thiophene core, in symmetric donor- $\pi$ -donor chromophores, 208–209
- Third-order nonlinear susceptibility, 122
- Three-color three-laser flash photolysis, 99–103  
stepwise bond cleavage of C–O bonds studied via, 100–103  
three-laser control of intermediate populations studied via, 99–100
- Three-dimensional data storage, 115, 293–304
- Three-dimensional data storage systems, two-photon initiated crosslinking polymerization in, 272
- Three-dimensional lithographic microfabrication (3DLM), 285–293
- Three-dimensional micro-/nanofabrication, 284–293  
two-photon initiated crosslinking polymerization in, 272, 283
- Three-dimensional object manufacture, 115, 284–293
- Three-dimensional periodic structures, fabrication of, 288–289
- Three-level TPA model, 136–139
- Three-photon absorption, 165
- Three-photon excitation, simultaneous, 179
- Time constants, 29, 36
- Time-dependent density functional theory (TDDFT), for two-photon absorption, 126, 127. *See also* TDDFT calculations
- Time-dependent DFT/B3LYP method, 190
- Time-resolved fluorescence measurements, under magic angle polarization, 15–18
- Time-resolved fluorescence polarization measurements, 18–25
- “Time-selective” processes, 55–56
- Ti:sapphire laser, 150, 151, 152, 255  
in fluorescence imaging, 313–314  
in nonlinear TPA measurements, 154–155  
in optical circuits, 309  
singlet oxygen production and, 268  
in 3D data storage, 294  
in 3D microfabrication, 286–288
- Tissue autofluorescence, 317

- Titanium dioxide (TiO<sub>2</sub>), photocatalytic reactions on surface of, 96–98
- T<sub>n</sub> states. *See also* T<sub>1</sub> state
- bond dissociation from, 72–75
  - direct observation of, 81–85
  - electron transfer from, 75–81, 82
  - energy gap law and, 70–71
  - energy transfer from, 66–70
  - stepwise bond cleavage via, 100–103
  - substituent effects on lifetimes of, 71–72
- TPA coefficient, in waveguide materials, 310.  
*See also* Two-photon absorption (TPA)
- TPA cross section ( $\delta$ ), 120, 121–122, 123, 125, 179–180, 180–181, 262
- of bioprobe chromophores, 256–257, 258–259
  - decreasing, 174–175
  - effective conjugation length and, 176–177
  - of fluorescent chromophore/metal ion complexes, 258–259
  - in ionic donor- $\pi$ -acceptor chromophores, 194–195, 196–197
  - limiting laser optical power and, 326
  - in neutral donor- $\pi$ -acceptor chromophores, 183–185, 186
  - in nonlinear absorbing chromophores, 250, 251–252, 253, 254
  - in numerical TPA simulations, 156
  - in octupolar chromophores, 233–247, 250
  - oligomer length and, 173
  - of phenyl-substituted ladder-type poly(*p*-phenylene), 166
  - of poly [1,6-bis(3,6-dihexadecyl-*N*-carbazolyl)2,4-hexadiyne], 167–168
  - of polyfluorene, 175
  - for polymers, 163
  - quantifying, 144–145, 148, 149, 150, 151, 152
  - in radical polymerization, 281
  - in relative measurement of, 155
  - surface plasmon effects and, 143
  - in symmetric acceptor- $\pi$ -acceptor chromophores, 216, 217–218, 219, 223
  - in symmetric donor-acceptor-donor chromophores, 226, 227–228, 229, 230, 231
  - in symmetric donor- $\pi$ -donor chromophores, 197, 198–201, 202, 203, 210
  - 3D optical memory and, 304
  - in three-level TPA model, 136–137
  - in TP initiators, 272, 275
  - in two-level TPA model, 133, 134
  - in unsaturated conjugated systems, 157–160
  - vibronic coupling and, 168
- TPA cross section action for fluorescence, 186
- TPA data, 156–157
- on bioprobe chromophores, 256, 257
  - from conjugated polymers/oligomers, 181–182
  - for ionic donor- $\pi$ -acceptor chromophores, 194–195
  - for neutral donor- $\pi$ -acceptor chromophores, 182, 183–185
  - for nonlinear absorbing chromophores, 253
  - for octupolar chromophores, 233, 234–235
  - on singlet oxygen chromophores, 269
  - for symmetric acceptor- $\pi$ -acceptor chromophores, 216–225
  - for symmetric donor-acceptor-donor chromophores, 227–228
  - for symmetric donor- $\pi$ -donor chromophores, 197–202
  - for TP initiators, 275
  - for triphenylmethane dyes, 237–238
- TPA equipment design, 156
- TPA microfabrication, 283
- TPA spectra, 181
- of conjugated polymers, 170
  - of nonlinear absorbing chromophores, 254
  - of octupolar chromophores, 240, 241
  - of radical polymer, 276
  - of symmetric donor-acceptor-donor chromophores, 232
  - of symmetric donor- $\pi$ -donor chromophores, 210–211
- TP chromophores, applications of, 114–115
- TPE action cross sections, 256. *See also* Two-photon excitation (TPE)
- TPE fluorescence spectra, 206
- TPEF materials, 255. *See also* Two-photon excited fluorescence (TPEF)
- TPE spectra, 181. *See also* Two-photon excitation spectra
- of fluorescent chromophores, 260
  - normalized, 170–171
  - of octupolar chromophores, 238
- TP excitation maxima. *See also* TPE entries
- in ionic donor- $\pi$ -acceptor chromophores, 194–195
  - in neutral donor- $\pi$ -acceptor chromophores, 183–185

- in nonlinear absorbing chromophores, 253, 254
- in octupolar chromophores, 234–235, 236–237, 239
- in symmetric acceptor- $\pi$ -acceptor chromophores, 216, 217–218
- in symmetric donor-acceptor-donor chromophores, 227–228, 232
- in symmetric donor- $\pi$ -donor chromophores, 197, 198–201, 202, 210, 212
- in TP initiators, 275
- TP excited states, 123–124
  - energetic ordering of, 157–159
  - of methyl-substituted ladder-type poly(*p*-phenylene), 163–165
  - of symmetric donor- $\pi$ -donor chromophores, 212
  - in three-level TPA model, 136–139
- TP fluorescence excitation spectra, 172
- TP fluorophores, 255
  - molecular engineering of, 204
- TP induced dual-channel fluorescence image formation, in 3D data storage, 295–296
- TP induced isomerization, in 3D data storage, 301
- TP initiated polymerization
  - in holographic recording, 306–307
  - optical band-gap materials and, 305
  - in 3D microfabrication, 284–285
  - in 3D optical data storage, 303–304
  - waveguide materials and, 308–310
- TP initiators, 274, 275, 276, 277, 278. *See also* Photoinitiators
- TP laser scanning microscopy (TPLSM), nanoclinics and, 317–318
- TP matrix elements, 125–126
- TP microlithography, positive, 290
- TP microscopy, 255, 259
- TP microstereolithography, 289
- TP PDT triade, 319–321
- TP photoinitiator, 190
- TP quantum yield, 178
- TP sensitizers, detecting singlet oxygen via, 268
- TP transitions, two-photon absorption and, 139–140
- TP tunneling microscope, 143
- TP two-beam picosecond “page-by-page” writing, in 3D data storage, 300–301
- TP writing and OP readout principle, in 3D data storage, 294
- trans-cis* photoisomerization. *See also* Photoisomerization
  - of diphenylbutadiene, 264–265
  - in holographic data storage, 302–303
  - of ionic donor- $\pi$ -acceptor chromophores, 196
  - of retinal, 265
  - of stilbene, 263–264
- Transient absorption, 82, 83, 86, 88–89
  - femtosecond measurements of, 37–45
  - in laser flash photolysis, 101, 102, 103
  - in 1-methoxynaphthalene, 76–77
  - in 1-NpCH<sub>2</sub>-OBP, 73
  - in oligothiophenes, 84
  - in stilbene, 58–61
  - in TiO<sub>2</sub> photocatalytic reactions, 97
- Transition densities, in three-level TPA model, 137–138
- Transition dipole, 178, 180, 186. *See also* Dipole moment (*M*)
- Transition dipole moment (*M<sub>nm</sub>*), 128–129, 140, 175–177, 180–181, 190, 216, 217–218
  - in octupolar chromophores, 233, 234–235, 237
  - in symmetric donor-acceptor-donor chromophores, 227–228
  - in symmetric donor- $\pi$ -donor chromophores, 197, 198–201, 202–203, 210, 211
  - in three-level TPA model, 136–137
  - in two-level TPA model, 133–134
- Transitions, forbidden, 168
- Transmission contrast ratio (*TCR*), limiting laser optical power and, 326
- Transmission dynamic range (*TDR*), limiting laser optical power and, 326
- Transmissivity, limiting laser optical power and, 326, 327
- Transparent multilayer polymer photochromic matrices, in 3D data storage, 301–302
- Trenches, 291
- Triarylamine dendrimer systems, 7
- Triarylmino group, 295
- Triarylsulfonium hexafluoroantimonate, in 3D data storage, 295
- Triazine(s)
  - as coinitiators, 272
  - in neutral donor- $\pi$ -acceptor chromophores, 193
- 1,3,5-Triazine, in octupolar chromophores, 239–240, 241–242, 243
- Triethanolamine coinitiator, 278

- Trifunctional acrylate monomer, in three-dimensional microfabrication, 289–290
- 1,3,5-Trimethoxybenzyl (TMB) radical cation, transient fluorescence spectrum of, 62–64, 91
- Trimethylolpropane triacrylate, in holographic recording, 307
- p*-Trimethylsilylmethylacetophenone, bond dissociation from  $T_n$  states in, 75
- p*-Trimethylsilylmethylbenzophenone, bond dissociation from  $T_n$  states in, 75
- Triphenylamine units, in octupolar chromophores, 244–246
- Triphenylmethane dyes, TPA data for, 237–238
- Triple bonds
- in neutral donor- $\pi$ -acceptor chromophores, 190–191
  - singlet oxygen generation and, 270–271
  - in symmetric acceptor- $\pi$ -acceptor chromophores, 221, 222
  - in symmetric donor- $\pi$ -donor chromophores, 205–206, 209–210, 213
- Triplet excited ( $T_n$ ) states, 55. *See also*  $T_n$  states energy transfer from, 66–70
- Triplets, in singlet oxygen generation, 269–270, 271
- Tris(2-hydroxyethyl)isocyanurate triacrylate, 293
- 1,3,5-Triscyanobenzene core, in octupolar chromophores, 238
- Trivalent boron, in neutral donor- $\pi$ -acceptor chromophores, 191
- Truncation, 291
- Tumors. *See* Cancer
- Tunneling microscope, 143
- “Twin chromophores,” 213–214
- Two-color experiment, 116
- Two-color two-laser DNA damaging, 91–94
- Two-color two-laser flash photolysis, 65–98
- control of photocatalytic reactions via, 96–98
  - excited radical cations studied via, 91
  - higher triplet excited states studied via, 66–85
  - ketyl radicals in the excited state studied via, 85–91
  - in rate constant determination, 94–96
  - of reactive intermediates, 91–98
- Two-level TPA model, 132–135
- Two-photon absorbing chromophores, 6
- Two-photon absorption (TPA), 116–156. *See also* TPA entries
- absolute evaluation of materials for, 148–155
  - applications of, 283–328
  - bond length alternation and, 130–132
  - chromophore design and optimization of, 156–255
  - covalent bonding and, 173–174
  - evaluating materials for, 144–155
  - experimental techniques in nonlinear, 144–148
  - fluorescence imaging and, 311–315
  - in ionic donor- $\pi$ -acceptor chromophores, 197
  - in limiting laser optical power, 323–328
  - mechanistic models of, 132–143
  - medical applications of, 315–321
  - by methyl-substituted ladder-type poly(*p*-phenylene), 163–165
  - in nonlinear absorbing chromophores, 250–255
  - nonlinear optical parameters and, 116–122
  - optical band-gap materials and, 305–308
  - outlook for, 328–330
  - by phenyl-substituted ladder-type poly(*p*-phenylene), 165–167
  - in photochemistry, 113–115
  - in photodynamic therapy, 318–321
  - by polyfluorenes, 167
  - prediction and discovery of, 113
  - pulse propagation in, 156
  - relative evaluation of materials for, 155
  - singlet oxygen via, 266–271
  - surface plasmons and, 140–143
  - symmetric chromophores with large, 197–233, 233–250
  - symmetry and, 128–130
  - theoretical background for, 116–132
  - theoretical methods for describing, 122–128
  - in 3D data storage, 293–304
  - three-level model of, 136–139
  - two-level model of, 132–135
  - in two-photon initiated cationic polymerization, 281–283
  - in unsaturated conjugated systems, 157–160
  - in upconverted lasing, 321–323
  - vibrational contributions to, 139–140
  - waveguide materials and, 308–310
- Two-photon active compounds, in 3D data storage, 302. *See also* TP entries
- Two-photon excitation (TPE), 116, 118–119, 121, 134–135. *See also* TPE entries

- applications of, 114–115  
efficient, 278  
fluorescence imaging and, 311  
nanoclinics and, 316–317  
organic reactions upon, 260–271  
outlook for, 328–330  
in photochemistry, 113–115  
singlet oxygen via, 267  
in three-level TPA model, 138–139
- Two-photon excitation microscope (TPEM), 311–312
- Two-photon excitation spectra, 178–179. *See also* TPE spectra  
of conjugated polymers, 171  
of fluorene dimer, 176  
of stilbene derivatives, 174
- Two-photon excited fluorescence (TPEF), 174–175, 255–260  
in absolute evaluation of TPA materials, 148–154  
of diphenylaminofluorene-based chromophores, 207  
excitation intensity and, 165, 166  
in neutral donor- $\pi$ -acceptor chromophores, 186  
quantifying TPA cross section via, 144, 147–148  
of symmetric acceptor- $\pi$ -acceptor chromophores, 221, 222  
of symmetric donor-acceptor-donor chromophores, 230–231
- Two-photon initiated cationic polymerization, 281–283
- Two-photon initiated crosslinking polymerization, 272–283
- Two-photon initiated polymerization, 274–281
- Two-photon photosciences, outlook for, 328–330
- Ultrafast energy hopping, 21. *See also* Femtosecond entries
- Ultrashort laser pulse, 298
- Ultraviolet (UV) absorption spectra, 325
- Ultraviolet excitation, cycloaddition via, 265
- Ultraviolet light, in 3D data storage, 295
- Unsaturated conjugated systems, TPA data for, 157–160
- Upconverted lasing, 321–323
- Urethane acrylate oligomers, in optical circuits, 309
- Urethane acrylates, TP initiated polymerization of, 286
- van der Auweraer, M., 1
- VB-CT model, for two-photon absorption, 128
- Vibrational contributions, to two-photon absorption, 139–140
- Vibronic coupling, TPA cross section and, 168
- Vibronic maxima, 15
- Vinylene bridges, in octupolar chromophores, 244–246
- Virtual states, in two-photon excitation, 118–119
- Volume pixels. *See* Voxel entries
- Voxels  
addressing, 300–301  
in optical band-gap materials, 305  
in waveguide materials, 309
- Voxel size ( $S$ ), 288, 289, 291, 292  
in 3D data storage, 296–298
- Wavefunctions, 118–119  
in octupolar chromophores, 236  
symmetry and, 129
- Waveguide materials, 308–310
- White-light continuum (WLC) fluorescence, 240  
of diphenylaminofluorene-based chromophores, 207
- Wire-frame objects, 289
- Write once, read many (WORM) technique, for 3D optical memory systems, 294
- Xanthene dyes, 277, 278
- Xanthone ( $X_n$ ), excited-state ketyl radicals of, 88
- Xylene moiety, in symmetric donor- $\pi$ -donor chromophores, 209–210, 210
- Yield. *See* Quantum yield entries
- Zero differential overlap (ZDO) transition densities, 138
- ZINDO/S package, 125, 128. *See also* CEO-INDO/S procedure; INDO-MRDCI/SOS method  
with three-level TPA model, 136, 138  
 $z$ -scan method, quantifying TPA cross section via, 146, 147, 182, 247  
“Z-scheme” of photosynthesis, 97
- Zwitterionic structures, bond length alternation and, 130–132



# CUMULATIVE INDEX

## VOLUMES 1-29

|  | VOL. | PAGE |
|--|------|------|
| Addition of Atoms to Olefins, in Gas Phase (Cvetanovic) . . . . .  | 1    | 115  |
| Advances in the Measurement of Correlation in Photoproduct Motion<br>(Morgan, Drabbels, and Wodtke) . . . . .                      | 23   | 279  |
| AFM and STM in Photochemistry Including Photon Tunneling (Kaupp)   | 19   | 119  |
| Alcohols, Ethers, and Amines, Photolysis of Saturated (von Sonntag and<br>Schuchmann) . . . . .                                    | 10   | 59   |
| Alkanes and Alkyl Radicals, Unimolecular Decomposition and Isotope<br>Effects of (Rabinovitch and Setser) . . . . .                | 3    | 1    |
| Alkyl Nitrites, Decomposition of and the Reactions of Alkoxy Radicals<br>(Heicklen) . . . . .                                      | 14   | 177  |
| Alternative Halocarbons, Atmospheric Photochemistry of (Francisco and<br>Maricq) . . . . .   | 20   | 79   |
| Anthracenes, Excited State Reactivity and Molecular Topology Relation-<br>ships in Chromophorically Substituted (Becker) . . . . . | 15   | 139  |
| Anti-Stokes Fluorescence, Cooling of a Dye Solution by (Zander and<br>Drexhage) . . . . .  | 20   | 59   |
| Aromatic Hydrocarbon Solutions, Photochemistry of (Bower) . . . . .  | 1    | 23   |
| Asymmetric Photoreactions of Conjugated Enones and Esters (Pete) . . .   | 21   | 135  |
| Atmospheric Reactions Involving Hydrocarbons, FTIR Studies of (Niki<br>and Maker) . . . . .  | 15   | 69   |
| Benzene, Excitation and Deexcitation of (Cundall, Robinson, and Pereira)   | 10   | 147  |
| Biocatalysis and Biomimetic Systems, Artificial Photosynthetic Trans-<br>formations Through (Willner and Willner) . . . . .        | 20   | 217  |
| Biochromophoric Systems, Excited State Behavior of Some (De Schryver,<br>Boens, and Put) . . . . .                                 | 10   | 359  |
| Cancer Treatment, Photochemistry in (Dougherty) . . . . .  | 17   | 275  |
| Carbonyl Compounds, The Photocycloaddition of, to Unsaturated Sys-<br>tems: The Syntheses of Oxetanes (Arnold) . . . . .           | 6    | 301  |

---

*Advances in Photochemistry, Volume 29* Edited by Douglas C. Neckers, William S. Jenks,  
and Thomas Wolff © 2007 John Wiley & Sons, Inc.

|  | VOL. | PAGE |
|--|------|------|
| Catalysis of Photoinduced Electron Transfer Reactions (Fukuzumi and Itoh)  | 25   | 107  |
| Cobalt (III) and Chromium (III) Complexes, the Photochemistry of, in<br>Solution (Valentine, Jr.) . . . . .                                      | 6    | 123  |
| Complexes, Photoinitiated Reactions in Weakly Bonded (Shin, Chen,<br>Nickolaissen, Sharpe, Beaudet, and Wittig) . . . . .                        | 16   | 249  |
| Cyclic Ketones, Photochemistry of (Srinivasan) . . . . .   | 1    | 83   |
| 1,2-Cycloaddition Reaction of Carbonyl Compounds and Pentaatomic<br>Heterocyclic Compounds (D'Auria, Emanuele, and Racioppi). . . . .            | 28   | 81   |
| Cyclobutanones, Solution Phase Photochemistry of (Morton and Turro). . . . .   | 9    | 197  |
| Cyclometallated Complexes, Photochemistry and Luminescence of<br>(Maestri, Balzani, Deuschel-Cornioley, and von Zelewsky). . . . .               | 17   | 1    |
| $\alpha$ -Dicarbonyl Compounds, The Photochemistry of (Monroe) . . . . .   | 8    | 77   |
| Diffusion-Controlled Reactions, Spin-Statistical Factors in (Saltiel and<br>Atwater). . . . .  | 14   | 1    |
| Electron and Energy Transfer, Mimicking of Photosynthetic (Gust and Moore)   | 16   | 1    |
| Electron Energy Transfer between Organic Molecules in Solution (Wilkinson)   | 3    | 241  |
| Electronically Excited Halogen Atoms (Hussain and Donovan) . . . . .   | 8    | 1    |
| Electron Spin Resonance Spectroscopy. Application of to Photochemistry<br>(Wan). . . . .   | 9    | 1    |
| Electron Transfer, Photoinduced in Organic Systems, Control of Back<br>Electron Transfer of (Fox). . . . .                                       | 13   | 237  |
| Electron Transfer Luminescence in Solution (Zweig) . . . . .   | 6    | 425  |
| Elementary Photoprocesses in Designed Chromophore Sequence on $\alpha$ -<br>Helical Polypeptides (Sisido). . . . .                               | 22   | 197  |
| Ensemble Photophysics of Rigid Polyphenylene Based Dendritic Structures<br>(Lor, Schweitzer, van der Anweaer, Hofkens, and De Schryver). . . . . | 29   | 1    |
| Ethyleneic Bonds, Present Status of the Photoisomerization Abut (Arai and<br>Tokumaru) . . . . .   | 20   | 1    |
| Excimers, What's New in (Yakhot, Cohen, and Ludmer) . . . . .  | 11   | 489  |
| Excited Electronic States, Electronic and Resonance Raman Spectroscopy<br>Determination of Molecular Distortions in (Zink and Shin). . . . .     | 16   | 119  |
| Flash Photolysis with Time-Resolved Mass Spectrometry (Carr) . . . . .   | 25   | 1    |
| Free Radical and Molecule Reaction in Gas Phase, Problems of Structure<br>and Reactivity (Benson) . . . . .                                      | 2    | 1    |
| FTIR Emission Studies, Time Resolved, of Photochemical Reactions<br>(Hancock and Heard) . . . . .  | 18   | 1    |
| Functional Molecular Glasses: Building Blocks for Future Optoelectronics<br>(Fuhrmann and Salbeck) . . . . .                                     | 27   | 83   |
| Gas Phase, Addition of Atoms of Olefins in (Cvetanovic) . . . . .  | 1    | 115  |
| Gas Phase Reaction, Photochemical, in Hydrogen-Oxygen System (Volman)  | 1    | 43   |
| Gas Phase Reactions, Involving Hydroxyl and Oxygen Atoms, Mech-<br>anisms and Rate Constants of (Avramenko and Kolesnika) . . . . .              | 2    | 25   |
| Halogenated Compounds, Photochemical Processes in (Major and<br>Simons) . . . . .  | 2    | 137  |
| Heterogeneous Catalysts, the Question of Artificial Photosynthesis of<br>Ammonia on (Davies, Boucher, and Edwards) . . . . .                     | 19   | 235  |

|   | VOL. | PAGE |
|---|------|------|
| Hydrogen-Oxygen Systems, Photochemical Gas Phase Reactions in (Volman) . . . . .  | 1    | 43   |
| Hydroxyl and Oxygen Atoms, Mechanisms and Rate Constants of Elementary Gas Phase Reactions Involving (Avramenko and Kolesnikova) . . . . .  | 2    | 25   |
| Hydroxyl Radical with Organic Compounds in the Gas Phase, Kinetics and Mechanisms of the Reactions of (Atkinson, Darnall, Winter, Lloyd, and Pitts) . . . . .                         | 11   | 375  |
| Hypericin and Its Perylene Quinone Analogs: Probing Structure, Dynamics, and Interactions with the Environment (Das, Halder, Chowdhury, Park, Alexeev, Gordon, and Petrich) . . . . . | 28   | 1    |
| Hypophalites, Developments in Photochemistry of (Akhtar) . . . . .  | 2    | 263  |
| Imaging Systems, Organic Photochemical (Delzenne) . . . . .   | 11   | 1    |
| Intramolecular Proton Transfer in Electronically Excited Molecules (Klöpffer) . . . . .   | 10   | 311  |
| Invention of Dylux <sup>®</sup> Instant-Access Imaging Materials and the Development of Habi Chemistry—A Personal History (Dessauer) . . . . .  | 28   | 129  |
| Ionic States, in Solid Saturated Hydrocarbons, Chemistry of (Kevan and Libby) . . . . .   | 2    | 183  |
| Isotopic Effects in Mercury Photosensitization (Gunning and Strausz) . . . . .  | 1    | 209  |
| Ketone Photochemistry, a Unified View of (Formosinho and Arnaut) . . . . .  | 16   | 67   |
| Lanthanide Complexes of Encapsulating Ligands at Luminescent Devices (Sabbatini, Guardigli, and Manet) . . . . .  | 23   | 213  |
| Laser Trapping-Spectroscopy-Electrochemistry of Individual Microdroplets in Solution (Nakatani, Chikami, and Kitamura) . . . . .  | 25   | 173  |
| Mechanism of Energy Transfer, in Mercury Photosensitization (Gunning and Strausz) . . . . .   | 1    | 209  |
| Mechanistic Organic Photochemistry, A New Approach to (Zimmerman) . . . . .   | 1    | 183  |
| Mercury Photosensitization, Isotopic Effects and the Mechanism of Energy Transfer in (Gunning and Strausz) . . . . .  | 1    | 209  |
| Metallocenes, Photochemistry in the (Bozak) . . . . .   | 8    | 227  |
| Methylene, Preparation, Properties, and Reactivities of (De More and Benson) . . . . .  | 2    | 219  |
| Molecular Distortions in Excited Electronic States, Electronic and Resonance Raman Spectroscopy Determination of (Zink and Shin) . . . . .  | 16   | 119  |
| Neutral Oxides and Sulfides of Carbon, Vapor Phase Photochemistry of the (Fileeth) . . . . .  | 10   | 1    |
| Nitric Oxide, Role in Photochemistry (Heicklen and Cohen) . . . . .   | 5    | 157  |
| Noyes, W., A., Jr., a Tribute (Heicklen) . . . . .  | 13   | vii  |
| Nucleic Acid Derivatives, Advances in the Photochemistry of (Burr) . . . . .  | 6    | 193  |
| Olefins, Photolysis of Simple, Chemistry of Electronic Excited States or Hot Ground States? (Colling) . . . . .   | 14   | 135  |
| Onium Salts, Photochemistry and Photophysics of (DeVoe, Olofson, and Sahyun) . . . . .  | 17   | 313  |
| Optical Single-Molecule Detection at Room Temperature (Meixner) . . . . .   | 24   | 1    |
| Organic Molecules, Photochemical Rearrangements of (Chapman) . . . . .  | 1    | 323  |

|  | VOL. | PAGE |
|--|------|------|
| Organic Molecules in Adsorbed or Other Perturbing Polar Environments, Photochemical and Spectroscopic Properties of (Nichollas and Leermakers) . . . . . | 8    | 315  |
| Organic Molecules in their Triplet States, Properties and Reactions of (Wagner and Hammond) . . . . .  | 5    | 21   |
| Organic Nitrites, Developments in Photochemistry of (Akhtar) . . . . .   | 2    | 263  |
| Organic Photochemical Refractive-Index Image Recording Systems (Tomlinson and Chandross) . . . . .   | 12   | 201  |
| Organized Media on Photochemical Reactions, A Model for the Influence of (Ramamurthy, Weiss, and Hammond) . . . . .                                      | 18   | 67   |
| Organo-Transition Metal Compounds, Primary Photoprocesses of (Bock and von Gustorf) . . . . .  | 10   | 221  |
| Perhalocarbons, Gas Phase Oxidation of (Heicklen) . . . . .  | 7    | 57   |
| Phenyl Azide, Photochemistry of (Schuster and Platz) . . . . .   | 17   | 69   |
| Phosphorescence and Delayed Fluorescence from Solutions (Parker) . . . . .   | 2    | 305  |
| Phosphorescence-Microwave Multiple Resonance Spectroscopy (El-Sayed) . . . . .   | 9    | 311  |
| Photoassociation in Aromatic Systems (Stevens) . . . . .   | 8    | 161  |
| Photochemical Mechanisms, Highly Complex (Johnston and Cramarossa) . . . . .   | 4    | 1    |
| Photochemical Oxidation of Aldehydes by Molecular Oxygen, Kinetics and Mechanism of (Niclause, Lemaire, and Letort) . . . . .                            | 4    | 25   |
| Photochemical Reactivity, Reflections on (Hammond) . . . . .   | 7    | 373  |
| Photochemical Rearrangements of Conjugated Cyclic Ketones: The Present State of Investigations (Schaffner) . . . . .                                     | 4    | 81   |
| Photochemical Transformations of Polyenic Compounds (Mousseron) . . . . .  | 4    | 195  |
| Photochemically Induced Dynamic Nuclear Polarization (Goetz) . . . . .   | 23   | 63   |
| Photochemistry in Cyclodextrin Cavities (Bortolus and Monti) . . . . .   | 21   | 1    |
| Photochemistry of Triarylmethane Dye Leuconitriles (Jarikov and Neckers) . . . . .   | 26   | 1    |
| Photochemistry of Conjugated Dienes and Trienes (Srinivasan) . . . . .   | 4    | 113  |
| Photochemistry of Rhodopsins, The (Ottolenghi) . . . . .   | 12   | 97   |
| Photochemistry of Short-Lived Species Using Multibeam Irradiation (Fujitsuka and Majima) . . . . .   | 29   | 53   |
| Photochemistry of Simple Aldehydes and Ketones in the Gas Phase (Lee and Lewis) . . . . .  | 12   | 1    |
| Photochemistry of the Troposphere (Levy) . . . . .   | 9    | 369  |
| Photochemistry of Vitamin D and Its Isomers and of Simple Trienes (Jacobs and Havinga) . . . . .   | 11   | 305  |
| Photochemistry, Vocabulary of (Pitts, Wilkinson, Hammond) . . . . .  | 1    | 1    |
| Photochromism (Dessauer and Paris) . . . . .   | 1    | 275  |
| Photodissociation Dynamics of Hydride Molecules: H Atom Photofragment Translational Spectroscopy (Ashfold, Mordaunt, and Wilson) . . . . .               | 21   | 217  |
| Photo-Fries Rearrangement and Related Photochemical (l,j) Shifts of (j = 3,5,7) of Carbonyl and Sulfonyl Groups (Bellus) . . . . .                       | 8    | 109  |
| Photography, Silver Halide, Chemical Sensitization, Spectral Sensitization, Latent Image Formation (James) . . . . .                                     | 13   | 329  |
| Photo-Induced and Spontaneous Proton Tunneling in Molecular Solids (Trommsdorff) . . . . .   | 24   | 147  |

|   | VOL. | PAGE |
|---|------|------|
| Photoionization and Photodissociation of Aromatic Molecules, by Ultra violet Radiation (Terenin and Vilesov) . . . . .  | 2    | 385  |
| Photoluminescence Methods in Polymer Science (Beavan, Hargreaves, and Phillips) . . . . .   | 11   | 207  |
| Photolysis of the Diazirines (Frey) . . . . .   | 4    | 225  |
| Photooxidation Reactions, Gaseous (Hoare and Pearson) . . . . .   | 3    | 83   |
| Photooxygenation Reactions, Type II, in Solution (Gollnick) . . . . .   | 6    | 1    |
| Photophysical Probes of DNA Sequence-Directed Structure and Dynamics (Murphy) . . . . .   | 26   | 145  |
| Photophysics of Gaseous Aromatic Molecules: Excess Vibrational Energy Dependence of Radiationless Processes (Lim) . . . . .   | 23   | 165  |
| Photopolymerization, Dye Sensitized (Eaton) . . . . .   | 13   | 427  |
| Photoreactive Organic Thin Films in the Light of Bound Electromagnetic Waves (Sekkat and Knoll) . . . . .   | 22   | 117  |
| Photosensitized Reactions, Complications in (Engel and Monroe) . . . . .  | 8    | 245  |
| Photosynthetic Electron and Energy Transfer, Mimicking of (Gust and Moore) . . . . .  | 16   | 1    |
| Photothermal Studies of Photophysical and Photochemical Processes by the Transient Grating Method (Terazima) . . . . .  | 24   | 255  |
| Phytochrome, Photophysics and Photochemistry of (Schaffner, Branslavsky, and Holzwarth) . . . . .   | 15   | 229  |
| Polymers, Photochemistry and Molecular Motion in Solid Amorphous (Guillet) . . . . .  | 14   | 91   |
| Pressure-Tuning Photochemistry of Metal Complexes in Solution (Eldik and Ford) . . . . .  | 24   | 61   |
| Proton-Transfer Reactions in Benzophenone/ <i>N,N</i> -Dimethylaniline Photochemistry (Peters) . . . . .  | 27   | 51   |
| Primary Processes and Energy Transfer: Consistent Terms and Definitions (Porter, Balzani and Moggi) . . . . .   | 9    | 147  |
| Quantized Matter, Photochemistry and Photoelectrochemistry of: Properties of Semiconductor Nanoparticles in Solution and Thin-Film Electrodes (Weller and Eychmüller) . . . . . | 20   | 165  |
| Quantum Theory of Polyatomic Photodissociation (Kreslin and Lester) . . . . .   | 13   | 95   |
| Radiationless Transitions, Isomerization as a Route for (Phillips, Lamaire, Burton, and Noyes, Jr.) . . . . .   | 5    | 329  |
| Radiationless Transitions in Photochemistry (Jortner and Rice) . . . . .  | 7    | 149  |
| Semiconductor Nanoclusters, Photophysical and Photochemical Processes of (Wang) . . . . .   | 19   | 179  |
| Semiconductor Photocatalysis for Organic Synthesis (Kisch) . . . . .  | 26   | 93   |
| Silver Halides, Photochemistry and Photophysics of (Marchetti and Eachus) . . . . .   | 17   | 145  |
| Single Crystals, Photochemical Mechanism in: FTIR Studies of Diacyl Peroxides (Hollingsworth and McBride) . . . . .   | 15   | 279  |
| Singlet Molecular Oxygen (Wayne) . . . . .  | 7    | 311  |
| Singlet Molecular Oxygen, Bimolecular Reactivity of (Gorman) . . . . .  | 17   | 217  |
| Singlet Molecular Oxygen, Physical Quenchers of (Bellus) . . . . .  | 11   | 105  |

|  | VOL. | PAGE |
|--|------|------|
| Singlet and Triple States: Benzene and Simple Aromatic Compounds (Noyes and Unger) . . . . .   | 4    | 49   |
| Small Molecules, Photodissociation of (Jackson and Okabe) . . . . .  | 13   | 1    |
| Solid Saturated Hydrocarbons, Chemistry of Ionic States in (Kevan and Libby) . . . . .   | 2    | 183  |
| Solvation, Ultrafast Photochemical Intramolecular Charge Transfer and Excited State (Barbara and Jarzeba). . . . .   | 15   | 1    |
| Spectroscopy and (Photochemistry of Polyatomic Alkaline Earth Containing Molecules (Bernath) . . . . .   | 23   | 1    |
| Spin Conservation (Matsen and Klein) . . . . .   | 7    | 1    |
| Stilbenes, Bimolecular Photochemical Reactions of (Lewis) . . . . .  | 13   | 165  |
| Stilbenes and Stilbene-Like Molecules, <i>Cis-Trans</i> Photoisomerization of (Görner and Kuhn) . . . . .  | 19   | 1    |
| Structure and Reactivity of Organic Intermediates as Revealed by Time-Resolved Infrared Spectroscopy (Toscano) . . . . .   | 26   | 41   |
| Sulfur Atoms, Reactions of (Gunning and Strausz) . . . . .   | 4    | 143  |
| Sulfur and Nitrogen Heteroatomic Organic Compounds, Photochemical Reaction of (Mustafa) . . . . .  | 2    | 63   |
| Supramolecularly Organized Luminescent Dye Molecules in the Channels of Zeolite L (Calzaferri, Maas, Pauchard, Pfenniger, Megelski, Devaux) . . . . .                        | 27   | 1    |
| Surfactant Solutions, Photochemistry in (von Büнау and Wolff) . . . . .  | 14   | 273  |
| The EPR Spectroscopic <i>D</i> Parameter of Localized Triplet Diradicals as Probe for Electronic Effects in Benzyl-type Monoradicals (Adam, Harer, Kita, and Nau) . . . . .  | 24   | 205  |
| The Photochemistry of Indoles (Weedon) . . . . .   | 22   | 229  |
| Theory and Applications of Chemically Induced Magnetic Polarization in Photochemistry (Wan). . . . .   | 12   | 283  |
| Thiophosgene: A Tailor-Made Molecule for Photochemical and Photophysical Studies (Moule, Fujiwara, and Lim). . . . .   | 28   | 27   |
| Transition Metal Complexes, Primary Processes in (Forster) . . . . .   | 16   | 215  |
| Triatomic Free Radicals, Spectra and Structures of (Herzberg) . . . . .  | 5    | 1    |
| Two-Photon Physical, Organic, and Polymer Chemistry: Theory, Techniques, Chromophore Design, and Applications (Strehmel and Strehmel) . . . . .                              | 29   | 111  |
| Ultraviolet Photochemistry, Vacuum (McNesby and Okabe) . . . . .   | 3    | 157  |
| Ultraviolet Photodissociation Studies of Organosulfur Molecules and Radicals: Energetics, Structure Identification, and Internal State Distribution (Cheuk-Yiu Ng) . . . . . | 22   | 1    |
| Ultraviolet Radiation, Photoionizational and Photodissociation of Aromatic Molecules by (Terenin and Vilesov) . . . . .  | 2    | 385  |
| Up-Scaling Photochemical Reactions (Braun, Jakob, Oliveros, Oller do Nascimento) . . . . .   | 18   | 235  |
| Velocity Mapping of UV Multiphoton Excited Molecules (Chandler and Parker) . . . . .   | 25   | 59   |
| Weakly Bonded Complexes, Photoinitiated Reactions in (Shin, Chen, Nickolaisen, Sharpe, Beaudet, and Wittig) . . . . .  | 16   | 249  |
| Xanthine Dyes, Photochemistry of the (Neckers and Valdes-Aguilera) . . . . .   | 18   | 315  |