

Contents

| | |
|--|-------|
| Preface | xxi |
| Taking an Evidence-Based Approach to the Care of Patients | xxii |
| How to Become an Information Master: Feeling Good about NOT Knowing Everything | xxvii |
| InfoPOEMs (Patient-Oriented Evidence that Matters) | 1 |
| Cancer | 3 |
| Breast cancer | 3 |
| Most women with breast cancer have no risk factors | 3 |
| Raloxifene reduces breast cancer risk (CORE) | 4 |
| Raloxifene and tamoxifen equally effective in reducing breast cancer risk | 5 |
| Raloxifene decreases breast cancer risk, no effect on cardiovascular disease risk (RUTH) | 6 |
| MRI more accurate than mammogram in high-risk patients | 7 |
| MRI more sensitive than mammography in high-risk women (MARIBS) | 8 |
| Digital mammography more sensitive for younger women | 9 |
| Mammography results in overdiagnosis | 10 |
| Annual mammography starting at 40 doesn't lower breast cancer mortality | 11 |
| Gynecologic cancers | 12 |
| Many unnecessary Pap smears are performed after hysterectomy | 12 |
| Specific HPV strains are associated with cervical cancer risk | 13 |
| Vaccine effective against HPV | 14 |
| Bivalent vaccine against HPV effective for 4.5 years | 15 |
| Liquid-based not better than conventional Pap | 16 |
| Endometrial sampling adequate for diagnosing endometrial cancer | 17 |
| Melanoma | 18 |
| Melanoma incidence really not rising | 18 |
| Lifetime risk of mole transforming to melanoma very low | 19 |
| 0.7% of congenital melanocytic nevi become malignant | 20 |
| Dermoscopy with validated criteria more sensitive than unaided eye | 21 |
| Larger margins better in melanoma >2 mm thick | 22 |
| Prevention and screening | 23 |
| Antioxidants don't prevent GI cancers, and may increase overall mortality | 23 |
| Vitamin E doesn't lower women's risk of cardiovascular disease or cancer | 24 |
| Antioxidants don't prevent colorectal cancer | 25 |
| Mammography, FOBT and CXR don't reduce all-cause mortality | 26 |
| Spiral CT detects early lung cancer, but screening use is premature | 27 |
| Cancer linked to some diabetes treatments | 28 |

| | |
|---|----|
| Prostate cancer | 29 |
| Finasteride of mixed benefit in preventing prostate cancer | 29 |
| Prostate cancer screening every 4 years as good as annually | 30 |
| False-positive PSA associated with increased worry, fears | 31 |
| Elevated PSA should be confirmed before biopsy | 32 |
| Radical prostatectomy improves outcomes in symptom-detected prostate cancer | 33 |
| Cardiology | 34 |
| Acute myocardial infarction and acute coronary syndromes | 34 |
| Early metoprolol in acute MI: no benefit, possible harm (COMMIT; CCS-2) | 34 |
| Insulin/dextrose infusion ineffective in AMI (HI-5) | 35 |
| Early initiation of statins following ACS does not improve outcomes | 36 |
| Antibiotics not effective in acute coronary syndromes | 37 |
| ABCDE approach to non-ST-segment elevation ACS | 38 |
| Door-to-balloon time important in STEMI | 39 |
| Routine invasive strategy may be preferred for ACS | 40 |
| Intensive medical tx + selective PCI preferred for non-ST ACS | 41 |
| Invasive tx slightly better in non-ST-elevation ACS (RITA3) | 42 |
| In ACS, 5 years of invasive tx decreases MI but not all cause mortality (FRISC-II) | 43 |
| Captopril = losartan for reducing all-cause mortality after MI | 44 |
| Captopril better than valsartan or combination post-MI | 45 |
| Implantable defibrillators are not effective post-MI | 46 |
| Anticoagulation management | 47 |
| Start warfarin at 10 mg faster for outpatient anticoagulation | 47 |
| Use an algorithm to start warfarin in older patients | 48 |
| Self-monitoring of anticoagulation safe, effective | 49 |
| Self-monitoring anticoagulation superior at preventing venous thromboembolics events | 50 |
| INR 1.5–1.9 less effective than 2.0–3.0 for idiopathic DVT | 51 |
| Oral vitamin K works faster than subcutaneous | 52 |
| Oral vitamin K effective for warfarin overdose | 53 |
| Atrial fibrillation | 54 |
| Clinical decision rules accurately predict stroke risk in atrial fibrillation | 54 |
| Thromboembolism just as likely with atrial flutter | 55 |
| Warfarin prevents more strokes than clopidogrel + ASA in atrial fibrillation (ACTIVE) | 56 |
| Ximelagatran effective in preventing stroke in atrial fibrillation | 57 |
| Quality of life with rate or rhythm control in atrial fibrillation (AFFIRM) | 58 |
| Rate control better than rhythm control in atrial fibrillation (AFFIRM) | 59 |
| Amiodarone > sotalol > placebo for maintaining NSR in atrial fibrillation (SAFE-T) | 60 |
| Coronary artery disease | 61 |
| Mediterranean diet associated with lower all cause mortality | 61 |
| Coffee does not increase risk of developing CAD | 62 |
| Vitamin E has no effect on cardiovascular disease | 63 |
| Lowering homocysteine does not reduce cardiovascular disease (HOPE 2) | 64 |
| Omega 3 fatty acids do not affect mortality | 65 |

| | |
|--|-----|
| Folic acid supplementation does not reduce cardiovascular disease risk nor mortality | 66 |
| Fenofibrate doesn't prevent coronary events in DM (FIELD) | 67 |
| Optimal oral antiplatelet therapy for vascular disease | 68 |
| Clopidogrel + ASA no better than ASA alone for high-risk patients | 69 |
| Clopidogrel beneficial added to ASA, fibrinolytic in STEMI | 70 |
| Aspirin reduces risk of CV events, increases risk of bleeding | 71 |
| Aspirin + PPI safer than clopidogrel if history of GI bleed | 72 |
| Electron beam tomography not helpful | 73 |
| Chest pain relief by NTG doesn't predict active CAD | 74 |
| Ranolazine adds little to maximum antianginal therapy (ERICA) | 75 |
| Better outcomes with CABG than PCI with stent for 2,3 vessel disease | 76 |
| Adding ACEI doesn't improve outcomes in stable angina w/nl LVEF (PEACE) | 77 |
| COX-2 inhibitors and heart disease | 78 |
| Rofecoxib increases risk of cardiovascular events | 78 |
| Celecoxib increases risk of cardiovascular complications | 79 |
| Rofecoxib, diclofenac and indomethacin increase risk of CVD | 80 |
| Heart failure | 81 |
| BNP improves outcomes in evaluation of dyspnea | 81 |
| BNP testing beneficial with CHF + pulmonary dx | 82 |
| Serial BNP levels predict risk of death and CHF after ACS | 83 |
| Higher BNP and N-ANP predict CV events | 84 |
| ARBs = ACEIs for all-cause mortality in heart failure | 85 |
| Optimal digoxin range for men 0.5 to 0.8 ng/ml | 86 |
| Digoxin increases mortality in women with heart failure | 87 |
| Nesiritide for CHF may increase mortality risk | 88 |
| Implantable defibrillators reduces mortality in NYHA Class II heart failure | 89 |
| Hyperlipidemia | 90 |
| Herbs may reduce cholesterol, no data on clinical outcomes | 90 |
| Statins prevent CAD | 91 |
| Cholesterol lowering cost-effective in high-risk elderly | 92 |
| Intensive lipid lowering of marginal benefit even if high-risk | 93 |
| Varying effects of lipid drugs on overall mortality | 94 |
| Statins equivalent for CVD prevention | 95 |
| Intensive lipid lowering with statins unnecessary with stable CAD (IDEAL) | 96 |
| Hypertension | 97 |
| JNC 7 report on prevention/evaluation/treatment of hypertension | 97 |
| Work stress has no meaningful effect on blood pressure | 98 |
| Pseudoephedrine has a minimal effect on blood pressure | 99 |
| Habitual caffeine intake does not increase risk of hypertension in women | 100 |
| Home blood pressure monitoring valuable | 101 |
| Office measurements usually overestimate blood pressure | 102 |
| Hypertension follow-up: 3 months = 6 months | 103 |
| Patients with HTN + high lipids may benefit less from statins (ALLHAT) | 104 |
| Excessive lowering of blood pressure causes more harm than good | 105 |
| Diuretics clearly first line agent for HTN (ALLHAT) | 106 |

| | |
|---|-----|
| Meta-analysis supports diuretics as first line for HTN | 107 |
| Outcomes for thiazides similar | 108 |
| ACEI better than diuretic in older men for hypertension | 109 |
| Atenolol of questionable efficacy for HTN | 110 |
| Beta-blockers > placebo, not other drugs, in preventing HTN complications | 111 |
| Renal function similarly affected by antihypertensives (ALLHAT) | 112 |
| Peripheral vascular disease and aneurysm | 113 |
| Screening program for abdominal aortic aneurysm ineffective | 113 |
| Fewer aneurysm deaths but not overall deaths with AAA screening | 114 |
| CABG not helpful before AAA or peripheral vascular surgery | 115 |
| EVAR worse than open repair of AAA (EVAR Trial 1) | 116 |
| Bypass = angioplasty for severe leg ischemia, but costs more (BASIL) | 117 |
| Pulmonary hypertension | 118 |
| Sildenafil effective for primary pulmonary hypertension | 118 |
| Sildenafil = bosentan in pulmonary hypertension (SERAPH) | 119 |
| Thrombophilias | 120 |
| Testing for prothrombotic defects not necessary after first DVT | 120 |
| Management of antiphospholipid antibody syndrome | 121 |
| Venous thromboembolism | 122 |
| Fixed-dose, subcutaneous, unfractionated heparin effective for VTE | 122 |
| Compression stockings prevent post-thrombotic syndrome | 123 |
| Routine use of vena cava filters doesn't reduce mortality | 124 |
| At least 6 months of anticoagulation optimal to prevent recurrent VTE | 125 |
| LMWH better than warfarin in preventing recurrent DVT in cancer patients | 126 |
| Venous thromboembolism diagnosis and prognosis | 127 |
| Best tests to rule in, rule out PE | 127 |
| Clinical prediction rules accurate for PE diagnosis | 128 |
| Validated algorithm for evaluating suspected PE | 129 |
| Optimal algorithm for evaluating suspected DVT | 130 |
| Multidetector CT accurate for PE, but requires clinical context | 131 |
| Negative CT scan to rule out PE equal to angiography | 132 |
| Determining major bleeding risk with warfarin for DVT | 133 |
| D-dimer useful for excluding DVT and PE | 134 |
| D-dimer <250 ng/mL predicts low risk of VTE recurrence | 135 |
| Care of Infants and Children | 136 |
| ADHD | 136 |
| Restricted diet improves parental perception of hyperactive behavior | 136 |
| Stimulants similarly effective for ADHD | 137 |
| Responders to atomoxetine do well with lower doses | 138 |
| Anemia | 139 |
| Anemia not prevented by iron in infants | 139 |
| Anemia doesn't predict iron deficiency among toddlers | 140 |

| | |
|--|-----|
| Asthma | 141 |
| Long-term budesonide does not effect adrenal function in children | 141 |
| Amoxicillin for 3 days effective for pediatric pneumonia | 142 |
| Educational programs effective for young asthmatics | 143 |
| Diarrhea | 144 |
| Nitazoxanide reduces rotavirus duration in hospitalized kids | 144 |
| Oral rotavirus vaccines safe and effective | 145 |
| Intense diet-behavior-physical activity effective for obese children | 146 |
| Evidence lacking for milk's benefit in children | 147 |
| Prevention | 148 |
| Vaccine exposure does not increase risk of infectious disease hospitalization | 148 |
| Dental restoration with amalgam (mercury) safe in children | 149 |
| School-based violence prevention programs not proved | 150 |
| Otitis media | 151 |
| Delayed prescription for AOM reduces unnecessary antibiotics | 151 |
| Parent satisfaction okay with no treatment of AOM | 152 |
| Parents prefer shared decision-making for AOM | 153 |
| Tubes marginally effective in otitis media with effusion | 154 |
| Early tympanostomy tubes do not improve outcomes after 3 or more years | 155 |
| Prompt tympanostomy tube insertion doesn't improve 9 year outcomes | 156 |
| Other pediatric problems | 157 |
| Bed and pillow covers ineffective for allergic rhinitis | 157 |
| Risk scoring system predicts mortality in pediatric ICU | 158 |
| Petroleum jelly doesn't reduce recurrent pediatric epistaxis | 159 |
| Effective and ineffective interventions for infant colic | 160 |
| Respiratory infections and fever | 161 |
| Ear thermometry is unreliable in detecting fever | 161 |
| Acetaminophen, ibuprofen comparable for pain, fever in children | 162 |
| Cough suppressants ineffective in children | 163 |
| Delayed prescriptions for URIs reduce antibiotic use | 164 |
| Amoxicillin for 3 days effective for pediatric pneumonia | 165 |
| Sore throat | 166 |
| Herbal tea effective for symptoms of acute pharyngitis | 166 |
| Steroids provide brief relief of pain in mononucleosis | 167 |
| Echinacea purpurea ineffective for treating URI in kids | 168 |
| Complementary and Alternative Medicine | 169 |
| Diarrhea | 169 |
| Yogurt prevents antibiotic-associated diarrhea | 169 |
| Probiotics helpful for antibiotic-associated diarrhea | 170 |
| Probiotics effective in preventing acute diarrhea, but not traveler's diarrhea | 171 |

| | |
|--|-----|
| Diet and exercise | 172 |
| Popular diets equally effective for losing weight | 172 |
| Mediterranean diet associated with lower all cause mortality | 173 |
| Omega 3 fatty acids do not affect mortality | 174 |
| Diet and exercise prevent diabetes in high-risk patients | 175 |
| Mental health problems | 176 |
| Restricted diet improves parental perception of hyperactive behavior | 176 |
| Variable support for CAM therapies for anxiety | 177 |
| Antioxidants do not prevent dementia | 178 |
| Lowering homocysteine with B vitamins doesn't improve cognition | 179 |
| Vitamin E, donepezil ineffective for mild cognitive impairment | 180 |
| Light therapy as effective as fluoxetine for seasonal affective disorder | 181 |
| Musculoskeletal problems | 182 |
| Acupuncture effective for chronic back pain | 182 |
| Tai chi improves symptoms of osteoarthritis | 183 |
| Acupuncture effective for OA of the knee | 184 |
| Acupuncture ineffective for fibromyalgia | 185 |
| Acupuncture better than sham treatment for neck pain | 186 |
| Prevention of cancer | 187 |
| Vitamin E not helpful, perhaps harmful | 187 |
| Vitamin E doesn't lower women's risk of cardiovascular disease or cancer | 188 |
| Vitamin E has no effect on cardiovascular disease | 189 |
| Antioxidants don't prevent GI cancers, and may increase overall mortality | 190 |
| Antioxidants don't prevent colorectal cancer | 191 |
| Green tea consumption is associated with reduced mortality | 192 |
| DHEA, testosterone not effective in aging patients | 193 |
| Lowering homocysteine does not reduce cardiovascular disease (HOPE 2) | 194 |
| Folic acid supplementation does not reduce cardiovascular disease risk nor mortality | 195 |
| Herbs may reduce cholesterol, no data on clinical outcomes | 196 |
| Upper respiratory problems | 197 |
| Herbal tea effective for symptoms of acute pharyngitis | 197 |
| Petroleum jelly doesn't reduce recurrent pediatric epistaxis | 198 |
| Saline nasal irrigation effective for frequent sinusitis | 199 |
| Forcing fluids during a respiratory infection unsupported by studies | 200 |
| Zinc nasal gel reduces duration of common cold | 201 |
| Echinacea purpurea ineffective for treating URI in kids | 202 |
| Echinacea doesn't shorten or lessen cold symptoms | 203 |
| Women's health | 204 |
| Black cohosh ineffective for vasomotor symptoms | 204 |
| Acupuncture ineffective for hot flashes | 205 |
| Lactobacillus doesn't prevent post-antibiotic vaginitis | 206 |
| Immersion exercise reduces leg edema in pregnancy | 207 |

| | |
|--|-----|
| Endocrinology | 208 |
| Diabetes | 208 |
| Diet and exercise prevent diabetes in high-risk patients | 208 |
| Treating prediabetes does not affect progression | 209 |
| Rosiglitazone delays developing diabetes mellitus (DREAM) | 210 |
| Hypertension control most important aspect of diabetes care | 211 |
| Fenofibrate doesn't prevent coronary events in diabetes mellitus (FIELD) | 212 |
| Home glucose monitoring makes little difference in type 2 diabetes | 213 |
| Rosiglitazone more likely to succeed as monotherapy clinical outcomes no better | 214 |
| ACE inhibitors beneficial in diabetics | 215 |
| ARB no better than ACEI for prevention of nephropathy progression | 216 |
| Pioglitazone ineffective in secondary prevention of macrovascular complications (PROactive) | 217 |
| Cancer linked to some diabetes treatments | 218 |
| Duloxetine decreases pain of diabetic neuropathy | 219 |
| Obesity | 220 |
| Weight loss interventions have little effect on quality of life | 220 |
| Popular diets equally effective for losing weight | 221 |
| Intense diet-behavior-physical activity effective for obese children | 222 |
| Rimonabant minimally effective for weight loss | 223 |
| Topiramate beneficial for binge eating disorder with obesity | 224 |
| Bariatric surgery effective for morbid obesity | 225 |
| Eye, Ear, Nose and Throat | 226 |
| Allergic rhinitis | 226 |
| Bed and pillow covers ineffective for allergic rhinitis | 226 |
| Pollen blocker cream effective for allergic rhinitis | 227 |
| Montelukast alone ineffective for allergic rhinitis | 228 |
| Epistaxis | 229 |
| Petroleum jelly doesn't reduce recurrent pediatric epistaxis | 229 |
| Eye problems | 230 |
| Lutein improves vision with macular degeneration in men | 230 |
| Three indicators herald bacterial conjunctivitis in adults | 231 |
| Antibiotics for conjunctivitis decreases symptom duration | 232 |
| Diet high in beta carotene, vitamins C and E, and zinc may reduce risk of macular degeneration | 233 |
| Otitis media | 234 |
| Parent satisfaction okay with no treatment of AOM | 234 |
| Parents prefer shared decision-making for AOM | 235 |
| Delayed prescription for AOM reduces unnecessary antibiotics | 236 |
| Tubes marginally effective in otitis media with effusion | 237 |
| Early tympanostomy tubes do not improve outcomes after 3 or more years | 238 |
| Prompt tympanostomy tube insertion doesn't improve 9 year outcomes | 239 |

| | |
|--|-----|
| Sinusitis | 240 |
| Saline nasal irrigation effective for frequent sinusitis | 240 |
| Intranasal steroids alone effective for acute uncomplicated sinusitis | 241 |
| Sore throat | 242 |
| Herbal tea effective for symptoms of acute pharyngitis | 242 |
| Steroids provide brief relief of pain in mononucleosis | 243 |
| URI | 244 |
| Gargling with water prevents colds | 244 |
| Delayed prescriptions for URIs reduce antibiotic use | 245 |
| Forcing fluids during a respiratory infection unsupported by studies | 246 |
| Zinc nasal gel reduces duration of common cold | 247 |
| Echinacea purpurea ineffective for treating URI in kids | 248 |
| Antibiotics slightly effective for purulent rhinitis | 249 |
| Echinacea doesn't shorten or lessen cold symptoms | 250 |
| Gastrointestinal Problems | 251 |
| Diarrhea and gastroenteritis | 251 |
| Yogurt prevents antibiotic-associated diarrhea | 251 |
| Probiotics helpful for antibiotic-associated diarrhea | 252 |
| Probiotics effective in preventing acute diarrhea, but not traveler's diarrhea | 253 |
| Nitazoxanide reduces rotavirus duration in hospitalized kids | 254 |
| Oral rotavirus vaccines safe and effective | 255 |
| Ondansetron effective for gastroenteritis with vomiting | 256 |
| Dyspepsia and gastrointestinal bleeding | 257 |
| Omeprazole 20 mg = 40 mg for primary care acid-related dyspepsia | 257 |
| Omeprazole, ranitidine effective for HP negative dyspepsia (CADET-HN) | 258 |
| 4 drugs for 1 day eliminates H. pylori | 259 |
| Aspirin + PPI safer than clopidogrel if history of GI bleed | 260 |
| Gastroesophageal reflux disease | 261 |
| A PPI test does not improve diagnosis of GERD | 261 |
| Lifestyle changes of little benefit in GERD | 262 |
| All PPIs equivalent for treatment of GERD | 263 |
| Hepatobiliary disorders | 264 |
| Acetaminophen does not affect liver function in alcoholics | 264 |
| Evaluation of abnormal liver chemistries | 265 |
| Enteral nutrition preferred in pancreatitis | 266 |
| Antibiotic prophylaxis reduces mortality in hospitalized cirrhotics | 267 |
| Gynecology | 268 |
| Cervical and endometrial cancer | 268 |
| Many unnecessary Pap smears are performed after hysterectomy | 268 |
| Specific HPV strain are associated with cervical cancer risk | 269 |

| | |
|---|-----|
| Vaccine effective against HPV | 270 |
| Bivalent vaccine against HPV effective for 4.5 years | 271 |
| Liquid-based not better than conventional Pap | 272 |
| Endometrial sampling adequate for diagnosing endometrial cancer | 273 |
| Infertility and contraception | 274 |
| Avoid nonoxynol-9 spermicides with less than 100g per dose | 274 |
| Metformin increases fertility in PCOS | 275 |
| Menopause | 276 |
| HRT with estrogen alone increases stroke, decreases fractures | 276 |
| Clonidine, gabapentin, and some SSRIs effective for hot flashes | 277 |
| Black cohosh ineffective for vasomotor symptoms | 278 |
| Acupuncture ineffective for hot flashes | 279 |
| HRT increases risk of stress, urge urinary incontinence | 280 |
| Vaginitis | 281 |
| Weekly fluconazole reduces recurrent candidiasis, though costly | 281 |
| Lactobacillus doesn't prevent post-antibiotic vaginitis | 282 |
| Hematology | 283 |
| Anemia | 283 |
| Anemia not prevented by iron in infants | 283 |
| Anemia doesn't predict iron deficiency among toddlers | 284 |
| Repeated blood testing causes anemia | 285 |
| Oral as good as IM vitamin B12 replacement | 286 |
| Thrombophilias | 287 |
| LMWH better than warfarin in preventing recurrent DVT in cancer patients | 287 |
| Testing for prothrombotic defects not necessary after first DVT | 288 |
| Management of antiphospholipid antibody syndrome | 289 |
| Infectious Disease | 290 |
| Prevention | 290 |
| Oral rotavirus vaccines safe and effective | 290 |
| Acellular pertussis vaccine 63% to 92% effective | 291 |
| Vaccine exposure does not increase risk of infectious disease hospitalization | 292 |
| Antibacterial household products don't reduce infectious symptoms | 293 |
| Prescribing practices | 294 |
| Cephalosporins often safe in PCN-allergic patients | 294 |
| Prescribing antibiotics does not save time | 295 |
| Delayed prescription for AOM reduces unnecessary antibiotics | 296 |
| Delayed prescriptions for URIs reduce antibiotic use | 297 |

| | |
|---|-----|
| Respiratory infections | 298 |
| Flu shots are effective in elderly | 298 |
| Useful signs and symptoms for diagnosis of influenza | 299 |
| Steroids provide brief relief of pain in mononucleosis | 300 |
| Kidney Diseases | 301 |
| Diabetic nephropathy | 301 |
| ACE inhibitors beneficial in diabetics | 301 |
| ARB no better than ACEI for prevention of nephropathy progression | 302 |
| Gout | 303 |
| Meat, seafood, and little dairy are risk factors for gout | 303 |
| Febuxostat = allopurinol for gout prophylaxis | 304 |
| Renal failure | 305 |
| Low-dose dopamine ineffective for acute renal failure | 305 |
| Furosemide doesn't prevent acute renal failure | 306 |
| Urinary tract infection and dysuria | 307 |
| Treating negative dipstick dysuria decreases symptoms | 307 |
| Immediate- and extended-release ciprofloxacin similar for UTI | 308 |
| Musculoskeletal Problems | 309 |
| Back and neck pain | 309 |
| Acupuncture effective for chronic back pain | 309 |
| Surgery better than no surgery for spinal stenosis | 310 |
| Acupuncture better than sham treatment for neck pain | 311 |
| Useful treatments for fibromyalgia syndrome | 312 |
| Acupuncture ineffective for fibromyalgia | 313 |
| Degenerative joint disease | 314 |
| Acetaminophen = celecoxib in DJD | 314 |
| Knee taping useful for osteoarthritis pain | 315 |
| Tai chi improves symptoms of osteoarthritis | 316 |
| Acupuncture effective for OA of the knee | 317 |
| Rofecoxib increases risk of cardiovascular events | 318 |
| Celecoxib increases risk of cardiovascular complications | 319 |
| Rofecoxib, diclofenac and indomethacin increase risk of CVD | 320 |
| Rofecoxib cardiovascular risks known in 2000 | 321 |
| Arthroscopy ineffective for osteoarthritis of knee | 322 |
| Obesity does not effect 5-year knee replacement outcomes | 323 |
| Osteoporosis and fractures | 324 |
| Monthly vitamin D prevents fractures | 324 |
| Calcium/vitamin D effective for prevention of first fracture | 325 |
| Calcium/vitamin D not effective for secondary prevention of fracture (RECORD) | 326 |
| 1000mg calcium/400 IU vit D not very effective for fracture prevention (WHI) | 327 |

| | |
|---|-----|
| HRT with estrogen alone increases stroke, decreases fractures | 328 |
| Long-term PPI use increases hip fracture risk | 329 |
| Most women quickly stop taking bisphosphonates | 330 |
| Evidence lacking for milk's benefit in children | 331 |
| Folate and vitamin B12 lowers fracture risk after stroke | 332 |
| Falls prevented in elderly by formal assessment or exercise program | 333 |
| Early mobilization better for acute limb injuries | 334 |
| Other musculoskeletal problems | 335 |
| Nonsurgical treatment effective for carpal tunnel | 335 |
| PET scan best in chronic osteomyelitis diagnosis | 336 |
| Extracorporeal shock wave therapy is ineffective for tennis elbow | 337 |
| Tennis elbow: injection better short-term, worse long-term | 338 |
| Neurology | 339 |
| Headache | 339 |
| Metoclopramide (Reglan) effective in migraine | 339 |
| Intranasal zolmitriptan effective for cluster headaches | 340 |
| Specific headache criteria useful for migraine diagnose, neuroimaging | 341 |
| Herpes zoster | 342 |
| Herpes zoster vaccine safe and effective for older adults | 342 |
| Tricyclics drug of choice for post-herpetic neuralgia | 343 |
| Insomnia | 344 |
| Melatonin effective for some sleep disorders | 344 |
| Cognitive therapy superior to zopiclone for insomnia | 345 |
| Syncope and vertigo | 346 |
| Simple maneuvers can decrease vasovagal syncope episodes | 346 |
| Steroid effective for vestibular neuritis, valacyclovir is not | 347 |
| Stroke | 348 |
| ASA prevents stroke, not MI, in women | 348 |
| ASA + dipyridamole better than ASA for cerebral ischemia (ESPRIT) | 349 |
| Warfarin increases mortality in intracranial arterial stenosis | 350 |
| Lipid-lowering prevents stroke in patients with or without CHD | 351 |
| High-dose atorvastatin reduces recurrent stroke, not mortality (SPARCL) | 352 |
| ABCD rule predicts 7-day stroke risk in TIA | 353 |
| ABCD rule predicts 7- and 30-day stroke risk in patients with TIA | 354 |
| Eight clinical findings distinguish stroke from stroke mimics | 355 |
| Noninvasive carotid imaging can replace invasive imaging | 356 |
| Three percent of TIA patients have a stroke within 30 days (BASIC) | 357 |
| Endarterectomy best within 2 weeks of stroke or TIA | 358 |
| Carotid stent inferior to carotid endarterectomy | 359 |
| PEG tubes worsen quality of life in stroke (FOOD) | 360 |

| | |
|---|-----|
| Pregnancy Care | 361 |
| Labor management | 361 |
| Fetal pulse oximetry does not reduce operative delivery | 361 |
| Suction before delivery of shoulders doesn't prevent meconium aspiration | 362 |
| VBAC with prior vaginal delivery safer than cesarean | 363 |
| Prenatal care | 364 |
| Risk-based screening for GDM effective | 364 |
| Steroids effective for hyperemesis gravidarum | 365 |
| Immersion exercise reduces leg edema in pregnancy | 366 |
| Aspirin = enoxaparin for prevention of recurrent miscarriage | 367 |
| Preterm labor | 368 |
| Prenatal vaginal infection screening + treating reduces preterm delivery | 368 |
| Oral clindamycin for BV prevents prematurity | 369 |
| Psychology and Substance Abuse | 370 |
| ADHD | 370 |
| Restricted diet improves parental perception of hyperactive behavior | 370 |
| Stimulants similarly effective for ADHD | 371 |
| Responders to atomoxetine do well with lower doses | 372 |
| Antipsychotics and schizophrenia | 373 |
| Newer antipsychotics similar to older agents (CATIE) | 373 |
| Atypical antipsychotics minimally effective, poorly tolerated in Alzheimer's disease | 374 |
| Older, newer antipsychotics both increase mortality in elderly | 375 |
| Anxiety and stress | 376 |
| Variable support for CAM therapies for anxiety | 376 |
| Work stress has no meaningful effect on blood pressure | 377 |
| Dementia | 378 |
| Antioxidants do not prevent dementia | 378 |
| Lowering homocysteine with B vitamins doesn't improve cognition | 379 |
| Reasoning training slows age-related functional decline | 380 |
| Statins not associated with decreased dementia | 381 |
| Vitamin E, donepezil ineffective for mild cognitive impairment | 382 |
| Drug therapy minimally effective for neuropsych symptoms of dementia | 383 |
| Minimal effect of cholinesterase inhibitors for Alzheimer's | 384 |
| Haloperidol shortens duration, intensity but not frequency of post-up delirium | 385 |
| Small increased risk of death with atypical antipsychotics for dementia | 386 |
| Older, newer antipsychotics both increase mortality in elderly | 387 |
| Depression | 388 |
| Two question screening for depression effective | 388 |
| No difference among new antidepressants | 389 |

| | |
|--|-----|
| Alternative medications are similarly effective for depression treatment failures (STAR*D) | 390 |
| Suicide rates equal regardless of antidepressant type | 391 |
| Antidepressant drugs increase suicidal behavior in children | 392 |
| Newer antidepressants increase suicidality in youths | 393 |
| Light therapy as effective as fluoxetine for seasonal affective disorder | 394 |
| Tobacco abuse | 395 |
| Smoking cessation 6–8 weeks before surgery reduces complications | 395 |
| 12 weeks of varenicline is effective for smoking cessation | 396 |
| 6 weeks of varenicline is also effective for smoking cessation | 397 |
| Respiratory Problems | 398 |
| Asthma | 398 |
| Educational programs effective for young asthmatics | 398 |
| Bed and pillow covers ineffective for asthma | 399 |
| Exhaled nitric oxide and sputum analysis useful for asthma diagnosis | 400 |
| Intermittent = continuous therapy for mild persistent asthma | 401 |
| Salmeterol + fluticasone better than fluticasone for asthma (GOAL) | 402 |
| Don't double steroid dose in deteriorating asthma | 403 |
| Long-term budesonide does not effect adrenal function in children | 404 |
| COPD | 405 |
| Procalcitonin test can reduce antibiotic use in COPD | 405 |
| Corticosteroids benefit patients with moderate to severe COPD | 406 |
| Anticholinergics are better than beta-agonists in COPD treatment | 407 |
| Lower respiratory infections | 408 |
| Cough suppressants ineffective in children | 408 |
| No antibiotics necessary for lower respiratory infection | 409 |
| "Chest cold" instead of "bronchitis" decreases unnecessary antibiotic use | 410 |
| Pelargonium effective for acute bronchitis symptoms | 411 |
| Single oral dose dexamethasone effective for even mild croup | 412 |
| Data on treating bronchiolitis severely limited | 413 |
| Pneumonia | 414 |
| Blood cultures add little to pneumonia care | 414 |
| Antibiotics with atypical bacteria coverage unnecessary | 415 |
| Outpatient = inpatient for low-risk patients with pneumonia | 416 |
| Antibiotic choice makes little difference in community-acquired pneumonia | 417 |
| Pneumonia: 3 days of antibiotics for uncomplicated course | 418 |
| Amoxicillin for 3 days effective for pediatric pneumonia | 419 |
| 7-day levofloxacin slightly better than 1-day azithromycin for pneumonia | 420 |
| Skin Diseases | 421 |
| Acne vulgaris | 421 |
| Milk intake correlates with increased risk of acne in women | 421 |
| Dapsone gel effective for acne vulgaris treatment | 422 |

| | |
|---|-----|
| Dermatitis | 423 |
| Effective prevention and treatment of contact dermatitis | 423 |
| Immunosuppressives = steroids for atopic dermatitis | 424 |
| Once-daily topical steroid dosing just as effective for atopic eczema | 425 |
| Herpes zoster | 426 |
| Herpes zoster vaccine safe and effective for older adults | 426 |
| Tricyclics drug of choice for post-herpetic neuralgia | 427 |
| Melanoma | 428 |
| Melanoma incidence really not rising | 428 |
| Lifetime risk of mole transforming to melanoma very low | 429 |
| 0.7% of congenital melanocytic nevi become malignant | 430 |
| Dermoscopy with validated criteria more sensitive than unaided eye | 431 |
| Other skin problems | 432 |
| Effective methods for preventing pressure ulcers | 432 |
| Duct tape seems effective for warts | 433 |
| Effective treatments for rosacea | 434 |
| Surgery + Procedures | 435 |
| Cancer | 435 |
| Larger margins better in melanoma >2mm thick | 435 |
| Radical prostatectomy improves outcomes in symptom-detected prostate cancer | 436 |
| Implantable defibrillators | 437 |
| Implantable defibrillators are not effective post-MI | 437 |
| Implantable defibrillators reduce mortality in NYHA Class II heart failure | 438 |
| Minor procedures | 439 |
| Small punch biopsies don't require sutures | 439 |
| Suturing of hand lacerations not necessary | 440 |
| Sterile gloves not necessary for laceration repair | 441 |
| Hot water (45C) immersion best for jellyfish, man-of-war stings | 442 |
| Orthopedics and sports medicine | 443 |
| Surgery better than no surgery for spinal stenosis | 443 |
| Knee taping useful for osteoarthritis pain | 444 |
| Arthroscopy ineffective for osteoarthritis of knee | 445 |
| Extracorporeal shock wave therapy is ineffective for tennis elbow | 446 |
| Tennis elbow: injection better short-term, worse long-term | 447 |
| Otitis media | 448 |
| Tubes marginally effective in otitis media with effusion | 448 |
| Early tympanostomy tubes do not improve outcomes after 3 or more years | 449 |
| Prompt tympanostomy tube insertion doesn't improve 9year outcomes | 450 |

| | |
|---|-----|
| Preoperative care | 451 |
| Beta-blockade reduces cardiac events in major noncardiac surgery | 451 |
| High-risk but not low-risk patients benefit from perioperative beta-blockade | 452 |
| Smoking cessation 6–8 weeks before surgery reduces complications | 453 |
| Vascular surgery and procedures | 454 |
| Better outcomes with CABG than PCI with stent for 2,3 vessel disease | 454 |
| Routine invasive strategy may be preferred for ACS | 455 |
| In ACS, 5 years of invasive tx decreases MI but not all cause mortality (FRISC-II) | 456 |
| CABG not helpful before AAA or peripheral vascular surgery | 457 |
| EVAR worse than open repair of AAA (EVAR Trial 1) | 458 |
| Bypass = angioplasty for severe leg ischemia, but costs more (BASIL) | 459 |
| Endarterectomy best within 2 weeks of stroke or TIA | 460 |
| Carotid stent inferior to carotid endarterectomy | 461 |
| Routine use of vena cava filters doesn't reduce mortality | 462 |
| Systems of Care | 463 |
| Care of children | 463 |
| School-based violence prevention programs not proved | 463 |
| Risk scoring system predicts mortality in pediatric ICU | 464 |
| Parent satisfaction okay with no treatment of AOM | 465 |
| Parents prefer shared decision-making for AOM | 466 |
| Critical care | 467 |
| Intensive insulin only helpful if MICU stay 3 days or longer | 467 |
| Normal saline similar to 4% albumin in ICU fluid resuscitation | 468 |
| No benefit from pulmonary artery catheters for critically ill patients | 469 |
| Pulmonary artery catheters: no harm, no benefit (PAC-Man) | 470 |
| Door-to-balloon time important in STEMI | 471 |
| Prescribing practices | 472 |
| Pharmaceutical ads are often inaccurate | 472 |
| Prescribing antibiotics does not save time | 473 |
| Delayed prescription for AOM reduces unnecessary antibiotics | 474 |
| Delayed prescriptions for URIs reduce antibiotic use | 475 |
| Process of care | 476 |
| Quality of care not related to spending | 476 |
| Disclosure of errors preferred by patients | 477 |
| Hypertension follow-up: 3 months = 6 months | 478 |
| Watchful waiting acceptable option for inguinal hernia | 479 |
| Self-monitoring | 480 |
| Self-monitoring of anticoagulation safe, effective | 480 |
| Self-monitoring anticoagulation superior at preventing venous thromboembolic events | 481 |

| | |
|---|-----|
| Urology | 482 |
| Dysuria and urinary tract infection | 482 |
| Immediate- and extended-release ciprofloxacin similar for UTI | 482 |
| Treating negative dipstick dysuria decreases symptoms | 483 |
| Incontinence | 484 |
| Urinary incontinence diagnosis with 3 questions | 484 |
| HRT increases risk of stress, urge urinary incontinence | 485 |
| PT cures postpartum stress incontinence | 486 |
| No long-term continence with pelvic floor muscle training | 487 |
| Fewer complications with condom than Foley catheters | 488 |
| Nephrolithiasis | 489 |
| Tamsulosin effective for renal colic | 489 |
| Renal colic: NSAIDs more effective, less noxious than opioid analgesics | 490 |
| Medications help passage of kidney stones | 491 |
| Prostate cancer | 492 |
| Finasteride of mixed benefit in preventing prostate cancer | 492 |
| Prostate cancer screening every 4 years as good as annually | 493 |
| False-positive PSA associated with increased worry, fears | 494 |
| Elevated PSA should be confirmed before biopsy | 495 |
| Glossary | 496 |
| Index | 498 |