

Contents

I. INTRODUCTION

1. Introduction to Virology 1

THE NATURE OF VIRUSES 2

Viruses consist of a nucleic acid genome packaged in a protein coat 2

Viruses are dependent on living cells for their replication 2

Virus particles break down and release their genomes inside the cell 2

Virus genomes are either RNA or DNA, but not both 2

WHY STUDY VIRUSES? 3

Viruses are important disease-causing agents 3

Viruses can infect all forms of life 3

Viruses are the most abundant form of life on Earth 4

The study of viruses has led to numerous discoveries in molecular and cell biology 4

A BRIEF HISTORY OF VIROLOGY: THE STUDY OF VIRUSES 5

The scientific study of viruses is very recent 5

Viruses were first distinguished from other microorganisms by filtration 5

The crystallization of tobacco mosaic virus challenged conventional notions about genes and the nature of living organisms 5

The “phage group” stimulated studies of bacteriophages and helped found the field of molecular biology 7

Study of tumor viruses led to discoveries in molecular biology and understanding of the nature of cancer 7

DETECTION AND TITRATION OF VIRUSES 8

Most viruses were first detected and studied by infection of intact organisms 8

The plaque assay arose from work with bacteriophages 8

Eukaryotic cells cultured in vitro have been adapted for plaque assays 8

Hemagglutination is a convenient and rapid assay for many viruses 9

Virus particles can be seen and counted by electron microscopy 10

The ratio of physical virus particles to infectious particles can be much greater than 1 10

THE VIRUS REPLICATION CYCLE: AN OVERVIEW 10

The single-cycle virus replication experiment 10

An example of a virus replication cycle: mouse polyomavirus 11

Analysis of viral macromolecules reveals the detailed pathways of virus replication 12

STEPS IN THE VIRUS REPLICATION CYCLE 12

1. Virions bind to receptors on the cell surface 12

2. The virion (or the viral genome) enters the cell 12

3. Early viral genes are expressed: the Baltimore classification of viruses 13

The six groups in the Baltimore classification system 13

4. Early viral proteins direct replication of viral genomes 14

5. Late messenger RNAs are made from newly-replicated genomes 15

6. Late viral proteins package viral genomes and assemble virions 15

7. Progeny virions are released from the host cell 15

2. Virus Structure 17

The molecular structure of virus particles 17

How virus structure is studied: viruses come in a variety of sizes and shapes 18

Small viruses come in simple, symmetrical packages 18

Many virus capsids have icosahedral symmetry 18

Some examples of virions with icosahedral symmetry 21

The concept of quasi equivalence 21

How many subunits can be accommodated on the capsid surface? 22

Other structures, large and small, display icosahedral symmetry 23

Many virus capsids are organized as helical tubes 23

Larger viruses come in more complex packages 24

Specific packaging signals direct incorporation of viral genomes into virions 25

Core proteins may accompany the viral genome inside the capsid 25

Scaffolding proteins help in virion assembly but are not incorporated into the mature virion 25

Viral envelopes are made from lipid bilayer membranes 26

Viral glycoproteins are inserted into the lipid membrane to form the envelope 26

Budding is driven by interactions between viral proteins 27

Assembly and disassembly of virions: the importance of an irreversible step 27

3. Virus Classification: The World of Viruses 30

VIRUS CLASSIFICATION 30

Many viruses, infecting virtually all known life forms, have been discovered 30

Virus classification is based on molecular architecture, genetic relatedness, and host organism 31

Viruses are grouped into species, genera, and families 31
 Distinct naming conventions and classification schemes have developed in different domains of virology 32

MAJOR VIRUS GROUPS 32

Study of the major groups of viruses leads to understanding of shared characteristics and replication pathways 32
 Viruses with single-stranded DNA genomes are small and have few genes 33
 Viruses with double-stranded DNA genomes include the largest known viruses 34
 Most plant viruses and many viruses of vertebrates have positive-strand RNA genomes 35
 All viruses with negative-strand RNA genomes have helical nucleocapsids and some have fragmented genomes 36
 Viruses with double-stranded RNA genomes have fragmented genomes packaged in capsids with icosahedral symmetry 37
 Viruses with a reverse transcription step in their replication cycle can have either RNA or DNA genomes 38
 Satellite viruses and satellite nucleic acids require a helper virus to replicate 39
 Viroids do not code for proteins, but replicate independently of other viruses 39

THE EVOLUTIONARY ORIGIN OF VIRUSES 40

The first steps in the development of life on earth: the RNA world 40
 Viroids and RNA viruses may have originated in the RNA world 40
 The transition to the DNA-based world 40
 Small and medium-sized DNA viruses could have arisen as independently-replicating genetic elements in cells 41
 Large DNA viruses could have evolved from cells that became obligatory intracellular parasites 41

4. Virus Entry 43

How do virions get into eukaryotic cells? 43
 Enveloped and nonenveloped viruses have distinct penetration strategies 44
 Some viruses can pass directly from cell to cell 44
 A variety of cell surface proteins can serve as specific virus receptors 45
 Receptors interact with viral glycoproteins, surface protrusions, or ‘canyons’ in the surface of the virion 45
 Many viruses enter the cell via receptor-mediated endocytosis 45
 Passage from endosomes to the cytosol is often triggered by low pH 47
 Membrane fusion is mediated by specific viral ‘fusion proteins’ 47
 Fusion proteins undergo major conformational changes that lead to membrane fusion 48
 Nonenveloped viruses penetrate by membrane lysis or pore formation 49

Virions and capsids are transported within the cell in vesicles or on microtubules 49
 Import of viral genomes into the nucleus 50
 The many ways in which viral genomes are uncoated and released 50

5. Single-Stranded RNA Bacteriophages 53

The discovery of RNA phages stimulated research into messenger RNA function and RNA replication 53
 RNA phages are among the simplest known organisms 54
 Two genera of RNA phages have subtle differences 54
 RNA phages bind to the F-pilus and use it to insert their RNA into the cell 55
 Phage RNA is translated and replicated in a regulated fashion 55
 RNA secondary structure controls translation of lysis and replicase genes 56
 Ribosomes translating the coat gene disrupt secondary structure, allowing replicase translation 57
 Ribosomes terminating coat translation can reinitiate at the lysis gene start site 57
 Replication versus translation: competition for the same RNA template 58
 Genome replication requires four host cell proteins plus the replicase 58
 A host ribosomal protein directs polymerase to the coat start site 59
 Polymerase skips the first A residue but adds a terminal A to the minus strand copy 59
 Synthesis of plus strands is less complex and more efficient than that of minus strands 59
 The start site for synthesis of maturation protein is normally inaccessible to ribosomes 61
 Synthesis of maturation protein is controlled by delayed RNA folding 61
 Assembly and release of virions 62

6. Bacteriophage ϕ X174 63

ϕ X174: a tiny virus with a big impact 63
 Overlapping reading frames allow efficient use of a small genome 64
 ϕ X174 binds to glucose residues in lipopolysaccharide on the cell surface 65
 ϕ X174 delivers its genome into the cell through spikes on the capsid surface 66
 Stage I DNA replication generates double-stranded replicative form DNA 66
 Gene expression is controlled by the strength of promoters and transcriptional terminators 66
 Replicative form DNAs are amplified via a rolling circle mechanism 67
 Summary of viral DNA replication mechanisms 67
 Procapsids are assembled by the use of scaffolding proteins 67

Scaffolding proteins have a flexible structure 68
 Single-stranded genomes are packaged into procapsids as they are synthesized 68
 Role of the J protein in DNA packaging 69
 Cell lysis caused by E protein leads to release of phage 69
 Did all icosahedral ssDNA virus families evolve from a common ancestor? 69

7. Bacteriophage T7 71

T7: a model phage for DNA replication, transcription, and RNA processing 71
 T7 genes are organized into three groups based on transcription and gene function 72
 Entry of T7 DNA into the cytoplasm is powered by transcription 73
 Transcription of class II and III genes requires a novel T7-coded RNA polymerase 73
 Class II genes code for enzymes involved in T7 DNA replication 74
 T7 RNAs are cleaved by host cell ribonuclease III to smaller, stable mRNAs 74
 Regulation of class III gene expression 74
 DNA replication starts at a unique internal origin and is primed by T7 RNA polymerase 75
 Large DNA concatemers are formed during replication 76
 Concatemers are processed and packaged into preformed proheads 76
 Special features of the T7 family of phages 76

8. Bacteriophage Lambda: A Piñata of Paradigms 79

In the beginning . . . 80
 Uptake of λ DNA depends on cellular proteins involved in sugar transport 80
 The λ lytic transcription program is controlled by termination and antitermination of RNA synthesis at specific sites on the genome 81
 The CI repressor blocks expression of the lytic program by regulating three nearby promoters: P_L , P_R , and P_{RM} 82
 Cleavage of CI repressor in cells with damaged DNA leads to prophage induction 83
 The Cro repressor suppresses CI synthesis and regulates early gene transcription 83
 Making the decision: Go lytic or lysogenize? 83
 A quick review 85
 Breaking and entering: The insertion of λ DNA into the bacterial chromosome 85
 The great escape: The liberation of λ DNA from the bacterial chromosome 86
 Int synthesis is controlled by retroregulation 86
 λ DNA Replication is directed by O and P, but carried out by host cell proteins 87
 Assembly of λ heads involves chaperone and scaffolding proteins 87

DNA is inserted into preformed proheads by an ATP-dependent mechanism 87
 Host cell lysis 88

9. Parvoviruses 89

Parvoviruses have very small virions and a linear, single-stranded DNA genome 89
 Parvoviruses replicate in cells that are going through the cell cycle 90
 Discovery of mammalian parvoviruses 90
 Parvoviruses have one of the simplest known virion structures 91
 Parvoviruses have very few genes 91
 Single-stranded parvovirus DNAs have unusual terminal structures 92
 Uncoating of parvovirus virions takes place in the nucleus and is cell-specific 92
 DNA replication begins by extension of the 3' end of the terminal hairpin 93
 The DNA "end replication" problem 93
 Steps in DNA replication 93
 Non-structural proteins are multifunctional 95
 Adenovirus functions that help AAV replication 96
 In the absence of helper virus, AAV DNA can integrate into the cell genome 96
 Parvovirus pathogenesis: the example of B19 virus 96

10. Polyomaviruses 98

Mouse polyomavirus was discovered as a tumor-producing infectious agent 98
 Simian virus 40 was found as a contaminant of Salk poliovirus vaccine 99
 Polyomaviruses are models for studying DNA virus replication and tumorigenesis 99
 Polyomavirus capsids are constructed from pentamers of the major capsid protein 99
 The circular DNA genome is packaged with cellular histones 100
 Circular DNA becomes supercoiled upon removal of histones 100
 Supercoiled DNA can be separated from relaxed or linear DNA molecules 101
 Polyomavirus genes are organized in two divergent transcription units 102
 Virions enter cells in caveolae and are transported to the nucleus 102
 The viral minichromosome is transcribed by cellular RNA polymerase II 103
 Four early mRNAs are made by differential splicing of a common transcript 104
 T antigens share common N-terminal sequences but have different C-terminal sequences 105
 T antigens bring resting cells into the DNA synthesis (S) phase of the cell cycle 105

- Small T antigen inhibits protein phosphatase 2A and induces cell cycling 105
- Middle T antigen stimulates protein tyrosine kinases that signal cell proliferation and division 105
- Large T antigen activates or suppresses transcription of cellular genes by binding to a number of important cellular regulatory proteins 107
- Large T antigen hexamers bind to the origin of DNA replication and locally unwind the two DNA strands 108
- Large T antigen hexamers assemble cellular DNA synthesis machinery to initiate viral DNA replication 110
- High levels of late transcripts are made after DNA replication begins 111
- Three late mRNAs are made by alternative splicing 112
- How do polyomaviruses transform cells in vitro and cause tumors *in vivo*? 112
- 11. Papillomaviruses 114**
- Papillomaviruses cause warts and other skin and mucosal lesions 114
- Oncogenic human papillomaviruses are a major cause of genital tract cancers 115
- Papillomaviruses are not easily grown in cell culture 115
- Papillomavirus genomes are circular, double-stranded DNA 116
- The infectious cycle follows differentiation of epithelial cells 116
- Viral mRNAs are made from two promoters and two polyadenylation signals 117
- Viral E1 and E2 proteins bind to the replication origin and direct initiation of DNA replication 118
- Viral E7 protein interacts with cell cycle regulatory proteins, particularly Rb 118
- Viral E6 protein controls the level of cellular p53 protein 120
- Synergism between E6 and E7 and the predisposition to cancer 121
- Cells transformed by papillomaviruses express E6 and E7 gene products from integrated viral DNA 121
- Future prospects for diagnosis and treatment of diseases caused by papillomaviruses 121
- 12. Adenoviruses 123**
- Adenoviruses cause respiratory and enteric infections in humans 124
- Adenoviruses can be oncogenic, but not in humans 124
- Virions have icosahedral symmetry and are studded with knobbed fibers 124
- Fibers make contact with cellular receptor proteins to initiate infection 124
- Expression of adenovirus genes is controlled at the level of transcription 126
- E1A proteins are the kingpins of the adenovirus growth cycle 127
- E1A proteins bind to the retinoblastoma protein and activate E2F, a cellular transcription factor 127
- E1A proteins also activate other cellular transcription factors 128
- E1A proteins indirectly induce apoptosis by activation of cellular p53 protein 128
- E1B proteins suppress E1A-induced apoptosis, allowing viral replication to proceed 128
- The preterminal protein primes DNA synthesis carried out by viral DNA polymerase 129
- Single-stranded DNA is circularized via the inverted terminal repeat 130
- The major late promoter is activated after DNA replication begins 131
- Five different poly(A) sites and alternative splicing generate multiple late mRNAs 131
- The tripartite leader ensures efficient transport of late mRNAs to the cytoplasm 132
- The tripartite leader directs efficient translation of late adenovirus proteins 132
- Adenoviruses kill cells by apoptosis, aiding virus release 132
- Cell transformation and oncogenesis by human adenoviruses 132
- 13. Herpes Simplex Virus 134**
- Herpesviruses are important human pathogens 135
- Most herpesviruses can establish latent infections 135
- Herpes simplex virus genomes contain both unique and repeated sequence elements 135
- Nomenclature of herpes simplex virus genes and proteins 137
- The icosahedral capsid is enclosed in an envelope along with tegument proteins 137
- Entry by fusion is mediated by envelope glycoproteins and may occur at the plasma membrane or in endosomes 138
- Viral genes are sequentially expressed during the replication cycle 138
- Tegument proteins interact with cellular machinery to activate viral gene expression and to degrade cellular messenger RNAs 139
- Immediate early (α) genes regulate expression of other herpesvirus genes 140
- β gene products set the stage for viral DNA replication 140
- Herpesvirus begins with bidirectional DNA replication 141
- Rolling-circle replication subsequently produces multimeric concatemers of viral DNA 141
- DNA replication leads to activation of γ_1 and γ_2 genes 142
- Viral nucleocapsids are assembled on a scaffold in the nucleus 143
- Envelopment and egress: three possible routes 143
- Many viral genes are involved in blocking host responses to infection 143
- The establishment and maintenance of virus latency 145
- Latency-associated transcripts include stable introns 145

14. Baculoviruses 147

- Insect viruses were first discovered as pathogens of silkworms 148
- Baculoviruses are used for pest control and to express eukaryotic proteins 148
- Baculoviruses produce two kinds of particles: “budded” and “occlusion-derived” virions 149
- Baculoviruses have large, circular DNA genomes and encode many proteins 150
- Insects are infected by ingesting occlusion bodies; infection spreads within the insect via budded virions 151
- Viral proteins are expressed in a timed cascade regulated at the transcription level 152
- Immediate early gene products control expression of early genes 152
- Early gene products regulate DNA replication, late transcription, and apoptosis 153
- Late genes are transcribed by a novel virus-coded RNA polymerase 153
- Baculoviruses are widely used to express foreign proteins 156

15. Poxviruses 157

- Smallpox was a debilitating and fatal worldwide disease 158
- Variolation led to vaccination, which has eradicated smallpox worldwide 158
- Poxviruses remain a subject of intense research interest 159
- Linear vaccinia virus genomes have covalently sealed hairpin ends and lack introns 159
- Two forms of vaccinia virions have different roles in spreading infection 160
- Poxviruses replicate in the cytoplasm 162
- Poxvirus genes are expressed in a regulated transcriptional cascade controlled by viral transcription factors 162
- Virus-coded enzymes packaged in the core carry out early RNA synthesis and processing 162
- Enzymes that direct DNA replication are encoded by early mRNAs 163
- Poxviruses produce large concatemeric DNA molecules that are resolved into monomers 164
- Postreplicative mRNAs have 5' end poly(A) extensions and 3' end heterogeneity 164
- Mature virions are formed within virus “factories” 165
- Extracellular virions are extruded through the plasma membrane by actin tails 166
- Poxviruses make several proteins that target host immune defenses 167

16. Picornaviruses 169

- Picornaviruses cause a variety of human and animal diseases including poliomyelitis and the common cold 170
- Poliovirus: a model picornavirus for vaccine development and studies of replication 170

- Picornavirus virions bind to cellular receptors via depressions or loop regions on their surface 171
- Genome RNA may pass through pores formed in cell membranes by capsid proteins 171
- Translation initiates on picornavirus RNAs by a novel internal ribosome entry mechanism 172
- Essential features of picornavirus IRES elements 173
- Interaction of picornavirus IRES elements with host cell proteins 175
- Picornavirus proteins are made as a single precursor polyprotein that is autocatalytically cleaved by viral proteinases 176
- Picornaviruses make a variety of proteinases that cleave the polyprotein and some cellular proteins 176
- Replication of picornavirus RNAs is initiated in a multiprotein complex bound to proliferated cellular vesicles 176
- RNA synthesis is primed by VPg covalently bound to uridine residues 177
- Virus assembly involves cleavage of VP0 to VP2 plus VP4 178
- Inhibition of host cell macromolecular functions 179

17. Flaviviruses 181

- Flaviviruses cause several important human diseases 182
- Yellow fever is a devastating human disease transmitted by mosquitoes 182
- A live, attenuated yellow fever virus vaccine is available and widely used 183
- Hepatitis C virus: a recently discovered member of the Flaviviridae 183
- The flavivirus virion contains an icosahedral nucleocapsid wrapped in a tightly fitted envelope 183
- Flavivirus E protein directs both binding to receptors and membrane fusion 184
- Flaviviruses enter the cell by pH-dependent fusion 185
- Flavivirus genome organization resembles that of picornaviruses 185
- The polyprotein is processed by both viral and cellular proteinases 186
- Nonstructural proteins organize protein processing, viral RNA replication, and capping 187
- Flavivirus RNA synthesis is carried out on membranes in the cytoplasm 188
- Virus assembly also takes place at intracellular membranes 189

18. Togaviruses 191

- Most togaviruses are arthropod borne, transmitted between vertebrate hosts by mosquitoes 192
- Togavirus virions contain a nucleocapsid with icosahedral symmetry wrapped in an envelope of the same symmetry 192
- Togaviruses enter cells by low pH-induced fusion inside endosome vesicles 193

Nonstructural proteins are made as a polyprotein that is cleaved by a viral protease 193
 Partly-cleaved nonstructural proteins catalyze synthesis of full-length antigenome RNA 194
 Replication and transcription: synthesis of genome and subgenomic RNAs 196
 Structural proteins are cleaved during translation and directed to different cellular locations 196
 Assembly of virions and egress at the plasma membrane 197
 Effects of mutations in viral proteins on cytopathic effects and on pathogenesis 198
 Alphaviruses have been modified to serve as vectors for the expression of heterologous proteins 199
 Alphavirus vectors have multiple potential uses 199

19. Coronaviruses 201

Coronaviruses cause common colds in humans and important veterinary diseases 202
 A newly emerged coronavirus caused a worldwide epidemic of severe acute respiratory syndrome (SARS) 202
 The SARS coronavirus may have passed from animals to humans via direct contact 202
 Coronaviruses have large, single-stranded, positive sense RNA genomes 203
 Coronaviruses fall into three groups based on genome sequences 203
 Coronaviruses have enveloped virions containing helical nucleocapsids 204
 Coronavirus virions contain multiple envelope proteins 204
 Coronavirus spike proteins bind to a variety cellular receptors 205
 The virus envelope fuses with the plasma membrane or an endosomal membrane 206
 The replicase gene is translated from genome RNA into a polyprotein that is processed by viral proteinases 206
 RNA polymerase, RNA helicase, and RNA modifying enzymes are coded by the replicase gene 207
 Replication complexes are associated with cytoplasmic membranes 207
 Genome replication proceeds via a full-length negative-strand intermediate 208
 Transcription produces a nested set of subgenomic mRNAs 208
 Subgenomic mRNAs are most likely transcribed from subgenomic negative-sense RNA templates 208
 The alternative model of discontinuous transcription of antigenome RNA is unlikely to be correct 209
 Assembly of virions takes place at intracellular membrane structures 211
 Adaptability of Coronaviruses 212

20. Paramyxoviruses and Rhabdoviruses 214

The mononegaviruses: a group of related negative-strand RNA viruses 215

Rabies is a fatal human encephalitis caused by a rhabdovirus 215
 Measles is a serious childhood disease caused by a paramyxovirus 215
 Paramyxovirus and rhabdovirus virions have distinct morphologies 216
 Viral envelope proteins are responsible for receptor binding and fusion with cellular membranes 217
 Genome RNA is contained within helical nucleocapsids 218
 Paramyxoviruses enter the cell by fusion with the plasma membrane at neutral pH 218
 Gene order is conserved among different paramyxoviruses and rhabdoviruses 219
 Viral messenger RNAs are synthesized by an RNA polymerase packaged in the virion 220
 Viral RNA polymerase initiates transcription exclusively at the 3' end of the viral genome 220
 The promoter for plus-strand RNA synthesis consists of two sequence elements separated by one turn of the ribonucleoprotein helix 220
 mRNAs are synthesized sequentially from the 3' to the 5' end of the genome RNA 222
 The P/C/V gene codes for several proteins by using alternative translational starts and by mRNA "editing" 223
 Functions of P, C and V proteins 224
 N protein levels control the switch from transcription to genome replication 224
 Virions are assembled at the plasma membrane 224

21. Filoviruses 226

Marburg and Ebola viruses: sporadically emerging viruses that cause severe, often fatal disease 227
 Filoviruses are related to paramyxoviruses and rhabdoviruses 228
 Filoviruses cause hemorrhagic fever 228
 Filovirus genomes contain seven genes in a conserved order 228
 Filovirus transcription, replication and assembly 230
 Cloned cDNA copies of viral mRNAs and viral genome RNA are used for study of filoviruses 230
 Multi-plasmid transfection systems allow recovery of infectious filoviruses 230
 Filovirus glycoprotein mediates both receptor-binding and entry by fusion 231
 Ebola virus uses RNA editing to make two glycoproteins from the same gene 232
 Does the secreted glycoprotein play a role in virus pathogenesis? 232
 Minor nucleocapsid protein VP30 activates viral mRNA synthesis in Ebola virus 233
 Matrix protein VP40 directs budding and formation of filamentous particles 234
 Most filovirus outbreaks have occurred in equatorial Africa 234

Filovirus infections are transmitted to humans from an unknown animal origin 235
 Spread of filovirus infections among humans is limited to close contacts 235
 Pathogenesis of filovirus infections 236
 Clinical features of infection 236

22. Bunyaviruses 238

Most bunyaviruses are transmitted by arthropod vectors, including mosquitoes and ticks 239
 Some bunyaviruses cause severe hemorrhagic fever, respiratory disease, or encephalitis 240
 Bunyaviruses encapsidate a segmented RNA genome in a simple enveloped particle 240
 Bunyavirus protein coding strategies: negative-strand and ambisense RNAs 240
 L RNA codes for viral RNA polymerase 241
 M RNA codes for virion envelope glycoproteins 242
 S RNA codes for nucleocapsid protein and a nonstructural protein 243
 After attachment via the virion glycoproteins, bunyaviruses enter the cell by endocytosis 243
 Bunyavirus mRNA synthesis is primed by the capped 5' ends of cellular mRNAs 243
 Coupled translation and transcription may prevent premature termination of mRNAs 244
 Genome replication begins once sufficient N protein is made 244
 Virus assembly takes place at Golgi membranes 245
 Evolutionary potential of bunyaviruses via genome reassortment 246

23. Orthomyxoviruses 248

Influenza viruses cause serious acute disease in humans, and occasional pandemics 249
 Influenza virus infections of the respiratory tract can lead to secondary bacterial infections 249
 Orthomyxoviruses are negative-strand RNA viruses with segmented genomes 249
 Eight influenza virus genome segments code for a total of ten different viral proteins 251
 Hemagglutinin protein binds to cell receptors and mediates fusion of the envelope with the endosomal membrane 252
 M₂ is an ion channel that facilitates release of nucleocapsids from the virion 252
 Nucleocapsids enter the nucleus, where mRNA synthesis and RNA replication occur 253
 Capped 5' ends of cellular pre-messenger RNAs are used as primers for synthesis of viral mRNAs 254
 Viral mRNAs terminate in poly(A) tails generated by "stuttering" transcription 255
 Two influenza A mRNAs undergo alternative splicing in the nucleus 255

Genome replication begins when newly synthesized NP protein enters the nucleus 255
 Nucleocapsids are exported from the nucleus in a complex with matrix protein and NS2 256
 The NS1 protein interferes with polyadenylation of cellular mRNAs 256
 NS1 also inhibits activation of PKR, an important antiviral pathway induced by interferon 257
 Viral envelope proteins assemble in the plasma membrane and direct budding of virions 257
 Neuraminidase cleaves sialic acid, the cellular receptor that binds to HA 257
 Influenza virus strains vary in both transmissibility and pathogenicity 257
 Genetic variability generates new virus strains that can cause pandemics 258
 The 1918 pandemic influenza A virus was probably not a reassortant virus 258
 Genome sequences from some previous influenza A virus strains confirm the antigenic shift hypothesis 258
 Highly pathogenic influenza A strains in poultry farms could lead to a new pandemic 259

24. Reoviruses 261

Reoviruses were the first double-stranded RNA viruses discovered 262
 Some members of the Reoviridae are important pathogens 262
 Reoviridae have segmented genomes made of double-stranded RNA 262
 Reovirus virions contain concentric layers of capsid proteins 263
 The attachment protein binds to one or two cellular receptors 265
 During entry, the outer capsid is stripped from virions and the core is released into the cytoplasm 265
 Enzymes in the viral core synthesize and cap messenger RNAs 266
 Translation of reovirus mRNAs is regulated 267
 Interferon and PKR: effects on viral and cellular protein synthesis 267
 Synthesis of progeny double-stranded genomes occurs within subviral particles 268
 Reoviruses induce apoptosis via activation of transcription factor NF- κ B 269
 Studies of reovirus pathogenesis in mice 270

25. Retroviruses 272

Retroviruses have a unique replication cycle based on reverse transcription and integration of their genomes 273
 Viral proteins derived from the gag, pol and env genes are incorporated in virions 273
 Retroviruses enter cells by the fusion pathway 274

Viral RNA is converted into a double-stranded DNA copy by reverse transcription 275

A copy of proviral DNA is integrated into the cellular genome at a random site 277

Sequence elements in the long terminal repeats direct transcription and polyadenylation by host cell enzymes 277

Differential splicing generates multiple mRNAs 279

The Gag/Pol polyprotein is made by suppression of termination and use of alternative reading frames 279

Virions mature into infectious particles after budding from the plasma membrane 280

Acute transforming retroviruses express mutated forms of cellular growth signalling proteins 281

Retroviruses lacking oncogenes can transform cells by insertion of proviral DNA near a proto-oncogene 282

26. Human Immunodeficiency Virus Type 1 284

Human immunodeficiency virus type 1 (HIV-1) and acquired immunodeficiency syndrome 285

HIV-1 infection leads to a progressive loss of cellular immunity and increased susceptibility to opportunistic infections 285

HIV-1 is a complex retrovirus 287

HIV-1 targets cells of the immune system by recognizing CD4 antigen and chemokine receptors 287

Virus mutants arise rapidly because of errors generated during reverse transcription 288

Unlike other retroviruses, HIV-1 directs transport of proviral DNA into the cell nucleus 288

Latent infection complicates the elimination of HIV-1 289

The Tat protein increases HIV-1 transcription by stimulating elongation by RNA polymerase II 289

The Rev protein mediates cytoplasmic transport of viral mRNAs that code for HIV-1 structural proteins 290

Together, the Tat and Rev proteins strongly upregulate viral protein expression 291

The Vif protein increases virion infectivity by counteracting a cellular deoxycytidine deaminase 291

The Vpr protein enables the preintegration complex to be transported to the nucleus 291

The Vpu protein enhances release of progeny virions from infected cells 291

The Nef protein is an important mediator of pathogenesis 292

27. Human T-Cell Leukemia Virus Type 1 294

Discovery of the first human retrovirus 294

Like lentiviruses, HTLV-1 codes for regulatory proteins by producing doubly-spliced mRNAs 295

HTLV-1 Rex regulates polyadenylation, splicing, and nuclear export of viral RNAs 296

HTLV-1 Tax regulates transcription of viral and cellular genes 297

Cell transformation by HTLV-I is mediated by Tax 299

The interleukin-2†autocrine loop stimulates T-cell proliferation 299

Activation of the Jak-Stat pathway by p12^I mimics interleukin-2 stimulation 299

Cell Cycle Progression: p16^{INK4A} and cyclin-dependent kinases 300

The mitotic spindle checkpoint and MAD1 300

Downregulation of p53 activity by Tax allows T cell proliferation 301

Diseases caused by HTLV-1 develop slowly and can be severe 301

Coinfection by HIV-1 and HTLV-1 is an emerging problem 302

Antiviral therapy of disease caused by HTLV-1 has not met with great success 302

28. Hepadnaviruses 303

At least seven distinct viruses cause human hepatitis 304

The discovery of hepatitis B virus 304

Dane particles are infectious virions; abundant noninfectious particles lack nucleocapsids 305

The viral genome is a circular, partly single-stranded DNA with overlapping reading frames 305

Nucleocapsids enter the cytoplasm via fusion and are transported to the nucleus 305

Transcription of viral DNA gives rise to several mRNAs and a pregenome RNA 307

The roles of hepatitis B virus proteins 308

The pregenome RNA is packaged by interaction with polymerase and core proteins 309

Genome replication occurs via reverse transcription of pregenome RNA 309

Virions are formed by budding in the endoplasmic reticulum 311

Hepatitis B virus can cause chronic or acute hepatitis, cirrhosis, and liver cancer 312

Hepatitis B virus is transmitted by blood transfusions, contaminated needles, and unprotected sex 313

A recombinant vaccine is available 313

Antiviral drug treatment has real but limited success 313

29. Viroids and Hepatitis Delta Virus 315

Viroids are small, circular RNAs that do not encode proteins 316

Group A and group B viroids have distinct properties 316

Viroids replicate via linear multimeric RNA intermediates 317

Three enzymatic activities are needed for viroid replication 317

How do viroids cause disease? 318
 Interaction of viroid RNA with cellular RNAs or proteins
 may disrupt cell metabolism 318
 Plant satellite RNAs resemble viroids but are
 encapsidated 320
 Hepatitis delta virus is a human viroid-like satellite
 virus 320
 Hepatitis delta virus may use two different cellular RNA
 polymerases to replicate 320
 RNA editing generates two forms of hepatitis delta
 antigen 322
 Conclusion: viroids may be a link to the ancient RNA
 world 322

30. Prions 323

Prions are proteins that cause fatal brain diseases 323
 Prion diseases were first detected in domestic
 ruminants 324
 Human prion diseases can be either inherited or
 transmitted 324
 The infectious agent of prion diseases contains protein but
 no detectable nucleic acid 325
 PrP^{Sc} is encoded by a host cell gene 325
 Differences between PrP^C and PrP^{Sc} 326
 The prion hypothesis: formation of infectious and
 pathogenic prions from normal PrP^C 326
 Is the prion hypothesis correct? 327
 Pathology and diagnosis of prion diseases 330
 Genetics of prion diseases 330
 Prion diseases are not usually transmitted among different
 species 330
 Strain variation and crossing of the species barrier 331
 The nature of the prion infectious agent 331

31. Interferons 333

Virus-infected cells secrete interferons, which protect nearby
 cells against virus infection 334
 Interferons are a first line of host defense against viruses but
 therapeutic use has been limited 334
 Interferons α , β , and γ are made by different cells and have
 distinct functions 335
 Transcription of interferon genes is activated by virus
 infection or double-stranded RNA 335
 Transcriptional activation occurs by binding of transcription
 factors to interferon gene enhancers 335
 Interferon signal transduction is carried out via the Jak-Stat
 pathway 337
 Antiviral activities induced by interferon 338
 Interferons have diverse effects on the immune system 340
 The adaptive immune system 340
 Interferons stimulate antigen processing and
 presentation 342

Interferon and the development of CD4-positive helper
 T-cells 342
 The role of interferon in macrophage activation and cellular
 immunity 343
 Effects of interferons on antibody production 343
 Interferons regulate cell growth and apoptosis 343
 Viruses have developed numerous strategies to evade the
 interferon response 343
 Conclusion: interferons are a first line of defense against
 virus infection 344

32. Antiviral Chemotherapy 346

The discovery and widespread use of antiviral compounds
 began only recently 346
 Importance of antiviral drugs for basic science 347
 How are antiviral drugs obtained? 347
 Targeting drugs to specific steps of virus infection 347
 Capsid-binding drugs prevent attachment and entry of
 virions 348
 Amantadine blocks ion channels and inhibits uncoating of
 influenza virions 349
 Nucleoside analogues target viral DNA polymerases 349
 Acyclovir is selectively phosphorylated by herpesvirus
 thymidine kinases 350
 Acyclovir is preferentially incorporated by herpesvirus DNA
 polymerases 351
 Cytomegalovirus encodes a protein kinase that
 phosphorylates ganciclovir 351
 HIV-1 reverse transcriptase preferentially incorporates
 azidothymidine into DNA, leading to chain
 termination 354
 Nonnucleoside inhibitors selectively target viral replication
 enzymes 354
 Protease inhibitors can interfere with virus assembly and
 maturation 354
 Ritonavir: a successful protease inhibitor of HIV-1 that was
 developed by rational methods 354
 Neuraminidase inhibitors suppress release and spread of
 influenza virus 355
 Antiviral chemotherapy shows promise for the
 future 357

33. Eukaryotic Virus Vectors 358

Many viruses can be engineered to deliver and express
 specific genes 358
 Virus vectors are used to produce high levels of specific
 proteins in cultured cells 359
 Gene therapy is an expanding application of virus
 vectors 360
 Virus vectors are produced by transfection of cells with
 plasmids containing deleted genomes 360
 Virus vectors are engineered to produce optimal levels of
 gene products 361

ADENOVIRUS VECTORS 362

Adenovirus vectors are widely used in studies of gene transfer and antitumor therapy 362

Replication-defective adenovirus vectors are propagated in complementing cell lines 362

Replication-competent adenovirus vectors are useful tools in antitumor therapy 363

Advantages and limitations of adenovirus vectors 363

RETROVIRUS VECTORS 364

Retrovirus vectors incorporate transgenes into the cell chromosome 364

Packaging cell lines express retrovirus enzymatic and structural proteins 364

Strategies for controlling transgene transcription 364

Lentivirus vectors are used for gene delivery to nondividing cells 365

Production of lentivirus vectors requires additional cis-acting sequences 366

Applications of retrovirus vectors: treatment of blood disorders 367

Advantages and limitations of retrovirus vectors 367

ADENO-ASSOCIATED VIRUS VECTORS 367

Adeno-associated virus vectors can insert transgenes into a specific chromosomal locus 367

Production of AAV vectors usually requires a helper virus 367

Applications of adeno-associated virus vectors: treatment of hemophilia 368

Advantages and limitations of AAV vectors 368

34. Viral Vaccines 370

A brief history of viral vaccines 371

Early vaccine technology was crude but effective 372

Embryonated chicken eggs and cell culture played major roles in recent vaccine development 373

Major categories of viral vaccines 373

Advantages and drawbacks of vaccine types 374

How do viral vaccines work? 375

The role of the immune system in fighting viral infections 375

Some vaccines target mainly antibody production; others target the cellular immune response 376

New approaches to vaccine development show great promise 376

The changing vaccine paradigm 378

Vaccine-associated adverse events 378

Ethical issues in the use of viral vaccines 380