

Index

• Numerics •

3rd ventricle, 31
4th ventricle, 31
911, calling, 13–14, 192, 193–195, 238

• A •

AAA (American Automobile Association), 286
AARP (American Association of Retired Persons), 286
accident, 58
ACE (angiotensin converting enzyme) inhibitors, 149
active, less, 241
acute ischemic stroke, 199. *See also* white stroke
acute stroke, 329
Administration on Aging, 270
advertising, 173, 181
affairs, getting in order, 305–306
African Americans
 high blood pressure, 140
 stroke risk and, 171
age
 blood vessels, 91
 brain hemorrhage, 86
 dementia, 119, 124
 dizziness, evaluating, 76
 job limitations, accepting, 252
 stroke risk, 170
 subarachnoid hemorrhage, 110
Aggrenox (dypiramide), 176, 325
aides, nurses', 268–269
alcohol
 abuse, 172
 high blood pressure, 142
 stroke dementia, preventing, 130
Alzheimer's disease, 120
amaurosis fugax (fleeting blindness), 72
American Association of Retired Persons (AARP), 286
American Automobile Association (AAA), 286
American Heart Association, 310
amyloid angiopathy, 91, 96, 329
aneurysm
 described, 20, 101, 103, 329
 finding, 112–113
 high blood pressure, 139
 rupture, 104–107
 second opinion, seeking, 212
 warning signs, 104
angiogram
 aneurysm, 112–113
 atherosclerosis, 209
 AVMs, 90
 described, 43
 white stroke, 65
angiography, 329
angiotensin converting enzyme (ACE) inhibitors, 149
anterior cerebral artery, 38
antibiotics, 290
anticoagulant, 129, 329
antidepressant, 173
aphasia, 230, 259, 329
appointment, medical, 245
arachnoid, 102
arm
 brain portion controlling, 33
 falls asleep, 75
 sudden numbness or weakness, 191, 213
arteriogram, 90, 183
arteriovenous malformations (AVMs), 90, 211
artery
 brain workings, 38
 clotting, 51
 defined, 329
 narrowed, finding with carotid ultrasound, 208–209
 red stroke (SAH) and, 103
 spasm, treating, 115–116
 stenosis, 209–210

artery, carotid
 bleeding, 52
 described, 37, 72–73, 330
 evaluating, 326–327
 ultrasound, 80, 208–209

artificial heart valve, 176

aspirin, 80, 129, 176, 325

assets, listing, 264–265

assisted-living facilities
 adjusting to life, 271–274
 care needs, defining, 260–264
 community, 271
 costs, 270–271
 game plan, 259–260
 grappling with decision, 258–259
 helping family choose, 257–258
 problems, handling, 271
 resources, determining, 264–267
 transportation to appointments, 271

asthma, 149

atherosclerosis
 angiogram finding, 209
 dementia, 121
 described, 156–157, 329
 dietary fat, 54
 high blood pressure with, 136–137, 139
 risk for white stroke, 54
 surgery, 54, 81, 209
 ultrasound testing for, 183
 white stroke and, 16–17, 72

atherosclerotic plaque, 330

atherothrombotic stroke. *See* white stroke

Atkins diet, 164

Atorvastatin, 163

atrial fibrillation
 checking for, 79, 248
 described, 175, 330
 medication, 89
 risk for white stroke, 55–56
 treating, 326

attention deficit, 123

attorney, medical power of, 131

attorneys, 305

averaging blood pressure, 138

AVMs (arteriovenous malformations),
 90, 211

• B •

backsliding, rehabilitation, 227

Baclofen (Liorisal), 279

balance problems, 191

basilar artery, 37

bed sores, 218, 273

behavior modification, 174

behavioral changes, 109–110

benzodiazepines, 279–280

berry aneurysm, 88, 330

beta blockers, 146, 147–149

bile, 153

birth control pills, 55–56, 172, 211–212, 245

bladder infection, 290

blebs, 19, 87–88

bleeding
 around brain, 20
 within brain, 19
 second SAH, 204
 TPA, usefulness of, 202
 into vessel wall, 52

blind spot, discovering, 287

blood clot
 brain workings, 38–40
 deficiencies, 92
 defined, 330
 dissolving, 66
 encouraging to treat red stroke, 98
 forming, 73–74
 hardening, 51–52
 heart and vascular disease, 176–178
 medication, 80–81, 129
 preventing, 325
 ventricles, 32
 white stroke, 50, 51

blood clots in legs
 pulmonary embolism, 215
 risk, 214–215
 treatments, 215–216

blood flow, 37–38

blood pressure
 checking, 79
 heart, monitoring, 217
 lowering, 129, 131
 preventing red strokes, 99

- reducing, 97
 - slightly higher, 182
 - small strokes and, 121
 - TPA, usefulness of, 202
 - tracking, 183
 - blood pressure, high
 - with atherosclerosis, 136–137
 - averaging, 138
 - brain hemorrhage, 86
 - checking, 139
 - defining, 137–138
 - described, 331
 - lowering with medication, 129, 131, 145–150
 - lowering without medication, 144–146
 - measuring, 137
 - medication causing stroke, 139, 247
 - prevalence, 135
 - red stroke, 87–88, 211
 - risk factors, 140–144
 - sudden and severe increase, 92
 - treating, 324–325
 - white stroke and, 53–54
 - blood pressure, low
 - measuring, 138
 - medications going too far, 145–146
 - blood sugar, 77
 - blood tests
 - dementia, diagnosing, 127
 - diet, monitoring effects of, 164
 - doctor's visit, scheduling, 80
 - medical problems related to high blood pressure, 142–143
 - blood vessel. *See also* dissection
 - abnormalities, 90
 - brain workings, 36–37
 - damage, 16–17
 - lining, splitting, 16, 17–18
 - scarring, 122
 - small strokes and, 121
 - blood-pressure medication, 57
 - blood-thinning medication, 89
 - BMI (Body Mass Index) table, 157–158
 - body
 - brain portion controlling, 33
 - shape, 157
 - botox (botulinum toxin), 280
 - bowel problems
 - control, regaining, 289–290
 - nursing home solution, 290
 - prevalence, 288
 - braces, wearing, 224–225, 278
 - Brain Attack Coalition, 13, 190–191
 - brain bleeding, 60. *See also* red stroke
 - brain cells, exercising, 25
 - brain edema, 108
 - brain hemorrhage. *See* red stroke
 - brain injury, 276–277
 - brain packaging, 102–103
 - brain recovery, 225–226
 - brain stem
 - described, 31, 330
 - heart, monitoring, 217
 - subarachnoid hemorrhage, 111
 - brain swelling
 - deformation, 93–94
 - injury, white stroke, 59–60
 - monitoring, 213–214
 - red stroke (SAH), 108
 - relieving, 115
 - ventricles, 32
 - brain workings
 - arteries, 38
 - blood clots, 38–40
 - blood flow, 37–38
 - blood vessels, 36–37
 - illustration, 30
 - left and right hemispheres, 35–36
 - nerves, 32–34
 - parts, 31–32
 - response to injury, 40–41
 - senses, managing, 34–35
 - viewing technologies, 41–44
 - breakfast, 246
 - broken blood vessels, 15
 - bruises, 273
 - bruit sound, 53, 80
 - bubble-like growths, 19
 - Bupropion, 173
 - burnout, caregiver, 299–300
- C •
- calcium, 141
 - calcium antagonists, 150
 - calcium channel blockers, 115, 150

- calling 911, 13–14, 192, 193–195, 238
- calories, 180
- cancer, 98
- capable, less, 240
- capillaries, 37
- car, driving
 - alternatives, identifying, 286–287
 - consequences, 285
 - evaluation, 285–286
 - signs, reading, 284
 - work, returning to, 253
- Cardio Pulmonary Rescue (CPR), 193
- cardioembolic stroke. *See* white stroke
- care manager, 260
- care needs, defining, 260–264
- caregiver
 - burnout, preventing, 299–300
 - depression, discussing, 243
 - medical care, 300
 - priorities, including, 263–264
 - rehabilitation, 231–232
 - special advice, 301
- care-planning meetings, nursing home, 269
- carotid artery
 - bleeding, 52
 - described, 37, 72–73, 330
 - evaluating, 326–327
 - ultrasound, 80, 208–209
- carotid endarterectomy, 81, 209, 330
- CAT (computerized axial tomography) scan
 - aneurysm, finding, 112
 - brain hemorrhage, 86, 88, 93, 95
 - dementia, 126
 - described, 41–42, 43, 330
 - doctor's visit, 80
 - in emergency room, 200
 - following headache, 105
 - infarct, 49
 - subarachnoid hemorrhage, 106, 111
 - ventricles, 32
 - white stroke, diagnosing, 64
- catheter, 291
- Caverly, Brian (*Estate Planning For Dummies*), 306
- cavities, fluid-filled, 31–32
- cerebellum
 - clot, removing, 96
 - illustrated, 31
- cerebral angiogram, 209, 211
- cerebral infarct, 48
- cerebrospinal fluid (CSF), 102–103, 201, 330
- cerebrovascular accident (CVA), 16, 48, 199. *See also* white stroke
- cerebrum, 31
- certification, stroke center, 25
- champions, promoting stories about, 314
- changing needs, planning for, 263
- checking blood pressure, 139
- checklist
 - daily homecoming, 255
 - leaving rehabilitation center, 234
 - risk factors, 184
- check-up
 - dementia, diagnosing, 126
 - following TIA, 79–80
- chest x-ray, 184
- children
 - adult, 299
 - young, 294
- choking, 216, 230
- cholesterol
 - described, 330
 - diet, 154
 - HDL, ratio to, 161
 - reducing, 325–326
 - slightly higher, 182
 - stroke dementia, preventing, 130
 - stroke risk, 171
 - thiazide diuretics raising, 147
- cholesterol test
 - described, 158–159
 - evidence, collecting, 183
 - fasting before, 159–160
 - in hospital, 210
 - results, interpreting, 160–161
- chylomicrons, 156
- cigarettes. *See* smoking tobacco
- cleanliness, nursing home, 269
- clinical trials
 - list of, 314
 - volunteering for, 315
- clips
 - aneurysms, sealing off, 113–114
 - choosing, 115
 - second opinion, seeking, 212
- clock, timing stroke, 193
- clonazepam (Rivotril), 279–280
- clopidogrel (Plavix), 129, 176, 325

- clot
 - brain workings, 38–40
 - deficiencies, 92
 - defined, 330
 - dissolving, 66
 - encouraging to treat red stroke, 98
 - forming, 73–74
 - hardening, 51–52
 - heart and vascular disease, 176–178
 - medication, 80–81, 129
 - preventing, 325
 - ventricles, 32
 - white stroke, 50, 51
- clot-buster. *See* TPA
- clots in legs
 - pulmonary embolism, 215
 - risk, 214–215
 - treatments, 215–216
- clumsiness, 76
- coach class syndrome, 214–216
- cocaine, 92
- coffee, 246
- coils
 - aneurysms, sealing off, 114
 - choosing, 115
- collateral flow, 61–62
- color-coding stroke types, 15
- communication problems, 123
- community
 - assisted-living facilities, 271
 - champions, promoting stories about, 314
 - emergency medical services, checking into, 312–313
 - friends and family, sharing knowledge with, 311
 - health policy, 315
 - local hospitals, working with, 311–312
 - local stroke clubs and support groups, 309–310
 - regional, state, and national organizations, 310
 - research, 314–315
 - schoolchildren, educating, 313
 - stroke recovery group, starting, 313
- community stroke center, 204
- complaints, nursing home, 269
- computerized tomography/computerized axial tomography scan (CT scan)
 - aneurysm, finding, 112
 - brain hemorrhage, 86, 88, 93, 95
 - dementia, 126
 - described, 41–42, 43, 330
 - doctor's visit, 80
 - in emergency room, 200
 - following headache, 105
 - infarct, 49
 - subarachnoid hemorrhage, 106, 111
 - ventricles, 32
 - white stroke, diagnosing, 64
- confidence, 241
- confusion, 123
- connection point, nerve cell, 32
- constipation, 147, 165, 289
- contractures, 277
- control, less in, 241
- controllable risks, 22–23
- conversation, difficulty following, 123
- convulsions
 - drug withdrawal, sudden, 279
 - evaluating, 76–77
 - hospital treatment, 214
 - injury, white stroke, 60
 - subarachnoid hemorrhage, 109
- coordination problems, 191
- costs
 - assisted-living facilities, 270–271
 - nursing home, 267
- Coumadin, 177
- couples, 294
- CPR (Cardio Pulmonary Rescue), 193
- C-reactive protein, 159
- cross circulation minimizing stroke, 63–64
- crossing legs, 139
- carotid endarterectomy, 81
- CSF (cerebrospinal fluid), 102–103, 201, 330
- CT (computerized tomography) scan
 - aneurysm, finding, 112
 - brain hemorrhage, 86, 88, 93, 95
 - dementia, 126
 - described, 41–42, 43, 330
 - doctor's visit, 80
 - in emergency room, 200
 - following headache, 105
 - infarct, 49
 - subarachnoid hemorrhage, 106, 111
 - ventricles, 32
 - white stroke, diagnosing, 64

curriculum, rehabilitation, 227–228
 CVA (cerebrovascular accident), 16, 48,
 199. *See also* white stroke



daily checklist, 255
 dairy products, 141
 Dantrolene, 280
 daycare facility, 299
 death rate
 overall stroke, reduction in, 169
 red stroke, 85
 SAH red stroke, 102, 203
 smokers, 172
 three-stroke rule, 238
 decongestants, 143
 deforming brain, 93–94
 dehydration
 risk for white stroke, 56
 thiazide diuretics, 147
 dementia
 benefits, determining, 266
 described, 119–120, 330
 diagnosing, 125–129
 families, 297
 life, getting in order, 131
 preventing, 129–131
 prognosis, 120
 recognizing, 123–125
 small strokes and, 121–123
 depression
 alcohol consumption, 142
 case study, 62
 families, 297
 homecoming, 242–243
 left side of brain, 35
 recovery, 281–284
 diabetes
 checking after stroke, 211
 defined, 330
 future white strokes, preventing, 211
 monitoring, 99
 risk factors, 179–180
 stroke dementia, preventing, 130
 stroke risk, 171
 symptoms, TIA, 78
 testing for, 183
 thiazide diuretics, 147

diagnosis
 dementia, 125–129
 language problems, 191
 swallowing problems, 292
 white stroke, 64–65
 dialysis, 98
 diarrhea, 165
 diastolic blood pressure, 137, 138, 331
 Diazepam, 279–280
 diet
 cholesterol consumption, 154, 165
 drugs, 143
 exercise and, 327
 fecal incontinence, handling, 289
 high blood pressure, 141, 144–145
 right, 163
 supplements, 144
 digestive system, 152–153
 dining arrangements, nursing home, 271
 disabilities
 families, 297
 Medicare, 266
 discharge summary, 219–220
 dissection
 causes, 57–58
 described, 16, 17–18
 dissolving clot, 66
 diuretics, 146
 dizzy spells, 75–76
 doctor. *See also* neurosurgeon; second
 opinions
 homecoming checklist, 255
 importance of visiting after stroke,
 247–248
 medical attention, seeking, 78–79
 neurologist, 281, 332
 other problems, discussing, 248–249
 patient's role in appointments, 249
 physiatrist, 228
 radiologist, 114–115
 rehabilitation, factors considered by, 223
 stroke specialists, consulting, 25
 testing, 248
 transportation to appointments, 271
 draining brain, 95–96
 driving limitations
 alternatives, identifying, 286–287
 consequences, 285
 evaluation, 285–286

signs, reading, 284
 work, returning to, 253
 drug abuse, 172
 drugs, prescription. *See* medication
 dypiramide (Aggrenox), 176, 325

• E •

easygoing nature, loss of, 241
 eating habits, changing, 180–181
 eating right versus eating less, 162
 eddies, atrial fibrillation, 55
 educating families, 306
 EKG (electrocardiogram)
 atrial fibrillation, 55, 79
 described, 183, 331
 in emergency room, 200
 heart status, checking, 210
 elder law, 305
 embolic stroke. *See* white stroke
 embolism, 17
 embolus, 331
 Emergency Medical Service (EMS)
 scoop and run approach, 194
 stroke response, evaluating, 204, 312–313
 Emergency Medical Technicians
 (EMTs), 194
 emergency room
 complications, treating, 66
 going with patient, 192
 hospital, choosing, 195–197
 patient, advocating for, 197–198
 red strokes, treating, 203–205
 stroke strategy, understanding, 198–200
 symptoms of stroke, 190–193
 timing, 86
 transient stroke, 78
 treating red stroke (SAH), 111–112
 treatment plan, 23
 when to use, 189–190
 white strokes, treating, 201–203
 emotions, 35, 38, 294–295
 employment
 driving, 253
 limitations, 252
 social workers' interaction, 231
 speed, 251–252
 EMS (Emergency Medical Service)

scoop and run approach, 194
 stroke response, evaluating, 204, 312–313
 EMTs (Emergency Medical
 Technicians), 194
 endarterectomy, carotid, 209, 330
 energy, 241
 enzymes, 74, 153
Estate Planning For Dummies (Simon and
 Caverly), 306
 estrogen
 risk for white stroke, 55–56
 stroke risk, 172, 211–212
 evidence, collecting, 183–184
 executive function, 296
 exercise
 high blood pressure, lowering, 144–145
 injured brain cells, 225
 muscles and brain, 327–328
 obesity, 181–182
 stroke dementia, preventing, 129
 experts, connecting with best, 25
 eye pain, 104

• F •

face
 brain portion controlling, 33
 pain, 104
 family
 affairs, getting in order, 305–306
 caregiver, caring for, 299–301
 dementia, recognizing, 124, 126, 131, 297
 depression, handling, 242–243, 297
 disabilities and, 297
 educating, 306, 311
 finances, 303–304
 help, finding, 25–26
 importance of caring for, 293–294
 life support, handling, 98–99
 moving patient, 218
 nursing home proximity to, 268
 personality change, dealing with, 296
 rehabilitation, 231–232
 seeking help, 297–298
 of smokers, 173
 strengths, 306
 stress, signs of, 295–296
 stroke's affect on, 294–295
 team work, 301–303

- family history
 - high blood pressure, 140
 - stroke, 170
 - fast-food fries, 141
 - fasting, pre-cholesterol test, 159–160
 - fat
 - cholesterol and triglycerides, 153–155
 - consequences, 151–152
 - diet and cholesterol control, 162–164
 - in digestive system, 152–153
 - HDL, LDL and other transporters, 155–157
 - risk for white stroke, 54
 - as slow-burning fuel, 152
 - statin drugs, 164–167
 - testing, 157–161
 - fatigue, 252
 - fecal incontinence
 - control, regaining, 289–290
 - nursing home solution, 290
 - prevalence, 288
 - females
 - cerebellum, 31
 - estrogen risks, 56, 211–212, 245
 - stroke risk, 171
 - finances
 - assisted-living facilities, 270–271
 - discussing with family, 304
 - nursing home, 267
 - professional guidance, 303–304
 - rehabilitation, 232–233
 - finding
 - aneurysm, 112–113
 - narrowed artery, 208–209
 - fine-motor control, 123
 - finger
 - difficulty controlling, 189
 - vessels in, 55
 - fleeting blindness (amaurosis fugax), 72
 - food, nursing home, 269
 - 4th ventricle, 31
 - fracture, 202
 - friends, sharing knowledge with, 311
 - frontal lobes, 288, 290
 - fruits, 141
 - fungal infections, medications treating, 167
 - fusing joints, 280–281
 - future strokes, preventing, 226
- **G** •
- gemfibrozil (Lopid), 166, 167
 - glucose, 50, 152, 179
 - goals, prevention, 184–185
 - government organizations, 310
 - grapefruit juice, 166
 - guilt, nursing home or assisted-living facility placement, 258
- **H** •
- hand, weakness of, 76
 - Handel, George Frideric (composer), 254, 319–320
 - hands
 - brain portion controlling, 33
 - fine-motor control, 123
 - hardening clots, 51–52
 - hardening of the arteries
 - angiogram finding, 209
 - dementia, 121
 - described, 156–157, 329
 - dietary fat, 54
 - high blood pressure with, 136–137, 139
 - risk for white stroke, 54
 - surgery, 54, 81, 209
 - ultrasound testing for, 183
 - white stroke and, 16–17, 72
 - HDL (high-density lipoprotein)
 - cholesterol test, interpreting, 161
 - described, 155, 331
 - head trauma, 92
 - headache
 - aneurysm, revealing, 117
 - migraine, 76, 104, 205
 - real-life examples, 12
 - subarachnoid hemorrhage, 20, 87, 107
 - health maintenance organizations (HMOs), 233
 - health policy, 315
 - healthy future, prospect for, 186
 - hearing, 33
 - hearing cortex, 35
 - heart
 - checking, 210
 - exam, 80
 - failure, 178

- high blood pressure, 139
 - monitoring, 217
 - red stroke (SAH), 108
 - rhythm, 111
 - valve disease, 176
- heart beat, irregular
 - checking for, 79, 248
 - described, 175, 330
 - medication, 89
 - risk for white stroke, 55–56
 - treating, 326
- heart disease
 - atrial fibrillation, 175
 - attack, 175–176
 - clotting, avoiding, 176–178
 - managing, 99
 - medication, 143
 - status, checking, 210
 - stroke risk, 171
 - TPA, usefulness of, 202
 - valve disease, 176
- help, asking for, 25–26, 299
- helping someone else, 191–192
- hemorrhagic conversion of infarction, 89, 331
- hemorrhagic stroke. *See* red stroke
- Heparin, 89
- high blood pressure
 - with atherosclerosis, 136–137
 - averaging, 138
 - brain hemorrhage, 86
 - checking, 139
 - defining, 137–138
 - described, 331
 - lowering with medication, 129, 131, 145–150
 - lowering without medication, 144–146
 - measuring, 137
 - medication causing stroke, 139
 - prevalence, 135
 - red stroke, 87–88, 211
 - risk factors, 140–144
 - sudden and severe increase, 92
 - treating, 324–325
 - white stroke and, 53–54
- high-density lipoprotein (HDL)
 - cholesterol test, interpreting, 161
 - described, 155, 331
- HMOs (health maintenance organizations), 233
- holes in brain, 17, 48, 71, 331
- homecoming
 - avoiding future strokes, 238
 - basics, 254
 - daily checklist, 255
 - depression, 242–243
 - doctor's visit checklist, 255
 - medical therapy, 244–246
 - new habits, forming, 254–255
 - organization, 253–254
 - outlook, 237–238
 - personality changes, 240–241
 - physician, visiting, 247–249
 - rehabilitation therapy, 249–251
 - rehabilitation versus, 222–223
 - rewiring brain, 239
 - smoking, 247
 - spouse or caregiving, impact on, 243–244
 - timetable, recovery, 238–239
 - work, returning to, 251–253
- homocysteine, 159
- hospital
 - choosing, 195–197
 - primary stroke center designation, 196
 - hospital insurance, Medicare Part A, 232
 - hospital treatment
 - blood clots in legs, 66, 214–216, 215–216
 - brain swelling, monitoring, 213–214
 - discharge summary, 219–220
 - heart monitoring, 217
 - pressure sores, avoiding, 218
 - prognosis, 207
 - psychological, 283–284
 - seizures, 214
 - speed, critical need for, 23
 - stay, getting most out of, 219
 - stomach ulcers, bleeding, 218
 - swallowing difficulties, 66, 216–217
 - testing, 208–210
 - transient stroke, 78
 - white stroke, reducing risk of another, 210–211
- hospital treatment, red stroke (ICH)
 - draining brain, 95–96
 - drugs to reduce swelling, 95
 - hyperventilating, 95
 - repeat, reducing risk of, 211–212
 - unconsciousness, 94

hospital treatment, red stroke (SAH)
 artery spasm, 115–116
 brain swelling, relieving, 115
 emergency, 111–112
 finding aneurysm, 112–113
 prognosis, 110
 sealing off aneurysm, 113–115
 severity, measuring, 111
 hospital treatment, white stroke. *See also* TPA
 complications, 66
 dissolving clot, 66
 preventing recurrence, 67
 hydration, 327
 hydrocephalus, 108, 115
 hyperaldosteronism, 143
 hypertension, 53–54, 171, 331
 hypertensive crisis, 92
 hyperthyroidism, 127, 143
 hyperventilating, 95

• I •

ICH (intracerebral hemorrhage)
 hospital treatment, 94–99
 prevention, 99
 reasons for, 86–92
 identifying stroke type, 199
 imaging. *See also* CT scan; MRI scan;
 ultrasound
 angiogram, 43
 dementia, diagnosing, 126
 PET scan, 44
 impotence, blood pressure medications
 and, 146
 impulsive behavior, 38
 incapacitation, 191
 incontinence
 discussing with doctor, 248
 fecal, 288–290
 prevalence, 287–288
 urinary, 290–291
 independence, 241
 infarction, 17, 48, 71, 331
 infections, 91, 98
 information, gathering, 323–324
 injury, white stroke
 brain bleeding, 60
 brain swelling, 59–60

response, 40–41
 seizures, 60
 INR (International Normalized Ratio), 177
 insulin, 77, 179
 insurance, medical
 inadequacy, 266
 rehabilitation, 233
 intelligence, 296
 intracerebral, 331
 intracerebral hemorrhage (ICH)
 hospital treatment, 94–99
 prevention, 99
 reasons for, 86–92
 intracranial, 331
 irregular heart beat
 checking for, 79, 248
 described, 175, 330
 medication, 89
 risk for white stroke, 55–56
 treating, 326
 ischemia, 48, 332
 ischemic stroke, 16, 48, 332. *See also* white
 stroke

• J •

job, returning to
 driving, 253
 limitations, 252
 social workers' interaction, 231
 speed, 251–252
 Joint Commission on Accreditation of
 Healthcare Organizations, 204

• K •

Kesey, Ken (novelist), 320
 kidneys
 blood clotting, 98
 high blood pressure, 139, 142
 table salt, 141
 Knight, J. (*Wills, Probate, and Inheritance
 Tax For Dummies*), 306

• L •

laboratories, ultrasound, 184
 lacunar stroke. *See* white stroke

- language problems
 - brain portion, 33
 - diagnosing stroke, 191
 - nursing homes handling, 259
 - recognizing, 125
 - recovery pace, 238–239
 - speech therapy, 230
 - large vessel stroke. *See* white stroke
 - lateral ventricles, 31
 - Latino Americans, 171
 - LDL (low-density lipoprotein)
 - cholesterol test, interpreting, 161
 - defined, 332
 - described, 156
 - statin medicines, effectiveness of, 326
 - learning, 34
 - left and right hemispheres, 35–36
 - leg muscles
 - brain portion controlling, 33
 - not working, 14
 - legs
 - blood clots in, 66
 - crossing, raising blood pressure by, 139
 - falls asleep, 75
 - sudden numbness or weakness, 191
 - length of time, rehabilitation, 223–224
 - Lescol, 165
 - life, getting in order, 131
 - life support
 - hospital treatment, red stroke, 98–99
 - hyperventilation, 95
 - subarachnoid hemorrhage, 112
 - unconsciousness, 94
 - limb, jerking, 77
 - limitations, caregiving, 298
 - lining, blood vessel, 16, 17–18, 57–58
 - Liorisal (Baclofen), 279
 - lipase, 153
 - lipid
 - described, 99, 332
 - diet controlling, 163
 - stroke dementia, preventing, 130
 - stroke risk, 171
 - Lipitor, 165
 - lipoproteins, 155
 - liver
 - clotting failure, 98
 - fat, breaking down, 152–153
 - side effects of statin drugs, 165, 166
 - local hospitals, working with, 311–312
 - local stroke clubs and support groups, 309–310
 - location, nursing home, 268
 - logical half of brain, 35
 - Lopid (gemfibrozil), 166, 167
 - low blood pressure
 - measuring, 138
 - medications going too far, 145–146
 - low blood sugar, 77
 - low-density lipoprotein (LDL)
 - cholesterol test, interpreting, 161
 - defined, 332
 - described, 156
 - statin medicines, effectiveness of, 326
 - low-income medical coverage, 233, 266
 - lumbar puncture, 128, 200, 333
 - lung infection, 216
- M ●
- magnetic resonance angiography (MRA), 43
 - magnetic resonance imaging scan (MRI scan)
 - aneurysm, finding, 112
 - AVMs, 90
 - brain hemorrhage, 93
 - dementia, 126
 - described, 42–43, 332
 - in emergency room, 200
 - physician's visit, 80
 - white stroke, diagnosing, 64
 - maintenance plan, 328
 - malnutrition, 216
 - Mayer, Jacqueline (Miss America), 319
 - mechanical heart valve, 176
 - Medicaid, 233, 266
 - medical and professional staff, meeting, 268–269
 - medical care, 300
 - medical insurance
 - inadequacy, 266
 - rehabilitation, 233
 - medical power of attorney, 131
 - medical therapy, 244–246
 - Medicare, 232, 264–265

- medication
 - atrial fibrillation, 175
 - bladder infection, 290
 - blood clots, avoiding, 80–81
 - blood-thinning, 89
 - causing high blood pressure, 139, 143–144
 - cholesterol test, affecting, 159
 - continuing, 247
 - depression, 242, 283–284
 - diabetes, 179
 - doctor's staff, pressing, 245
 - lowering high blood pressure, 145–150
 - muscle spasticity, 279–280
 - at nursing home or assisted-living facility, 274
 - quitting smoking, 173
 - risk for white stroke, 57
 - swelling, reducing, 95
 - TPA, 24
 - Mediterranean diet, 163
 - medium (branches of main trunk arteries blocked) white stroke, 62–63
 - memory loss, 123, 124–125, 296, 297
 - men
 - high blood pressure medications, 146
 - stroke risk, 171
 - Messiah* (Handel), 319–320
 - Mevacor, 165
 - microaneurysm, 88, 332
 - microatheroma, 121
 - migraine, 76, 104, 205
 - mobility issues, 289–290
 - money issues
 - assisted-living facilities, 270–271
 - discussing with family, 304
 - nursing home, 267
 - professional guidance, 303–304
 - rehabilitation, 232–233
 - MRA (magnetic resonance angiography), 43
 - MRI (magnetic resonance imaging) scan
 - aneurysm, finding, 112
 - AVMs, 90
 - brain hemorrhage, 93
 - dementia, 126
 - described, 42–43, 332
 - in emergency room, 200
 - physician's visit, 80
 - white stroke, diagnosing, 64
 - muscle spasticity
 - braces or splits, wearing, 278
 - brain injury and uncontrolled muscles, 276–277
 - contractures, 277
 - described, 276, 333
 - medication, 279–280
 - physical therapy, 278
 - recognizing and treating, 277–278
 - reflexes, controlling, 278–279
 - surgery, 280–281
 - muscles
 - aching, 165
 - exercising, 327–328
 - injury, 166
 - rehabilitation, 224
 - muscles, leg
 - brain portion controlling, 33
 - not working, 14
- N ●
- narrowed artery, 209–210
 - National Association for Incontinence, 291
 - National Institute of Health Stroke Scale (NIHSS), 195
 - national organizations, 310
 - National Stroke Association, 319
 - nausea, 165
 - Neal, Patricia (actress), 321
 - neck pain, 104
 - nerves, 32–34
 - nervousness, 191
 - neurologist, 281, 332
 - neurosurgeon
 - age, 110–111
 - clipping or coiling aneurysm, 113–115
 - second opinion, seeking, 212
 - new habits, forming, 254–255
 - nicotine patches and gum, 174
 - NIHSS (National Institute of Health Stroke Scale), 195
 - 911, calling, 13–14, 192, 193–195, 238
 - non-brain symptoms, 14
 - nonmedical staff, 268
 - numbness
 - limbs, 75, 189, 191
 - pressure sores, avoiding, 218
 - nurses, 230–231, 268, 272

nursing home
 adjusting to life, 271–274
 care needs, defining, 260–264
 costs, investigating, 267
 fecal incontinence, 290
 game plan, 259–260
 grappling with decision, 258–259
 helping family choose, 257–258
 investigating, 269
 location, 268
 medical and professional staff, meeting,
 268–269
 Medicare payment, 265
 resources, determining, 264–267
 second opinion, getting, 270
 urinary incontinence, 291
 when to choose, 259

● 0 ●

obesity
 advertising, 181
 eating habits, changing, 180–181
 exercising, 181–182
 high blood pressure, 141, 144–145
 stroke dementia, preventing, 129
 stroke risk, 171
 weight gain, principle of, 180
 occlusive stroke. *See* white stroke
 occupational therapist
 defined, 332
 nursing home, 268
 rehabilitation, 229
 ophthalmologist, 287
 optimism, 26–27
 organization, homecoming, 253–254
 outlook, homecoming, 237–238
 outpatient services, 232
 outpatient therapy, 250
 overactive muscles, 276
 oxygen needs
 blood vessels, carrying, 36–37
 white stroke, 50–51

● p ●

pain, lack of, 13
 pancreas, 153
 patience, 240

patient, advocating for
 in emergency room, 197–198
 at nursing home or assisted-living facility,
 273–274
 pear-shaped figures, 157
 personality change
 families, dealing with, 296
 homecoming, 240–241
 testing for depression, 282
 PET (positron emission tomography)
 scan, 44
 pheochromocytoma, 143
 photographing nursing home staff, 272
 physiatrist, 228
 physical activity, 225
 physical therapist, 332
 physical therapy
 monitoring, 274
 muscle spasticity, 278
 nursing home, 268
 rehabilitation, 228–229
 physician. *See also* neurosurgeon; second
 opinions
 homecoming checklist, 255
 importance of visiting after stroke,
 247–248
 medical attention, seeking, 78–79
 neurologist, 281, 332
 other problems, discussing, 248–249
 patient's role in appointments, 249
 physiatrist, 228
 radiologist, 114–115
 rehabilitation, factors considered by, 223
 stroke specialists, consulting, 25
 testing, 248
 transportation to appointments, 271
 pia mater, 20, 102
 pictures of the brain. *See* imaging
 plans, disorganization, 123
 plaque
 dementia, 121
 described, 16–17, 332
 high blood pressure and, 136–137
 kidneys, 142
 sound, 52
 surgery to remove, 54, 81, 209
 transient stroke mechanics, 72
 vessel wall bleeding, 52
 platelets, 51–52, 176

- Plavix (clopidogrel), 129, 176, 325
 pneumonia, 216, 291–292
 positron emission tomography (PET)
 scan, 44
 potassium
 ACE inhibitors raising, 149
 high blood pressure, 141
 medication depleting, 147
 power of attorney, 131
 Pravachol, 165
 predicting red stroke (SAH), 116–117
 pre-hypertensive blood pressure, 138
 prescription drugs
 atrial fibrillation, 175
 bladder infection, 290
 blood clots, avoiding, 80–81
 blood-thinning, 89
 causing high blood pressure, 139, 143–144
 cholesterol test, affecting, 159
 continuing, 247
 depression, 242, 283–284
 diabetes, 179
 doctor’s staff, pressing, 245
 lowering high blood pressure, 145–150
 muscle spasticity, 279–280
 at nursing home or assisted-living
 facility, 274
 quitting smoking, 173
 risk for white stroke, 57
 swelling, reducing, 95
 TPA, 24
 pressure sores, avoiding, 218
 preventing
 dementia, 129–131
 recurrence of white stroke, 67
 primary stroke centers, 196
 problems with nursing home staff, 273
 problem-solving skills, 124, 296
 professional financial guidance, 303–304
 prognosis
 dementia, 120
 hospital treatment, 207
 quitting smoking, 174–175
 red stroke (SAH), 110
 promises, decision-making about nursing
 home, 258
 prothrombin time or protime (PT), 177
 Prozac, 284
 psychological changes, 281–284
 psychological tests, 127
 pulmonary embolism, 177, 215
 pulse rate, 149, 249
 pupil, large in one eye, 104
- *Q* •
- quitting smoking, 174–175
- *R* •
- race, stroke risk and, 171
 radiologist, 114–115
 range-of-motion therapy, 251
 reading, 33
 rebleeding, SAH red stroke, 107
 recalling words, 189
 recovery
 brain cells, exercising, 25
 depression and other psychological
 changes, 281–284
 driving limitations, 284–287
 experts, connecting with best, 25
 help, asking for, 25–26
 incontinence, 287–291
 muscle spasticity, 276–281
 swallowing difficulties, 291–292
 telling about your own, 321
 “use it or lose it” principal, 26
 recurrence, stroke, 171
 red stroke
 bleeding within brain, 19
 CT image, 42
 described, 15, 85–86, 333
 emergency room treatment, 203–205
 repeat, reducing risk, 211
 scenarios, 21
 treatment, 24
 with white stroke, 21
 white stroke, causing, 58–59
 red stroke (ICH)
 hospital treatment, 94–99
 prevention, 99
 reasons for, 86–92
 red stroke (SAH)
 arteries and, 103
 behavioral changes, 109–110
 bleeding around brain, 20

- brain packaging, 102–103
 - brain swelling, 108
 - CT scan, 42
 - described, 20, 101–102, 333
 - heart problems, 108
 - identifying, 199
 - outlook following, 110
 - predicting, 116–117
 - rebleeding, 107
 - red strokes, 92
 - repeat, reducing risk, 211–212
 - rupture, 104–107
 - seizures, 109
 - spinal tap, 201
 - transcranial doppler test, 210
 - treating, 110–116, 203
 - warning signs, 104
 - white stroke causing, 58
 - white strokes following, 109
 - reflexes, controlling, 278–279
 - regional organizations, 310
 - rehabilitation
 - backsliding, 227
 - brain recovery, 225–226
 - curriculum, 227–228
 - doctor, factors considered by, 223
 - family and other caregivers, 231–232
 - financing, 232–233
 - future strokes, preventing, 226
 - graduating, 233–234
 - at home, 249–251
 - home versus, 222–223
 - length of time, 223–224
 - muscles, maintaining, 224
 - need for, 221–222
 - nurses, 230–231
 - occupational therapists, 229
 - physical therapists, 228–229
 - rewiring brain, 41
 - social workers, 231
 - speech therapists, 230
 - substitution, 224–225
 - success story, 226
 - removing clot, 96–97
 - research, 314–315
 - residents, nursing home, 270
 - resources, determining, 264–267
 - response time, 14
 - response to injury, 40–41
 - results, cholesterol test, 160–161
 - retina, 72, 139
 - retirement, 252
 - rewiring brain, 40–41, 239
 - right hemispheres, 35–36
 - risk factors
 - accepting unchangeable, 170–171
 - angiogram, 43, 112
 - AVMs, 90
 - benefits of treating, 182
 - checklist, 184
 - cholesterol levels, high, 167
 - clinical trials, volunteering, 315
 - clot, removing, 96
 - dementia, 128–129
 - diabetes, 179–180
 - eliminating, reducing, or treating, 171–172
 - evidence, collecting, 183–184
 - goals, setting prevention, 184–185
 - healthy future, prospect for, 186
 - heart and vascular disease, 175–178
 - high blood pressure, 139–144
 - obesity, 180–182
 - tobacco, 172–175
 - TPA, 24
 - white stroke, 53–57
 - Rivotril (clonazepam), 279–280
 - rules, nursing home, 269
 - rupture, red stroke (SAH), 104–107
- S ●
- SAH (subarachnoid hemorrhage)
 - arteries and, 103
 - behavioral changes, 109–110
 - bleeding around brain, 20
 - brain packaging, 102–103
 - brain swelling, 108
 - CT scan, 42
 - described, 20, 101–102, 333
 - heart problems, 108
 - identifying, 199
 - outlook following, 110
 - predicting, 116–117
 - rebleeding, 107
 - red strokes, 92
 - repeat, reducing risk, 211–212
 - rupture, 104–107
 - seizures, 109

- SAH (subarachnoid hemorrhage) (*continued*)
 - spinal tap, 201
 - transcranial doppler test, 210
 - treating, 110–116, 203
 - warning signs, 104
 - white stroke causing, 58
 - white strokes following, 109
- salt, 141
- saturated fats, 154, 333
- saying words, 189
- scan, showing white stroke, 49
- schoolchildren, educating, 313
- scoop and run EMS approach, 194, 312
- sealing off aneurysm, 113–115
- second opinions
 - dementia, diagnosing, 128
 - nursing home, 270
 - stroke treatment, 212
- second-hand smoke, 173
- sedimentation rate, 127
- seizures
 - drug withdrawal, sudden, 279
 - evaluating, 76–77
 - hospital treatment, 214
 - injury, white stroke, 60
 - subarachnoid hemorrhage, 109
- selective serotonin reuptake inhibitors (SSRIs), 284
- senses, 32, 34–35
- Serzone antidepressant, 167
- severity, measuring red stroke (SAH), 111
- sex problems, 146, 248
- shaking, 191
- shoulder blades, sudden pain, 104
- shrinking brain, 122–123
- side effects
 - beta blockers, 149
 - blood thinners, 89
 - statin drugs, 165–166
- sight
 - brain portion controlling, 33
 - dementia, 124
 - diagnosing stroke, 191
 - fleeting blindness, 72
 - injury, response to, 40
- signs of stroke, 13
- Simon, Jordan (*Estate Planning For Dummies*), 306
- situation, evaluating, 298
- sleep
 - dehydration following, 56
 - low blood pressure, measuring, 138
 - seizures in, 77
 - stroke during, 193
- sleep apnea, 143
- small (cross circulation minimizing) white stroke, 63–64
- small injuries, 218
- small strokes, 121–123
- small vessel stroke. *See* white stroke
- smell, 35
- smoking tobacco
 - blood and, 172–173
 - carotid artery, 327
 - estrogen and, 56
 - homecoming, 247
 - prognosis for quitting, 174–175
 - risk for white stroke, 55
 - stroke dementia, preventing, 130
- social activities, nursing home, 269
- social engineering, 174
- social pressure, 173
- social worker
 - discussion, preparing, 265, 333
 - recovery group, starting, 313
 - rehabilitation, 231
 - resources, determining, 264
- sodium, 141, 333
- sound wave picture. *See* ultrasound
- spasticity, muscle
 - braces or splits, wearing, 278
 - brain injury and uncontrolled muscles, 276–277
 - contractures, 277
 - described, 276, 333
 - medication, 279–280
 - physical therapy, 278
 - recognizing and treating, 277–278
 - reflexes, controlling, 278–279
 - surgery, 280–281
- speech problems
 - brain portion controlling, 33
 - diagnosing stroke, 191
 - nursing homes handling, 259
 - recognizing, 125
 - recovery pace, 238–239
 - speech therapy, 230

- speech therapist
 - rehabilitation, 230
 - swallowing difficulties, diagnosing, 217, 292
- speed, critical need for, 23
- spinal cord, 31
- spinal tap
 - defined, 333
 - dementia, diagnosing, 128
 - in emergency room, 200, 201
- splints, wearing, 278
- spouse, impact on, 243–244. *See also* caregiver
- SSRIs (selective serotonin reuptake inhibitors), 284
- stable nature, 241
- staff, handling, 273
- state organizations, 310
- statins/statin drugs
 - continuing, 247
 - described, 333
 - pros and cons of using, 164–167
- stay, getting most out of hospital, 219
- stem, brain
 - described, 31, 330
 - heart, monitoring, 217
 - subarachnoid hemorrhage, 111
- stenosis, 81, 209–210
- stents, 82
- steroids, 143
- stiffness, 104
- stimulant drugs, 92
- stomach ulcers, bleeding, 218
- stress
 - families, signs of, 295–296
 - personality changes, 241
 - risk for white stroke, 56–57
- stroke
 - defined, 333
 - protocols, 195
 - recovery group, starting, 313
 - strategy, understanding, 198–200
 - support groups, 244, 266, 299
- Stroke Belt Consortium, 310
- stroke risk
 - controllable factors, 22–23
 - rates, 21–22
 - unchangeable factors, 22
- subarachnoid hemorrhage (SAH)
 - arteries and, 103
 - behavioral changes, 109–110
 - bleeding around brain, 20
 - brain packaging, 102–103
 - brain swelling, 108
 - CT scan, 42
 - described, 20, 101–102, 333
 - heart problems, 108
 - identifying, 199
 - outlook following, 110
 - predicting, 116–117
 - rebleeding, 107
 - red strokes, 92
 - repeat, reducing risk, 211–212
 - rupture, 104–107
 - seizures, 109
 - spinal tap, 201
 - transcranial doppler test, 210
 - treating, 110–116, 203
 - warning signs, 104
 - white stroke causing, 58
 - white strokes following, 109
- subarachnoid space, 103, 333
- substitution
 - quitting smoking, 174
 - rehabilitation, 224–225
- supplementary medical insurance, Medicare Part B, 232
- support groups, 244, 266, 299
- surgery
 - clot, removing, 96–97
 - following TIA, 81
 - muscle spasticity, 280–281
 - plaques, removing, 54, 81, 209
- swallowing difficulties
 - discerning, 230
 - recovery, 291–292
 - treating, 66, 216–217
- swelling, brain
 - deformation, 93–94
 - injury, white stroke, 59–60
 - monitoring, 213–214
 - red stroke (SAH), 108
 - relieving, 115
 - ventricles, 32
- swelling, brain edema, 108
- symptoms, 190–193
- synapses, 32
- systolic blood pressure, 137, 334

• T •

- table salt, 141
- taste, 35
- tea, 246
- team work
 - families, 301–303
 - TPA use, 66
- telephone numbers, 272
- television, 181
- tests/testing
 - after headaches, 105
 - aneurysm, 117
 - cerebral angiogram, 209
 - for depression, 282–283
 - in emergency room, 199–201
 - fat, 157–161
 - hospital treatment, 208–210
 - for liver damage from statins, 166
 - physician, 248
 - for swallowing problems, 217
 - transcranial doppler test, 209–210
 - ultrasound, 208–209
 - vessel causing symptoms, 38
- therapy, nursing home, 274
- therapy, rehabilitative
 - backsliding, 227
 - brain recovery, 225–226
 - curriculum, 227–228
 - doctor, factors considered by, 223
 - family and other caregivers, 231–232
 - financing, 232–233
 - future strokes, preventing, 226
 - graduating, 233–234
 - at home, 249–251
 - length of time, 223–224
 - muscles, maintaining, 224
 - need for, 221–222
 - nurses, 230–231
 - occupational therapists, 229
 - physical therapists, 228–229
 - rewiring brain, 41
 - social workers, 231
 - speech therapists, 230
 - substitution, 224–225
 - success story, 226
- thiazide diuretics, 147
- 3rd ventricle, 31
- three-stroke rule, 238
- thrombotic stroke. *See* white stroke
- thrombophlebitis, 215
- thyroid disease, 127, 143
- TIA (transient ischemic attack)
 - described, 18, 69–70, 334
 - identifying, 199
 - mechanics, 72–74
 - recognizing, 74–77
 - reducing risk for bigger stroke following, 80–82
 - responding to, 78–80
 - symptoms and brain injury, 72
 - TPA contraindicated, 202
 - as wake-up call, 82–83
- timetable, recovery, 238–239
- timing
 - improving condition, 237–238
 - physical therapy, 229
 - plugged artery, length of, 70
 - stroke, help for, 47–48, 193
- tissue plasminogen activator. *See* TPA
- Tizanidine, 280
- tobacco, smoking
 - blood and, 172–173
 - carotid artery, 327
 - estrogen and, 56
 - homecoming, 247
 - prognosis for quitting, 174–175
 - risk for white stroke, 55
 - stroke dementia, preventing, 130
- today's needs, planning for, 263
- tools, special, 224–225
- tooth infection, 91
- total cholesterol, 161
- touch, 35, 40
- TPA (tissue plasminogen activator)
 - described, 24, 39–40, 334
 - effectiveness, 66
 - patients who can benefit from, 202
 - risks, 201
 - success story, 320
- trans fats, 141, 154, 325–326
- transcranial doppler test, 209–210
- Transcranial Doppler ultrasound, 44
- transient ischemic attack (TIA)
 - described, 18, 69–70, 334
 - identifying, 199
 - mechanics, 72–74
 - recognizing, 74–77

reducing risk for bigger stroke following, 80–82
 responding to, 78–80
 symptoms and brain injury, 72
 TPA contraindicated, 202
 as wake-up call, 82–83
 transient monocular blindness, 72
 transportation to appointments, 271
 travel, 252
 treating red stroke (ICH)
 draining brain, 95–96
 drugs to reduce swelling, 95
 hyperventilating, 95
 repeat, reducing risk of, 211–212
 unconsciousness, 94
 treating red stroke (SAH)
 artery spasm, 115–116
 brain swelling, relieving, 115
 emergency, 111–112
 finding aneurysm, 112–113
 prognosis, 110
 sealing off aneurysm, 113–115
 severity, measuring, 111
 treating white stroke. *See also* TPA
 complications, 66
 dissolving clot, 66
 preventing recurrence, 67
 treatment
 blood clots in legs, 215–216
 psychological, 283–284
 speed, critical need for, 23
 trembling, 191
 trials, clinical
 list of, 314
 volunteering for, 315
 triglyceride, 154–155, 161, 334
 tubes
 clot, removing, 96
 feeding, 217, 292
 hydrocephalus, 115
 life support, 94
 tumors, brain, 91

• U •

ultrasound
 carotid, 326–327
 described, 43–44
 testing, 183, 208–209
 white stroke, diagnosing, 65

unchangeable risks, 22
 unconsciousness
 red stroke, 94
 subarachnoid hemorrhage, 107
 uncontrolled muscles, 276–277
 understanding, loss of, 189
 unsaturated triglycerides, 154
 urinary incontinence, 290–291
 “use it or lose it” principal, 26

• V •

Valium, 279–280
 vascular dementia
 benefits, determining, 266
 described, 18, 119–120, 330, 334
 diagnosing, 125–129
 families, 297
 infarction, 71
 life, getting in order, 131
 preventing, 129–131
 prognosis, 120
 recognizing, 123–125
 small strokes and, 121–123
 smoking and, 173
 vascular disease
 atrial fibrillation, 175
 attack, 175–176
 clotting, avoiding, 176–178
 heart attack, 175–176
 heart valve disease, 176
 managing, 99
 stroke risk, 171
 TPA, usefulness of, 202
 vasospasm, 59, 212
 vein, 334
 ventricles of the brain
 CT scan showing infarct, 49
 described, 31–32, 334
 draining, 95–96
 ventriculostomy, 96
 vertebral artery dissection, 58
 vertigo, 75
 very-low-density lipoprotein (VLDL), 156
 vessel, blood. *See also* dissection
 abnormalities, 90
 brain workings, 36–37
 damage, 16–17

vessel, blood (*continued*)
 lining, splitting, 16, 17–18
 scarring, 122
 small strokes and, 121
 vision
 brain portion controlling, 33
 dementia, 124
 diagnosing stroke, 191
 fleeting blindness, 72
 injury, response to, 40
 visiting time, 274
 visual cortex, 34–35
 visualization, smoking cessation, 174
 Vitamin K, 98, 177
 VLDL (very-low-density lipoprotein), 156
 volunteering for stroke research, 315

• W •

walking problems, 191
 warfarin
 described, 334
 details, paying attention to, 326
 hemorrhagic conversion of infarction, 89
 pulmonary embolism, 177–178
 red strokes, preventing, 99
 warning sign, transient stroke as, 78, 83
 Washington University, 314
 water pills, 147
 water, problem drinking, 217
 weak spots in arteries, 103
 weakness, 75, 191
 weight gain, principle of, 180
 Wellbutrin, 173
 white blood cells, 40
 white stroke. *See also* TIA
 big (major artery blocked), 61–62
 bleeding into vessel wall, 52
 blood-vessel damage, 16–17
 bruit sound, 53
 causes, 57–59
 clotting, 51

CT image, 42
 described, 15, 47–48, 334
 diagnosing, 64–65
 emergency room treatment, 201–203
 following SAH red stroke, 109
 hardening clots, 51–52
 hospital treatment, 210–211
 injuries, 59–60
 jargon, 48–49
 lining, blood vessel splits, 17–18
 medium (branches of main trunk arteries
 blocked), 62–63
 oxygen needs, 50–51
 red stroke, leading to, 88–89
 red stroke with, 21
 risk factors, 53–57
 scan showing, 49
 scenarios, 21
 small (cross circulation minimizing),
 63–64
 TIA, 18
 treating, 24, 65–67
 whoosh sound, hearing, 53, 80, 326–327
*Wills, Probate, and Inheritance Tax For
 Dummies* (Knight), 306
 Wilson, Woodrow (president), 317–318
 wine, 130
 withdrawal, sudden, 279
 women
 cerebellum, 31
 estrogen risks, 56, 211–212, 245
 stroke risk, 171
 work, returning to
 driving, 253
 limitations, 252
 social workers' interaction, 231
 speed, 251–252

• Z •

Zocor, 165
 Zyban, 173

