

INDEX

Note: Figures and Tables are indicated by *italic page numbers*

- active set algorithms 135, 136
- air quality (in environmental studies) 320, 321
 - see also* ALERT data; SCENES data
- ALERT data 318
 - chemical variables studied 319
 - interpretation of 322
- algebra 32–4
- algorithms 111–44
 - avoidance of numerical problems 229
- alternating least squares algorithms 111
 - benefits 112
 - for PARAFAC model 113–19
 - for Tucker1 model 124
 - for Tucker2 model 123–4
 - for Tucker3 model 121–2
- analysis of variance (ANOVA)
 - of data with complex interactions 340–6
 - in examples
 - designed data examples 324, 334–5
 - liquid chromatography example 312
 - modified-atmosphere packaging example 342–3
 - multiplicative approach 340–6
 - two-way 258
 - three-way array resulting from 259
 - see also* GEMANOVA
- analyte of interest, meaning of term 278
- ANOVA *see* analysis of variance
- anthropogenic loading, in VENICE data model 314
- applications 257–349
 - analysis of variance of data with complex interactions 340–6
 - chromatography 302–12
 - curve resolution of fluorescence data 259–75
 - exploratory analysis of designed data 323–40
 - exploratory analysis in environmental sciences 312–23
 - multi-way regression 285–8
 - process chemometrics 288–302
 - second-order calibration 276–85
- arctic air quality data 321
- array
 - definitions and properties 13–34
 - meaning of term 2
- associativity 32, 33
- autocorrelation plots 177
- autoscaling 234, 239
- bar charts/plots 176
 - examples 161, 168, 189, 190, 219, 330
 - scree plots presented as 159, 180, 186
- batch ester-synthesis examples
 - PARAFAC analysis 186–7, 189, 325–8
 - residual plots 216, 218
 - schematic view of data array 188
 - see also* BATCH example; SYNT example
- BATCH example 323–5
 - PARAFAC model 325–8
 - interpretation of 328–33
- batch polymerization example 291–301
 - background of example 291–3
 - building models 293–6
 - using models for on-line monitoring 296–9
 - process variables 292
 - typical time trajectories 290
- batch process monitoring 290–1
 - building models 293–6
 - fermentation example 301
 - on-line 296–9
 - polymerization example 291–3, 301–2
- batch processes 289
 - ‘ideal trajectory’ concept 291
- bilinear model, fitting 249–50

- bilinearity 21, 276–7
- biological axis, in VENICE data model 314
- biplots 176, 179, 206–13
 - compensation for number of objects and number of variables 207
 - factors affecting 206
 - fudge (zoom) factor in 207
 - example 211, 212
 - practical advice for 212–13
 - see also* joint plots
- block effect in PARAFAC model 324, 325, 326
 - cause in BATCH example 325
- block-scaling 235
- Box-Cox transformation 246
- bread sensory data 82–3, 286
 - choosing model for 155–6
 - influence analysis 172–3
 - multi-way regression 287–8
 - partial least squares regression 82–3
 - scree-like plot for Tucker3 model 162
 - sum of squared residuals 169
- 'broken stick' method 186, 329
- calibration 10, 46, 276
 - multivariate 10
 - second-order 10, 31, 106, 141, 257
 - example(s) 276–85
 - types 276
 - see also* second-order calibration
- Canonical Decomposition (CANDECOMP) model 58, 59
 - see also* PARAFAC model
- Carroll & Chang three-way model 58, 60
- cartesian plots 178–9
- categorization 6
- Cattell, Raymond 57, 58, 179
- centering of data
 - across first mode 225–6
 - across second mode 226–7
 - alternatives 254–5
 - compared with scaling 221, 244–5
 - definitions 222–7
 - geometric view 247–9
 - incorrect 251–4
 - interdependence with scaling
 - multi-way arrays 243–4
 - two-way arrays 238–9
 - method of operation 229–31
 - and missing data 232, 251
 - proper 244, 245
 - purpose 221
 - and rank reduction 229, 250–1
 - reason for wording 'across' 227
 - reasons for 228–9
 - rules for 228
 - situations when centering does not work
 - 232
 - three-way 240–1
 - as two-stage procedure for least squares fitting 231–2
 - two-way 228–32
- chemical rank 24
- chemical sources of variation 23–4, 31
- chromatography
 - multi-way analysis used in 257, 302–12
 - three-way array for 258
 - see also* normal-phase liquid chromatography; reversed-phase liquid chromatography
- classification methods 7
- closure
 - and rank 26–7
 - restriction due to 26
- color images 176
- column-centering 225–6, 229–30
- column-rank
 - of three-way arrays 30
 - of two-way arrays 22
- columns
 - in three-way arrays 14, 15, 15
 - in two-way arrays 2, 3
- comparative molecular field analysis (COMFA) 11
- component models 35–7, 53
 - cross-validation of 149–53
 - selecting number of components 156–7
 - determination from norm of components 160–1
 - residuals used 161–3
 - scree plots used 157–60
 - split-half analysis used 164–6
 - tools for 157–66
 - three-way one-block data analysis using 59–76
 - two-mode component analysis 45–6
 - two-way one-block data analysis using 35–46
 - see also* principal component analysis (PCA)
- components, number of
 - in component models 157–66
 - in examples 165–6, 264–5
 - selection of 156–66
 - verifying 165–6, 267–8
 - in regression models 166
- computational correctness 145
- conceptual aspects (of multi-way data analysis) 1–5
- confirmatory analysis 7
- constrained Tucker3 model 95
 - example 95–6
 - number of parameters 97
 - uniqueness 105–6
- continuous processes 289
 - monitoring of 289
- contour plots 176–7, 178
 - examples 182
- control chart(s) 10, 288, 289
- correlation spectroscopy 177
- CRAB data 215, 318, 319
 - chemical variables studied 319
 - interpretation of 321

- joint plot of data 215, 321
 - Tucker3 model 70, 321
- cross-validation 148–53
 - for determination of pseudo-rank 27
 - example comparing various approaches 152–3
 - expectation maximization approach 151–2, 153
 - in example(s) 268–70
 - leave-bar-out/leave-one-sample-out method 150–1, 153
 - in example(s) 270, 344
 - of multi-way component models 149–53
 - of multi-way regression models 153
 - of PARAFAC model in example(s) 268–70
- cumulative scree plot 184, 187
- curve resolution 8, 180, 302
 - examples
 - batch synthesis example 189
 - fluorescence spectroscopy examples 259–73, 274–5
 - mass spectrometry 274
 - UV-vis spectrometry 274
- cutoff values, in determination of pseudo-rank 27
- data base, meaning of term 1–2
- data collection 1
- data set(s)
 - features 6
 - large, visualization of 216, 218
- data transformations 245–7
 - see also* centering; scaling
- degeneracies
 - in PARAFAC models 89, 107–9
 - example 108–9
 - typical characteristic(s) 107
- dendrograms 177–8, 202–4
 - example 204
- designed data
 - BATCH example 323–5
 - interpretation of model 328–33
 - PARAFAC model 325–8
 - examples other than BATCH example 324, 334–5
 - model building and interpretation 336–9
 - exploratory analysis of 323–40
 - see also* BATCH example; MALT example; PEAT example; SEEDS example; SYNT example; XPS example
- diagnostics 146
- diffuse spectroscopy 246
- dimensionality vector, of three-way array 30
- direct trilinear decomposition (DTLD) 143
 - algorithm 143
 - second-order calibration problem solved using 282
- double-centering
 - three-way data 241
 - two-way data 227, 230
- dyad(s) 13, 17
- eigendecomposition 37
 - relationship with singular value decomposition 38
- ‘elbow’ method, example 329, 330
- emulsion batch polymerization, monitoring of 301–2
- environmental studies
 - exploratory analysis in 312–23
 - multi-way analysis used 70, 215, 312–23
 - second-order calibration example 280–3
 - typical three-way array for 259
 - VENICE data 313–14
 - interpretation of model 314, 316, 317
 - PARAFAC model 314, 315
- ester synthesis, batch process 186–9, 216, 218
- Euclidean distance(s) 205–6, 207
 - in example 209, 210, 211
- Euclidean plots 206
- expectation maximization
 - cross-validation using 151–2, 153
 - in example(s) 268–70
 - missing data handled using 151, 312
- explanatory validation 146
- exploratory analysis 7
 - in chromatography 302–12
 - of designed data 323–40
 - in environmental sciences 312–23
- fermentation example, batch process monitoring of 301
- first-order instruments 276
 - effect of interferent on calibration 278
- fit of model to data, and centering of data 229
- five-way data 11
- flow injection analysis, second-order calibration used in 284
- flow injection analysis example
 - constrained Tucker3 model for 95–6
 - PARAFAC model for 92–3
 - rank-deficiency
 - in three-way arrays 31–2
 - in two-way arrays 24–6
 - Tucker3 model for 92–3
- fluorescence spectroscopy
 - contour plots 182
 - curve resolution of data 259–73
- example
 - assessing validity of new samples 270–2
 - building of three-way model 262–70
 - cross-calibration of model 268–70
 - excitation-emission matrix (EEM) 263
 - finding scores of new data 270
 - low excitation wavelengths excluded 265–6
 - outlying samples and variables assessed 266–7
 - PARAFAC model for 260–1, 266
 - preparing data for analysis 261–2
 - preprocessing of data 262
 - selecting number of components 264–5
 - solvent-related Raman scatter 261–2
 - using model on new data 270–3

- fluorescence spectroscopy (*Continued*)
 using test sample scores in calibration model
 272–3
 verifying number of components 267–8
 excitation-emission matrix (EEM) 132, 261, 263
 handling of missing data 132–3
 line plots 183
 PARAFAC model 261–2
 rank annihilation factor analysis 138
 Rayleigh scatter in 132, 261, 263
 residuals used for assessing number of components
 162–3
 rotation of Tucker3 core-array 101–2
 second-order calibration in 285
 split-half analysis 165–6, 267–8
 uses 259
- food processing, predicting product quality 78–9
- four-way array(s), as fourth-order tensor(s) 32
- four-way data 11
- Fourier transforms 247
- Frobenius norm 111
- fudge (zoom) factor, in biplots 207, 211, 212
- GEMANOVA *see* generalized multiplicative analysis
 of variance
- generalized eigenvalue problem 139
 formulated in general terms 141
- generalized multiplicative analysis of variance
 (GEMANOVA) model 340–1
 application 344–6
- generalized rank annihilation method (GRAM) 136,
 138–44
 algorithm 142
 basic theory 139
 compared with least squares PARAFAC algorithm
 142
 second-order calibration problem solved using 279,
 280, 282, 346–7
 compared with other methods 282, 285
 in example 280, 282, 283
- graphing techniques
 not used 175
see also plotting techniques
- greyvalues, in visualization 176, 201, 202, 204, 205
- Hadamard product 17
 example 17
 notation 17, 134
 properties 17–18
- Harshman three-way model 58–9, 60
- heavy-metal analysis, in environmental studies 280–3
- heterogeneous variables 5
- heteroscedasticity
 in regression modelling 195
 scaling to accommodate 234
- hierarchy, meaning of term 90
- hierarchy of models 93–5, 154–5
- higher-way data 11
- histograms 177, 189
 three-dimensional 201
see also bar charts/plots
- history
 multi-way analysis 57–9
 plotting in chemical three-way analysis 180
 plotting in three-way analysis 179–80
- homogeneous variables 4–5
- Hotelling's T^2 statistic 172, 289
- hyphenated data
 examples 257
 PARAFAC models for 62, 63, 64–5
- ICP-AES analysis, in environmental study 70,
 215
- image analysis 201–2
 five-way array 11
 three-way arrays in 3, 11, 73
 Tucker1 model used 73–4
- image(s) 73, 176, 201
- influence analysis 171–3
 in example(s) 216, 219, 266–7
- influence measures 171, 216
- inner product(s), of vectors 13
- intercept(s) 222
see also offset(s)
- intercorrelations 5
- interferent
 example 280, 281
 meaning of term 278
- inverse problem 8
- iterative preprocessing 237
- joint plots 176, 179, 213–15
 in example 215, 321
see also biplots
- Khatri-Rao product 18, 115
 example 18
- Khatri-Rao product notation
 PARAFAC model in 85–6
 PCA model in 43, 44
- Kronecker product 16, 85
 example 16
 properties 16–17
 shorthand notation for 125
- Kronecker product notation
 PARAFAC model in 84
 PCA model in 43, 44
 Tucker1 model in 87
 Tucker3 model in 68
- Kruskal conditions 103, 279, 348, 349
- Lambert-Beer law 280
 deviations from 282
- large data sets, visualization of 216, 218

- latent variables
 - as combinations of manifest/measured variables 35–6, 69
 - estimation of 7–8, 53
 - and Tucker models 68–9
- least squares algorithms *see* alternating least squares algorithms
- least squares approach to missing data 133
- least squares minimization 48
- leverage 171–3
 - example using 172–3
 - visualization of 171, 172, 216, 219, 267
- Levin, Joseph 58
- line plots 176, 184–6
 - examples 183–5, 189, 218, 315, 316
- linear projection methods 7
- linearity 21
- liquid chromatography
 - exploratory analysis in 302–11
 - second-order calibration in 285, 302
 - see also* normal-phase liquid chromatography; reversed-phase liquid chromatography
- loading plots 190
 - in examples
 - batch-synthesis examples 189, 326
 - liquid chromatography example 307, 308, 309, 310
 - PEAT example 190, 191, 192, 193
 - SEEDS example 338, 339
 - VENICE data model 316, 317
 - wine example 209, 210
- loadings (vectors) 11, 12
- MacGregor's Tucker1 model 301
- Mahalanobis distance(s) 171, 207
 - in example 210
- MALT example 324, 334
 - model building and interpretation 337
- MANOVA, in examples 337–8
- many-fit diagnostics 146
- mass spectrometry
 - curve resolution in 274
 - MS/MS spectra, second-order calibration used in 284–5
- matrices
 - pictorial representation 13, 14
 - as second-order tensor(s) 32
- matricizing 19–20
 - of three-way arrays 30
- matrix product(s) 13, 16–18
 - pictorial representation 14
 - see also* Hadamard product; Khatri-Rao product; Kronecker product
- maximal variance structure 130
- mean-centering 221
 - see also* centering
- meat shelf life *see* modified-atmosphere packaging
- missing data
 - and centering 232, 251
 - handling 131–4
 - expectation maximization approach 151, 312
 - iterative data imputation approach 134
 - NIPALS approach compared with least squares method 132–3
 - weighted regression approach 134
- model hierarchy 93–5, 154–5
- modified-atmosphere packaging 341
 - example data 341–2
 - ANOVA model 342–3
 - GEMANOVA model 344–6
- Moore-Penrose inverse 17, 48
- movies 176
- multilinear engine, model for environmental data built using 321, 322
- multilinear partial least squares regression 79–83, 124–8
 - basis 124–5
 - predicting responses of new samples 127–8
 - see also* *N*-PLS regression
- multiple linear regression 48–9
 - in example 286
 - problems 47
- multiplicative multilinear models for ANOVA 340–1
 - application(s) 344–6
- multiplicative scatter correction (MSC)
 - transformation of spectra 247
- multivariate calibration 8, 10
- multivariate data structures 5–11
- multivariate image analysis 201, 216
- multivariate process monitoring
 - of batch processes 291
 - of continuous processes 289
- multivariate statistical process control (MSPC) 288
- multi-way arrays
 - addition of 32–3
 - outer products 33–4
 - subtraction of 32–3
- multi-way component models, cross-validation of 149–53
- multi-way covariates regression models 77, 128–30
 - alternating least squares algorithm 129
 - comparison with other regression models 83
 - example of use 78–9
 - extension to multi-way arrays 79
 - predicting responses of new samples 130
- multi-way data, analysis of 1
- multi-way regression application 285–8
 - background to theory 285
 - building a regression model 286–8
 - example data sets 286
- multi-way regression models, cross-validation of 153

- N*-mode principal component analysis *see* Tucker models
- N*-PLS regression 79–83
 comparison with other regression models 83
 example 82–3, 286, 287
 improved method 812
- near-infrared spectra
 in examples 189, 325
 interpretation of 324, 338
see also visible-NIR spectrometry
- nestedness, meaning of term 90
- NIGER data 318, 319
 chemical variables studied 319
 interpretation of 321–2
- NIPALS algorithm 40, 112
 and missing data 132–3
- NMR, second-order calibration in 285
- nonlinear multivariate analysis tools 7
- nonlinear transformations of data 245–6
- non-negative least squares (NNLS) algorithm 136
- non-negativity constraints 135–6
- nonorthogonal-*t* PLS regression 52–3
 algorithm 56
 generalization 79
- normal-phase liquid chromatography 302
 exploratory analysis in 311–12
- object(s), meaning of term 3
- object-space representation 248, 249
- Ockham's razor 154
- offset(s)
 across first mode 222, 224
 across second mode 222, 224, 225
 centering across first mode 225–6
 centering across second mode 226–7
 constant across columns and across rows 224–5
- on-line batch process monitoring 296–9
 'current deviation' approach 297
D test statistic 297
 charts 297, 298, 299, 299, 300
 pictorial representation 297
SPE test statistic 297
 charts 297, 298, 299, 299, 300
 pictorial representation 297
- one-fit diagnostics 146
- optimization techniques 111–13
- orthogonal-*t* PLS regression 51–2
 algorithm 56
- orthogonal transformation 99
 rotation of Tucker3 model by 100, 100–2
- orthogonal Tucker3 model 121
 least squares algorithm for 121–2
- outer product notation of PCA model 42
- outer product(s)
 of multi-way arrays 33–4
 of vectors 13
- outliers 192, 192, 208
 detection of 172, 270–2
- PARAFAC model 59–61
 alternating least squares algorithm 113–19
 block effect in 324, 325, 326
 closed-form solution 136–44
 compared with Tucker models 67–8, 101, 119, 154, 155–6
 cross-validation of, in example(s) 268–70
 degeneracies 89, 107–9, 307
 ways of dealing with 308
 example 62
 in examples
 BATCH data 325–8
 batch polymerization 300
 designed data 336–8
 fluorescence spectroscopy 260–1, 274–5, 286
 liquid chromatography 307–8
 PEAT example 198, 337
 VENICE data 314
 expression in subspaces 140
 in fluorescence spectroscopy 260–1, 274–5, 286
 cross-validation of model 268–70
 multi-way regression 287
 generalized expression 61, 91
 in heavy-metal environmental analysis 280
 intrinsic axes property 63–4
 in mass spectrometry 274
 multi-way regression, in example 286, 287, 288
 notation 60, 84–6
 Khatri-Rao product notation 85–6
 Kronecker product notation 84
 simultaneous components notation 62–3, 63, 64
 tensor products notation 84
 vec-operator notation 84
 number of parameters 97
 partial uniqueness 105
 pictorial representation(s) 61, 62, 63, 64
 preprocessing of data 229, 240
 relationship to Tucker3 model 67–8, 75, 86, 89, 91–3, 101, 154
 scatter plots 197–8
 scree plots 158–60
 sequential compared with simultaneous fitting 64–5
 uniqueness property 102–5
 in UV-vis spectrometry 274
- PARAFAC-ALS algorithm 114–15
 calculating scores of new samples 118–19
 convergence criterion 116
 initializing 116–18
 random start approach 118
 rational start approach 117
 semi-rational start approach 117–18

- more-efficient algorithm 115–16
- second-order calibration problem solved using 282
- parallel factor analysis (PARAFAC) 11–12, 59–66
 - see also* PARAFAC model
- parallel proportional profiles 57, 59
- parsimony principle 57, 154
 - and centering of data 229
- partial least squares (PLS) regression 50–3
 - algorithms 55–6, 125–7
 - basis 50–1
 - in example 286, 287
 - general scheme 53
 - multilinear 79–83, 124–8
 - nonorthogonal **t** vectors version 52–3, 56
 - orthogonal **t** vectors version 51–2, 56
 - three-way 79, 125–7
 - see also* *N*-PLS regression; tri-PLS algorithm
- partial uniqueness, of three-way component models 105
- PCA *see* principal component analysis
- PCA models
 - in examples
 - in batch polymerization example 294
 - liquid chromatography example 305–7
 - notation 40–5
 - Khatri-Rao product notation 43, 44
 - Kronecker product notation 43, 44
 - with matrices and vectors 41–2
 - outer product notation 42
 - with typical elements and summation 40–1
 - vec-operator notation 43, 44
 - pictorial representations 44, 45
- PEAT example 324, 334–5
 - bar plots 161, 190, 219
 - dendrogram 204
 - images of matricized arrays 203
 - images of Tucker core array 205
 - influence plots 216, 219
 - line plots 198
 - loading plots 191, 193, 198, 199
 - model building and interpretation 337
 - particle size distribution 187–9, 190
 - residual plots 203, 216, 217
 - scatter plots 191, 193, 198, 199, 219, 337
 - schematic view of data array 188
 - scree plots 159–60
 - squared Mahalanobis distance plot 219
 - three-dimensional loading plots 194
- Penrose problem 129
- permutational nonuniqueness 98
- pictorial representation, rules for 13, 14
- plotting techniques 175–9
 - biplots 176, 179, 206–13
 - dendrograms 177–8, 202–4
 - history 179–80
 - joint plots 176, 179, 213–15
 - leverage plots 216
 - line plots 176, 184–6
 - listed 176, 177
 - psychology of proportions 199, 200
 - residual plots 216
 - scatter plots 176, 178, 190–2
 - problems 192–201
 - scree plot(s) 157–8, 180–1, 184
- polyaromatic compounds
 - contour plots 182
 - line plots 183
- polymerization, batch process monitoring of 291–3, 301–2
- prediction error sum of squares (PRESS) 27, 148–9, 152
- preprocessing 221–55
 - factors affecting 2
 - iterative 237
 - other types 245–7
 - three-way 239–44
 - two-way 228–38
 - see also* centering; scaling
- principal component analysis (PCA) 7, 11, 37–45
 - formal treatment 38–40
 - N*-mode *see* Tucker models
 - outer product notation 42
 - Pearson's ('variance explained by principal components') approach 38, 39–40, 54–5
 - preprocessing of data 253, 254
 - scatter plots in 192, 195–7
 - as singular value decomposition 40, 206–7
 - 'variance of principal components' approach 38, 39, 54
 - see also* PCA models
- principal component regression (PCR) 49–50
 - in example 286, 287
- problem definition 2
- process chemometrics 257, 288–302
 - typical three-way array for 258
 - see also* batch synthesis example
- process control, meaning of term 288
- process monitoring
 - batch processes 290–1
 - example 291–6
 - continuous processes 289
 - meaning of term 288
 - on-line 296–9
 - operational/test phase 289
 - purpose 289
 - training phase 289
 - type of variation 289
- process variable measurements, to control product quality 290
- product quality measurements 290
- projection methods 7
- proper centering 244, 245
- proper scaling 244, 245

- pseudo-rank
 of three-way arrays 30–1
 analysis in example 293, 294
 of two-way arrays 24, 25
 determination of 27
- quantile plots 177
- quantitative structure-activity relationships (QSARs) 8
 three-dimensional (3D-QSAR) 11
- rainwater example, scaling of data 233–4
- rank
 mathematical concept not convenient in chemical modelling 23–4
 of three-way arrays 28–32
 of two-way arrays 22–8
- k*-rank 28
- rank annihilation 59
- rank annihilation factor analysis (RAFA) 136–8
 automated version 138–42
 line plots after 180, 184
see also generalized rank annihilation method
- rank deficiency
 in three-way arrays 31–2
 in two-way arrays 24–5
- rank reduction 25–6, 137
 automation of 138
 and centering of data 229, 250–1
- rank-one second-order instruments 276
- receptor modeling 320
 examples of applications 318, 320–1
- regression methods
 optimization problems 9
 prediction problems 8
- regression models 46–8, 53
 cross-validation of 153
 nomenclature 47
 scatter plots for 192, 195
 selecting number of components 166
 three-way two-block data analysis using 76–83
 two-way two-block data analysis using 46–53
- residual analysis 166–70
- residual plots 216
 in examples
 liquid chromatography example 307
 PEAT example 203, 217
- residuals 167
 number of components determined using 161–3
 techniques for studying 169–70
 visualization of 163, 169–70
- restricted Tucker3 model 122
- reversed-phase liquid chromatography
 exploratory analysis in 302–11
 second-order calibration in 285
- reversed-phase liquid chromatography example 302–4
 building and using models 304–11
 initial calculations 304–7
- PARAFAC analysis 307–8
 principal component analysis 305–7
 Tucker3 analysis 308–11
- mobile phase in 303
 retention factors calculated and transformed 304–5
 solutes in 303, 304
 stationary phases in 303
 source of variation 304
- rotational freedom
 in PCA models 40
 in Tucker3 models 99
- row-centering 226–7
- row-rank
 of three-way arrays 30
 of two-way arrays 23
- rows
 in three-way arrays 14, 15, 15
 in two-way arrays 2, 3
- Savitzky-Golay smoothing derivative 186, 189, 247, 327
 vis-NIR spectra affected by 189, 328
- scalar(s)
 pictorial representation 13, 14
 as zero-order tensor(s) 32
- scaling of data 5, 178, 232–8
 alternatives 237–8
 compared with centering 221, 244–5
 definitions 222–7
 examples 233–4, 235, 236
 incorrect 243
 interdependence with centering
 multi-way arrays 243–4
 two-way arrays 238–9
 method of operation 235–7
 proper 244, 245
 purpose 221
 reason for wording ‘within’ 227
 reasons for use 234–5
 rules for 228
 situations when scaling does not work 237
 three-way arrays 242–3
 incorrect use of scaling 243
 two-way arrays 232–8
 within first mode
 three-way data 242
 two-way data 227, 235
 within second mode, two-way data 227, 235
- scaling nonuniqueness 98
- scatter plots 176, 178, 190–2
 examples 195–7, 333, 336
 of images 201
 problems 192–201
 including the origin 200–1
 (non)orthonormality of base 192, 195–8
 scale problems 199
 sign indeterminacy 200

- for regression plotting 192, 195
- three-dimensional 176, 192, 194
- see also* loading plots; score plots
- SCENES data 318
 - chemical variables studied 319
 - interpretation of 322
- scientific visualization 218
- score plots 190
 - examples 197, 208–10, 295–6, 307, 330, 333
 - batch polymerization example 295–6
 - batch synthesis example 330, 331
 - liquid chromatography example 307
 - wine example 208–10
- scores (vectors) 11–12
 - predicting for new data 127, 270
- scree plot(s) 157–8, 180–1, 184
 - examples 159, 186
 - hypothetical example 158
 - for PARAFAC model 158–60
 - for Tucker3 model 161, 162
- second-order advantage 106, 278
- second-order calibration 10, 31, 106, 141, 257
 - applications
 - in chromatography 302
 - flow injection analysis 284
 - heavy-metal environmental analysis 280–3, 284
 - background of theory 276–80
 - disadvantages 284
 - environmental example 280–5
 - building and using models 280, 282
 - principles 280, 281
 - typical three-way array for 258
- second-order calibration problems
 - case (ii): mixtures with more than one interferent 279, 347–8
 - case (iv): multiple mixtures 279
 - case (iii): multiple standards 279, 348–9
 - case (i): standard containing several analytes 279
 - generalized rank annihilation solution 279, 280, 282, 346–7
 - compared with other methods 282, 285
 - in example 280, 282, 283
- second-order instruments 276
 - effect of interferent on calibration 278
- SEEDS example 324, 335
 - model building and interpretation 337–8, 338, 339
- segmented cross-validation 149
- selection of model 154–6
- selection of number of components 156–66
 - in example(s) 165–6, 264–5, 294
- sensory analysis
 - comparing configurations 9
 - prediction methods 82–3
 - selection of model 155–6
- sign reversal problems 200, 331, 333
- simultaneous components notation
 - Tucker1 model in 87
 - Tucker2 model in 86
 - Tucker3 model in 86
- simultaneous two-way centering and scaling 238–9
- single-centering
 - of three-way data 241
 - of two-way data 227, 230
- singular value decomposition (SVD) 37
 - in examples
 - in batch synthesis example 328–9
 - wine example 208
 - principal component analysis as 40, 206–7
 - relationship with eigendecomposition 38
 - weights in tri-PLS1 obtained from 144
- slab-centering 227
- spectral analysis plots 177
- split-half analysis 164–6
 - examples 165–6, 267–8
- squared Mahalanobis distance(s)
 - calculation of 172, 271
 - leverage viewed as 171, 172, 216, 218
 - in outlier detection 172, 271–2
- Standard Normal Variate (SNV) transformation of spectra 247
- statistical reliability 146
- sub-arrays 20, 167
- subspaces
 - finding suitable subspaces from real data 140–1
 - PARAFAC model expressed in 140
- sufficient variation 105
- sugar quality example 286
 - multi-way regression 286, 287
- supervised classification 7
- 'swamp' (slow convergence) 117, 229
 - avoidance of 117, 229
- SYNT example 324, 336
 - model building and interpretation 338–9
 - PARAFAC analysis 186–7, 189, 338–9
- tensor analysis 32
- tensor product 18–19
- tensor product notation
 - in PARAFAC model 84
 - in Tucker2 model 87
 - in Tucker3 model 86
- test-set validation 147
 - in example(s) 271, 272
- theoretical appropriateness 145
- thin-layer chromatography, second-order calibration in 285
- third-order instruments 278
- three-dimensional histograms 201
- three-dimensional scatter plots 176, 192
 - examples 194
- three-dimensional visualization 177, 216

- three-way array(s)
 - columns 15, 15
 - decomposition of 11–12
 - degrees of freedom 97
 - dimensionality vector 30
 - frontal slices 3, 4, 14, 14
 - horizontal slices 3, 4, 14, 14
 - pictorial representation 13, 14
 - preprocessing of 239–44
 - rank of 28–32
 - border rank 29
 - definitions 28–9
 - determination of 32
 - individual rank 29
 - maximum rank 29
 - properties 29
 - typical rank 29
 - rows 15, 15
 - as third-order tensor(s) 32
 - tubes 15, 15
 - vertical slices 3, 4, 14, 14
- three-way component models 59–76
 - complexity 97
 - degrees of freedom 97
 - extensions for multi-way arrays 76
 - fitting errors 96
 - general framework for 75–6
 - hierarchy 90
 - nestedness 90–1
 - number of parameters in 96–7
 - PARAFAC models 59–66
 - properties 89–109
 - relationships between 89–97
 - Tucker models 66–75
 - uniqueness in 98, 102–6
 - see also* PARAFAC models; Tucker models
- three-way data, representation 2–3, 3
- three-way multiblock data 10
- three-way one-block data 9–10
 - calibration and resolution 10
 - exploration of 9–10
 - models for 59–76
- three-way partial least squares regression 79
 - algorithm(s) 125–7
 - see also* Tucker1-PLS model
- three-way regression models 76–83
 - sequential methods 76, 79–83
 - simultaneous methods 76, 77–9
- three-way two-block data 10
 - models for 76–83
- Thurstone's principle of parsimony 57
- tri-PLS1 algorithm 125–6
 - weights obtained from singular value decomposition 144
- tri-PLS2 algorithm 126–7
- triad(s) 18
 - visualization of 18
- trilinear decomposition *see* PARAFAC model
- trilinear decomposition (TLD) algorithm 143
- trilinearity 21, 277–8
- tube-rank, of three-way arrays 30
- tubes, in three-way arrays 14, 15, 15
- TUCKALS3 algorithm 121–2
 - initializing 122–3
- Tucker, Ledyard 58, 66, 179
- Tucker congruence coefficient 107, 229
 - calculation of 107
 - in examples 268
- Tucker models 66–75
 - algorithms for fitting 121–2, 123–4
 - hierarchy 93–5
 - interpretation in terms of new basis 71–2
 - pictorial representations 67, 73, 75
 - uniqueness in 98
- Tucker1 model 73–5
 - algorithm for fitting 124
 - alternating least squares algorithm for 124
 - derivation from Tucker2 model 94
 - differences between MacGregor and Wold approaches 301
 - in examples
 - batch polymerization process 294, 295, 296, 298, 299, 301
 - image analysis 73–4
 - in Kronecker product notation 87
 - number of parameters 97
 - pictorial representation 75
 - in simultaneous components notation 87
 - in summation notation 74
- Tucker1-PLS model 79
 - comparison with other regression models 83
- Tucker2 model 72–3
 - algorithm for fitting 123–4
 - alternating least squares algorithm for 123–4
 - derivation from Tucker3 model 94
 - number of parameters 97
 - pictorial representation 73
 - in simultaneous components notation 86
 - in tensor product notation 87
- Tucker3 model 66–72
 - algorithm for fitting 121–2
 - alternating least squares algorithm for 121–2
 - convergence criterion 122
 - initializing 122–3
 - calculating scores of new samples 123
 - closed-form solution 120–1
 - compared with PARAFAC model 67–8, 101, 119
 - constrained 95
 - in FIA example 95–6
 - number of parameters 97
 - uniqueness 105–6
 - core array in 66, 119
 - analysis 156
 - calculating 119–20
 - rank 107
 - rotating 101–2, 130–1, 156

- simplifying 99–101
 - size 30, 106
 - visualizing 204–5, 321
- decomposition of 66–7, 213–14
- deriving from latent structure 68–9
- in examples
 - batch polymerization process 294–5, 296, 296, 298, 300, 301
 - environmental studies 70, 321, 322
 - liquid chromatography example 308–11
- generalization for three-way arrays 67, 93
- Kronecker multiplication for 68
- in Kronecker product notation 68
- least squares algorithm 121–2
- number of parameters 97
- pictorial representation 67, 213
- properties 66, 70–1, 94, 106–7
- relationship to PARAFAC model 67–8, 75, 86, 89, 91–3, 95, 101, 154
- rotational freedom in 99
- scree plot for 161, 162
- in simultaneous components notation 86
- squared Mahalanobis distance 173
- in tensor product notation 86
- Tucker3 regression, in example 286, 287
- two-factor degeneracy 107, 116, 117
- two-mode component analysis 45–6
- two-way array(s)
 - decomposition of 11, 35, 36
 - preprocessing of 228–38
 - rank of 22–8
 - definitions 22–3
 - determination of 27
 - properties 23
 - rows and columns 2, 3
- two-way centering 228–32
- two-way data
 - centering of 228–32
 - representation 2, 3
 - scaling of 232–8
- two-way multiblock data 9
- two-way one-block data 6–8
 - analysis using component models 35–46
 - classification methods 7
 - estimation of underlying phenomena 7–8
 - exploration of 6–7
- two-way scaling 232–8
- two-way two-block data 8–9
 - analysis using regression models 46–53
 - exploring and optimization problems 8–9
 - prediction problems 8
- unfolding, meaning of term 19
- uniqueness property
 - component models 98, 102–6
 - meaning of term 98, 103
- constrained Tucker3 models 105–6
- PARAFAC models 102–5
- principal components 40
- Tucker models 98
- unsupervised classification 7
- UV-vis spectra
 - analysis of 274
 - examples of offsets 223–6
- UV-vis spectrometry
 - curve resolution in 274
 - second-order calibration in 285
- validation
 - meaning of term 145–7
 - see also* cross-validation
- variable(s)
 - latent 7–8
 - meaning of term 3
 - types 4–5
- variable-space representation 248–9
- variation, chemical sources 23–4
- vec-operator 15–16
 - notation in PARAFAC model 84
 - notation in PCA model 44
- vector(s)
 - as first-order tensor(s) 32
 - multiplication of 13, 14
 - pictorial representation 13, 14
- vector product 13, 14
- VENICE data 313–14, 318, 319–20
 - chemical variables studied 319
 - PARAFAC model 314, 315
 - interpretation of model 314, 316, 317, 321
- visible-NIR spectrometry, batch synthesis monitored using 186, 323, 336
- visualization 175–220
 - of large data sets 216, 218
 - of residuals 163, 169–70
 - see also* plotting techniques
- water quality, in environmental studies 215, 319–20
- wavelet transforms 247
- weather systems data, visualization of 216, 218
- weighted least squares fitting, and scaling 236–7
- wine example 208–12
- Wold's Tucker1 model 301
- X-ray photoelectron spectroscopy 334
 - see also* XPS example
- XPS example 324, 334
 - model building and interpretation 336–7
- zeroth-order instruments 276
 - effect of interferent on calibration 278