

## Subject Index

### a

acoustic branch 152  
 – longitudinal 131  
 adiabatic approximation 113 ff., 151  
 Ag 248, 329  
 AgBr 154  
 AlSb 36  
 Al 151, 243, 318, 325, 329  
 amorphous materials 105, 350  
 anharmonic constants 186  
 anharmonic effect 150, 182, 245, 321  
 Ar 139  
 argon 92  
 asymmetry parameter  $\eta$  40  
 a-Tb<sub>33</sub>Fe<sub>67</sub> 364  
 Au 329, 334, 342  
<sup>197</sup>Au 83, 181, 327, 333

### b

BaSnO<sub>3</sub> 205 ff.  
 bcc *see* cubic  
 Bessel function 11, 13, 269, 274  
 – modified 375  
 bis(3-methylpentadienyl)iron 356  
 black absorber 12  
 bond-charge model 139  
 Born–Mayer potential 139  
 Born–Oppenheimer theorem 113  
 Born–von Karman theory 113, 146, 165, 355  
 Bravais crystal 181, 184  
 Bravais lattice 149, 305, 310, 321  
 Breit–Wigner formula 9  
 Brillouin zone 126, 131, 135, 143, 155, 171

### c

C (diamond) 151  
 C<sub>6</sub>H<sub>4</sub>(OH)<sub>2</sub> 188  
 C<sub>8</sub>Cs 192

center shift 346  
 central force approximation 316, 384  
 Clebsch–Gordan (C-G) coefficients 52, 60, 193  
 (CN<sub>3</sub>H<sub>6</sub>)<sub>2</sub>[Fe(CN)<sub>5</sub>NO] 281, 299  
 coherent elastic scattering 229 ff.  
 coherent state 21  
 – harmonic oscillator 17  
 – pseudo-classical 18  
 constant-Q method 162  
 copper 164  
 correlation function, time-dependent 287  
 cosine effect 94  
 Cr 64, 319, 329  
<sup>133</sup>Cs 192, 195  
 Cu 149, 151, 246, 319, 325, 328, 329  
 Cu<sub>5</sub>FeS<sub>4</sub> 88  
 cubic  
 – body-centered (bcc) 120, 125, 128, 313, 345, 380, 393  
 – face-centered (fcc) 119, 125, 128, 150, 183, 201, 313, 380, 393  
 – simple 393  
 CuInSe<sub>2</sub> 172  
 CuRh<sub>1.95</sub>Sn<sub>0.05</sub>Se<sub>4</sub> 189  
 Cu–Zn alloy 347

### d

damped harmonic oscillator function 363  
 DB *see* dynamical beats  
 Debye  
 – model 24, 143, 145, 149, 190, 198, 202, 312, 345, 346, 348  
 – temperature 24, 26, 145, 149, 177, 190, 236, 249, 281, 309, 335, 342, 345, 352  
 Debye–Waller factor  $f_D$  151, 155, 160, 229 ff., 241, 245 ff.

- density functional theory (DFT) 165, 168, 355  
density of states (DOS) 24, 142 ff., 165  
– host 333  
– impurity 333  
– partial 172, 286, 292, 299, 359  
– partial vibrational 354, 366  
– phonon 190, 253, 259, 285, 291, 293, 295, 298, 311, 366  
– vibrational 317, 352, 358, 359, 362  
diamond 26, 150, 335 f.  
dispersion relation 124, 126, 138, 141  
Doppler  
– broadening 4  
– effect 2, 14, 79  
– energy shift 2  
– second-order effect 73  
– transverse Doppler effect 196 ff.  
– velocity 6, 64, 373  
Doppler shift, second-order 148, 177, 318, 322, 333, 345, 348  
DOS *see* density of states  
<sup>161</sup>Dy 56, 83, 258  
Dy<sub>3</sub>Fe<sub>5</sub>O<sub>12</sub> 358  
dynamic response function 341  
dynamical beats (DB) 265, 274, 281, 284  
dynamical matrix 123 ff., 165  
Dyson equation 308, 317
- e**  
E2-type radiation 195  
effective magnetic field 53 ff.  
effective thickness  $t_a$  10, 72, 85, 269, 274, 278  
effective vibrating mass 177, 198, 355  
Einstein model 143, 323, 326  
Einstein temperature 144, 177  
Einstein–Debye formula 346  
Einstein–Debye model 323  
electric field gradient (EFG) 31, 43 ff.  
electric quadrupole interaction 29, 38 ff.  
electronegativity 37  
electron–electron interaction 168  
enhancement of coherent channel 266 f.  
<sup>166</sup>Er 56  
<sup>151</sup>Eu 56, 83, 258, 294  
Eu<sub>2</sub>Ti<sub>2</sub>O<sub>7</sub> 193  
EuBa<sub>2</sub>Cu<sub>3</sub>O<sub>7</sub> 205, 206  
EuBa<sub>2</sub>Cu<sub>3</sub>O<sub>7- $\delta$</sub>  189  
exchange-correlation energy 169  
extrapolation method 165
- f**  
face-centered cubic (fcc) 119, 125, 128, 150, 183, 201, 313, 380, 393  
– crystal 134  
Fe 149, 294  
<sup>57</sup>Fe 8, 33, 39, 43, 51, 82, 180, 200, 283, 327, 334, 356  
 $\alpha$ -Fe 34, 36, 56, 57, 66, 100, 149, 191, 206, 265, 283, 294, 299, 342, 356  
Fe nucleus (<sup>57</sup>Fe) 3  
Fe(C<sub>5</sub>H<sub>5</sub>)<sub>2</sub> 88, 206, 356  
[Fe(C<sub>5</sub>H<sub>7</sub>O<sub>3</sub>)<sub>3</sub>]<sup>3+</sup> 355  
[Fe(CN)<sub>6</sub>]<sup>4-</sup> 355  
[Fe(H<sub>2</sub>O)<sub>6</sub>]<sup>3+</sup> 355  
Fe[Co(CN)<sub>6</sub>] 151  
Fe[Ir(CN)<sub>6</sub>] 151  
Fe[Rh(CN)<sub>6</sub>] 151  
Fe<sup>2+</sup> 45, 54, 189  
Fe<sub>2</sub>O<sub>3</sub> 35, 299  
 $\alpha$ -Fe<sub>2</sub>O<sub>3</sub> 88, 206, 216, 223  
 $\alpha$ -<sup>57</sup>Fe<sub>2</sub>O<sub>3</sub> 261  
Fe<sub>2</sub>Tb 294  
Fe<sup>3+</sup> 45, 54  
Fe<sub>3</sub>Al 294  
Fe<sub>3</sub>BO<sub>6</sub> 227  
Fe<sub>40</sub>Ni<sub>40</sub>P<sub>14</sub>B<sub>6</sub> 105, 351, 353  
Fe<sub>67</sub>Tb<sub>33</sub> 294  
Fe<sub>74</sub>Si<sub>10</sub>B<sub>16</sub> 351  
Fe<sub>75</sub>P<sub>15</sub>C<sub>10</sub> 351  
Fe<sub>80</sub>B<sub>20</sub> 353  
FeBO<sub>3</sub> 294, 364  
<sup>57</sup>FeBO<sub>3</sub> 261  
FeC<sub>2</sub>Ni 364  
FeCl<sub>2</sub> 189  
FeCO<sub>3</sub> 50  
FeF<sub>2</sub> 189  
FeF<sub>3</sub> 51  
FeGe powder 60  
Fe–Rh–Ni polarizer 64, 70  
Fermi contact field 54  
Fermi pseudopotential 157  
ferrocene 88, 356  
ferrous halides 37  
FeS<sub>2</sub> 206  
Fe–Si 64  
FeSO<sub>4</sub>·7H<sub>2</sub>O 88  
first-principles methods 165 ff., 299  
force constant 117, 138, 150, 152, 291  
– central forces 383 ff.  
– host–host 305, 310  
– impurity–host 305, 310, 322  
– matrix 166, 380  
– ratio 326, 333, 346  
– tensor 307  
Fourier transform 108, 184, 296, 306  
Fourier-logarithm decomposition 296  
fractional absorption intensity 12

**g**

GaP 152  
 GaSb 36  
 Gauss–Newton method 105, 108  
<sup>155</sup>Gd 56, 83, 195  
 Gd<sub>2</sub>Ti<sub>2</sub>O<sub>7</sub> 195  
 Ge 325, 330  
 Ge(Li) 92  
 g-factor 52  
 G–K effect *see* Goldanskii–Karyagin effect  
 Glauber’s formula 23  
 Goldanskii–Karyagin effect 192 ff., 281, 343  
 Green’s function 152, 184, 306, 308, 314, 385  
   – matrices 390  
 Grüneisen constant 185, 190  
 guanidinium nitroprusside 281

**h**

harmonic approximation 115 ff., 140, 166, 180, 234, 341  
 harmonic lattice 120  
 harmonic oscillator 16  
 Hartree functional 169  
 hcp-Fe 191  
 heat capacity 141  
 helicity 62, 66  
 Hellmann–Feynman theorem 166  
 Hesse method 105  
 Hessian 166  
 hexacyanoferrate(II) 359  
<sup>201</sup>Hg 258  
<sup>165</sup>Ho 56  
 HoFe<sub>2</sub> 66  
 hydroquinone 188  
 hyperfine interactions 72

**i**

<sup>129</sup>I 83  
 impurity dynamic response function 309  
 impurity site moments 320 ff.  
 impurity space 307  
 impurity, interstitial 333  
 impurity, substitutional 305, 382  
 indium 146  
 InSb 36  
 internal conversion coefficient 83  
   – partial 286  
 Ir 206, 329  
<sup>191</sup>Ir 5, 8, 56  
 irreducible representation 313, 394  
 isomer shift 29, 31 ff., 72, 105, 348  
   – calibration of 34 f.  
 isometric state 9

**k**

K 149  
<sup>40</sup>K 82  
 K<sub>3</sub>Fe(CN)<sub>6</sub> 88  
 K<sub>4</sub>Fe(CN)<sub>6</sub>·3H<sub>2</sub>O 88, 205, 206, 225  
 KCl 243, 246  
 Keating model 139  
 K-fluorescence photons 214, 286, 289  
 Kohn–Sham equations 170  
 Kr 139  
<sup>83</sup>Kr 258  
 Kramers–Kronig relations 389  
 krypton 92

**l**

Lamb–Mössbauer factor  $f_{LM}$  229 ff., 281, 291, 295, 299  
 Laplace equation 40  
 lattice vibrations 123 ff., 139, 311  
 lattice wave 115, 122  
<sup>6</sup>Li 154  
<sup>7</sup>Li 154  
 linear response 166 ff.  
 Lipkin’s sum rule 7, 25, 291  
 local density approximation 172  
 localized vibrations 152 ff.  
 Lorentzian distribution 9, 108  
 low-temperature anharmonicity 188

**m**

M1-type radiation 52  
 M1-type transitions 221  
 magnetic dipole interaction 29, 51 ff.  
 magnetic hyperfine field 55, 63, 73, 86, 105  
 magnetic hyperfine splitting 55  
 Mannheim model 313, 322, 324 ff., 333  
 Maradudin–Flinn model 323  
 mass absorption coefficients 10, 85, 396 ff.  
 mass defect approximation 317  
 McMillan ratio 200, 327  
 mean-square displacement 7, 151, 155, 177 ff., 189, 245, 309, 335, 343, 345, 358, 391 f.  
 mean-square velocity 155, 177 ff., 197, 310  
 Mo 329  
 mode  
   – acoustic 358  
   – composition factor 358  
   – gap 154  
   – intermolecular 355  
   – intramolecular 355  
   – localized 311, 317

- longitudinal 138
  - normal 122, 129, 131, 140, 318
  - resonance modes 311, 317
  - resonant 154
  - stretching 361
  - transverse 138
- moments of frequency distribution 146 ff.
- Morse pair potentials 139
- Mössbauer diffraction 216, 223 ff.
- Mössbauer effect 5, 6, 73, 151
- phonon-assisted 297, 361
- Mössbauer lattice temperature 356
- Mössbauer spectroscopy 6, 72, 253 ff., 295
- conventional 298
  - energy domain 253, 280
  - spectrometer 79 ff.
  - synchrotron 202, 253 ff., 298
  - time domain 253, 265, 280

**n**

- Na 149, 151
- $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}] \cdot 2\text{H}_2\text{O}$  88, 206
- $\text{Na}_4\text{Fe}(\text{CN})_6 \cdot 10\text{H}_2\text{O}$  336
- NaCl 246
- NaF 143
- Nb 329
- $\text{Nb}_3\text{Sn}$  189, 201
- $\text{Nd}_2\text{Fe}_{14}\text{B}$  56, 70
- Ne 139, 150
- neon 92
- neutron scattering 156 ff.
- neutron scattering, inelastic 366
- $(\text{NH}_4)_2\text{Mg}^{57}\text{Fe}(\text{CN})_6$  274
- $(\text{NH}_4)_2\text{MgFe}(\text{CN})_6$  359
- Ni 151, 243, 329, 342
- $^{61}\text{Ni}$  56
- $[\text{Ni}(\text{CN})_4]^{2-}$  355
- normal coordinates 22, 120
- normal mode 122
- $^{237}\text{Np}$  33, 56, 83
- nuclear Bragg scattering 220, 225, 229 ff., 259, 265, 278
- nuclear exciton 265 f., 271
- nuclear forward scattering 214, 220, 229 ff., 259, 271, 281, 295, 298
- nuclear resonant scattering 155, 214
- inelastic 155, 295, 298, 318, 362
  - inelastic incoherent 289

**o**

- operator
- annihilation 16, 20
  - creation 16, 20
  - density 18

- displacement 17, 20
  - Hermitian 18
- optical branch 152
- longitudinal 131
- optimal sample thickness 84
- $^{191}\text{Os}$  5

**p**

- Pb 151, 218, 248
- Pd 84, 328, 329
- $\text{Pd}_3\text{Sn}$  201
- $\text{Pd}_{80}\text{Si}_{20}$  353
- phonon annihilation probability 291
- phonon creation probability 291
- phonon–phonon interaction 182
- plane-wave basis 171
- polarization 62 ff.
- $\pi$ -polarization 64, 69, 262
  - $\sigma$ -polarization 64, 69, 262
  - vector 63, 122, 164
- principal axis system 31, 40, 48, 56
- pseudo-classical quantum states 16
- pseudoharmonic approximation 185 ff.
- pseudopotential approximation 171
- Pt 325, 328, 329
- Pt absorbers 6
- $\text{PtB}_2$  188
- $\text{PtCl}_2$  188

**q**

- QB *see* quantum beats
- quadrupole splitting 72
- quantum beats (QB) 265, 274 ff., 284
- quasiharmonic approximation 185

**r**

- Rayleigh scattering 214, 233
- reciprocal lattice 125
- recoil energy 2, 161
- recoilless fraction  $f$  2, 6, 16, 21, 26, 73, 148, 161, 177, 183, 192, 202, 295, 309, 328, 333, 342, 391 f.
- reduced rotation matrix 220
- refractive index 272
- resonance absorption 9
- coefficient 85
- resonance fluorescence 1
- resonance modes 311
- Rh 84, 329
- $^{99}\text{Ru}$  56

**s**

- saturation effect 13, 71 f., 86
- Sb 36

- $^{121}\text{Sb}$  36, 37, 39, 83  
 scattering amplitude 230  
 scattering process  
   – slow 215, 220, 232  
   – fast 215, 232  
 second-order Doppler shift 73, 196, 291  
 self-consistent field 170  
 Si 241, 325  
 Si(Li) 92  
 simple cubic 393  
 $^{149}\text{Sm}$  56  
 $^{119}\text{Sn}$  8, 37, 57, 82, 181, 201, 258, 294, 327, 330, 356  
 $\alpha$ -Sn 294, 325, 330  
 $\beta$ -Sn 294, 342  
 $\text{SnO}_2$  206, 294  
 SNP *see* sodium nitroprusside  
 SnSb 36  
 SnTe 201  
 sodium nitroprusside (SNP) 88  
   –  $\text{Na}_2\text{Fe}(\text{CN})_5\text{NO}\cdot 2\text{H}_2\text{O}$  34  
 specific heat 155  
 speed-up effect 265 ff., 269  
 $\text{SrFe}_{12}\text{O}_{19}$  70  
 $\text{SrFeO}_3$  294  
 stainless steel 294  
 supercell 171  
 symmetry coordinates 314, 393  
 synchrotron radiation 65, 155, 213, 253, 298
- t**  
 Ta 329  
 $^{181}\text{Ta}$  37, 83  
 $^{159}\text{Tb}$  56  
 $\text{Tb}_{33}\text{Fe}_{67}$  364  
 $\text{TbAl}_2$  206  
 $\text{TbO}_7$  206  
 $^{125}\text{Te}$  83, 92  
 thermal diffuse scattering (TDS) 235  
 thin absorber approximation 12, 102  
 thin film 363  
 Thomas–Raleigh formula 234  
 Ti(Fe) alloy ( $\beta$ -) 345  
 time domain Mössbauer spectroscopy 265, 280
- $^{169}\text{Tm}$  56  
 transmission integral 10, 108
- u**  
 unitary transformation 147
- v**  
 V 325, 329  
 van der Waals forces 355  
 VDOS *see* density of states, vibrational  
 velocity calibration 99 ff.  
   – absolute 100 ff.  
 velocity transducer 8, 80, 95  
 Visscher model 323
- w**  
 W 206, 329, 364  
 waveguide 362  
 waves  
   – evanescent 362  
   – longitudinal 133  
   – standing 136, 362  
   – transverse 133, 138  
 Window method 106
- x**  
 Xe 139  
 xenon 92  
 x-ray absorption edge 92  
 x-ray diffraction 151  
 x-ray scattering 155
- y**  
 $^{170}\text{Yb}$  56  
 $\text{YFe}_2$  353
- z**  
 Zeeman effect 51  
 zero phonon process 6, 160, 235  
 zero point energy 140  
 Zn 151, 246, 342  
 $^{67}\text{Zn}$  83, 180, 341, 347  
 $\text{ZnO}$  342  
 $\text{Zr}_{9,5}\text{Fe}_{90,5}$  353

