

INDEX

- A-stability, 375
- Abel
 - Niels Henrik, 470
- absolute error, 4
- acceleration, 133–137
- accuracy, 2, 4, 6, 7, 13, 15, 20, 23–24, 28, 32–33, 43–45, 47, 49–50, 61, 63, 83–84, 88, 90–91, 95, 99, 111, 112, 115, 117, 118, 120, 132, 138, 148–149, 153, 173, 177–178, 185, 188, 191, 196, 197, 210, 231, 250, 253–254, 261–262, 264, 266–267, 272, 276, 279, 286, 288, 292, 297, 302, 304–306, 308, 310, 315, 331, 337, 346, 350, 356, 360–361, 369, 377, 385, 392, 393, 412, 433, 437, 500, 513, 514, 516, 525
- Adams
 - John Crouch, 358
- Adams-Bashforth method, 358, 360, 362, 368–369, 371–372
- Adams-Bashforth-Moulton method, 358, 369, 378
- Adams-Moulton method, 358, 361, 363, 368, 370
- adaptive solvers, 377
- Aitken extrapolation, 133
- Aitken
 - Alexander Craig, 133
- Algol, 29–30
- approximate equality, 16, 32
- approximation error, 16
- associative law of arithmetic, 27
- asymptotic order notation, 16–18, 30, 32
- Atkinson
 - Kendall, 116
- B-splines, 198–200
- backsolve, 76, 409
- backward error analysis, 431
- backward Euler method, 335–336
- backward solution, 76, 409
- banded systems, 443
- Bashforth
 - Francis, 358
- BASIC, 29
- BDF (backward difference formula), 363
- Bernoulli
 - Johann, 2, 24, 294
- Big O notation, 16
- bisection method
 - convergence and error, 88–89
 - derivation, 86–87
 - disadvantages of, 90
 - pseudocode, 89
- bisection, 86–90, 99, 149–151, 155
- bits, 21
- boundary data, 506
- boundary value problem (BVP)
 - definition, 322
 - shooting methods, 395–397
 - simple difference methods, 80–82, 390–393
 - spline solution, 211–214
- boundary value problems, 389
- Boyer

- Carl, 93
- Braun
 - Martin, 122
- Brent's algorithm, 149, 151, 154–155
- Brent-Dekker algorithm, 154–155, 183
- bytes, 21
- C, 29
- C++, 29
- calculus, 2
 - extreme value theorems, 9–10
 - mean value theorems, 9–10
 - Taylor's Theorem, 2–4, 7, 9
- Cauchy form of Taylor remainder, 14
- Cavalieri
 - Bonaventura, 258
- characteristic polynomial, 464–465
- Chebyshev least squares approximation, 230–231
- Chebyshev nodes, 240, 242–246
- Chebyshev polynomials, 227
- Chebyshev
 - Pafnuty I.ovich, 227
- Cholesky factorization, 441, 443, 522–525
- Cholesky Theorem, 441
- Cholesky
 - Andre-Louis, 441
- chopping, 23
- chord method (root finding), 137–138
- cloud effect, 47
- collocation, 214
- complete pivoting, 410
- complete spline, 202
- composite trapezoid rule, 65
- computational mathematics, 2
- computer arithmetic, 21
- computer languages, 29
- condition number, 428
- conditioning, 425
- Conjugate gradient iteration, 526
- convergence
 - acceleration of, 133–137
 - chord method, 138–139
 - fixed point iteration, 123, 126, 128–131
 - of bisection method, 88–89
 - of Newton's method, 95–97, 99–100, 107–109, 112–114
 - secant method, 122, 141–144
- corrected trapezoid rule, 253
- corrector step, 341
- Cotes
 - Roger, 258
- Crank
 - John, 513
- Crank-Nicolson method, 513, 515–518
- cubic splines, 198
- data fitting, 216, 218–220
- Davis
 - Philip J., 227
- de Boor
 - Carl, 198
- defective matrices, 466, 479, 481, 501
- definite integral
 - adaptive quadrature, 304–307, 309–310
 - extrapolation methods, 286–291
 - Gaussian quadrature, 274–283
 - improved trapezoid rule, 253–255
 - midpoint rule, 269, 271
 - Peano estimates, 314
 - review, 250–251
 - Romberg integration, 293–299
 - Simpson's rule, 257–264, 266
 - trapezoid rule, 64–70, 73
- deflation, 488–489, 494
- degree of precision, 261
- Dekker's method, 154
- derivative approximations, 39, 44–47, 183–185
- diagonal dominance, 77–78, 202–203
- difference approximations, 44–45
- DIFSUB, 378
- Discrete Average Value Theorem, 11
- divided difference table, 167–170
- divided differences, 167–170
- double precision, 24
- efficiency, 2
- eigenpairs, 464
- eigenspace, 466
- eigenvalues, 463
 - Gerschgorin's Theorem, 466–467
 - power methods, 476–489, 492
 - QR method, 494–501
 - reduction to Hessenberg form, 470–473, 475–476
 - review, 464–467
- eigenvectors, 464–466, 469, 472–474, 476
- Einstein
 - Albert, 251
- EISPACK, 30
- elementary row operations, 406
- elimination step, 76
- elliptic function theory, 448
- equivalence relation, 16
- error function, 30, 59
- Euler
 - Leonhard, 162, 274
- Euler-Maclaurin formula, 294, 300
- evaluation points, 251
- explicit method, 341, 358, 360
- extended floating point number system, 21
- Fibonacci numbers, 143
- finite element method, 214
- fixed point iteration, 123–124, 126, 128–131
 - existence and convergence theory, 126, 128–131
 - multidimensional, 455–457
- fixed point, 124
- floating-point arithmetic, 24, 26–27
- FORTRAN, 29–30
- function norm, 175
- Galois
 - Evariste, 470
- gamma function, 62, 171–172, 186, 196, 209
- Gauss points, 275
- Gauss
 - Carl Friedrich, 274, 293
- Gauss-Kronrod quadrature, 317

- Gauss-Legendre quadrature, 282
- Gauss-Seidel iteration, 449, 451, 527–528, 530
- Gaussian quadrature, 251, 274–276, 278–287, 299–300, 310, 317
- Gerschgorin's Theorem, 485
- Givens transformations, 502
- global method of root finding, 90, 95
- Goldstine
 - Herman, 93, 326, 354
- Gram-Schmidt process, 224
- Gregory's series, 8
- Gregory
 - James, 8
- Hahn
 - Alexander J., 101
- Halley's method (root-finding), 140–141, 152–153, 183
- Halley
 - Edmund, 93, 140
- Hermite interpolation, 186–188, 190
- Hermite orthogonal polynomials, 227
- Hermite polynomials, 196
- Hermite
 - Charles, 186
- Hessenberg form, 470–474, 476, 480, 487, 489, 494
- homogeneous boundary data, 506
- Horner's rule, 39–42
- Horner
 - William George, 40
- Housholder matrices, 472–473
- Housholder transformations, 472–473
- Housholder
 - Alston, 472
- hybrid algorithms (root finding), 149–151
- IEEE arithmetic, 21
- implicit QR , 499
- IMSL library, 30
- index array, 421, 423
- index map, 524
- initial data (PDE), 506, 518
- initial value problems
 - Euler's method, 52–55
- initial value, 322–323, 365, 376, 388, 395
- inner product, 221–224, 232
- Intermediate Value Theorem, 10, 86, 128
- interpolation error, 175–176, 183–184, 188, 195
- interval of uncertainty, 87
- inverse quadratic interpolation, 180–183
- iteration matrix, 447
- iterative refinement, 437–439
- Jacobi iteration (linear systems), 448–452, 457, 526–527, 530
- Jacobi
 - Karl Gustav Jacob, 448
- Java, 29
- Kepler
 - Johannes, 2
- knots, 192–193, 199–200, 210
- Kronecker delta, 161
- Kronecker
 - Leopold, 161
- Kutta
 - Martin Wilhelm, 352
- L'Hôpital's rule, 24
- L'Hôpital
 - Guillaume Francois Antoine Marquis de, 24
- Lagrange form of Taylor remainder, 14
- Lagrange interpolation, 160, 162, 164
- Lagrange
 - Joseph-Louis, 162
- Laguerre polynomials, 227
- Laguerre
 - Edmond Nicolas, 227
- LAPACK, 30, 435, 460
- legacy code, 30
- Legendre polynomials, 226, 228–230, 232
- Legendre
 - Adrien-Marie, 226
- linear interpolation, 57–62
- linear operator, 252, 312
- LINPACK, 30
- Lipschitz continuity, 324–326
- Lipschitz
 - Rudolf Otto Sigismund, 324
- Lotka-Volterra population model, 370
- Machin
 - John, 13
- machine epsilon, 26–27, 29
- Maclaurin
 - Colin, 294
- Maple, 29–30
- MathCAD, 29–30
- Mathematica, 29–30
- Mathias
 - Roy, 275
- MATLAB, 29–30
- matrix exponential, 463
- matrix norm, 426–427
- mesh points, 250, 260, 304
- method of false position, 92
- method of Heun, 354
- midpoint method, 336
- Milne's method, 370
- modified Euler method, 354
- Moulton
 - Forrest Ray, 358
- Muller's method, 180–183
- multiple roots, 145
- NAG library, 30
- naive Gaussian elimination, 408, 411, 413, 417–418
- natural spline, 202–207
- NETLIB, 30, 246
- Newton interpolation, 165–167, 170
- Newton's method of root finding, 93
 - convergence, 96, 109, 112–114
 - derivation, 93–95
 - error formula, 107–109
 - multiple roots, 145–148
 - stopping, 99–100
 - variants, 137–139
- Newton
 - Isaac, 2, 8, 93

- Newton-Cotes formulas, 317
- Newton-Raphson method, 93
- Nicolson
 - Phyllis (Lockett), 513
- nonsingular matrix, 405
- operation counts, 413–414
- orthogonal matrix, 464–465, 471–472, 488, 493–496, 502
- overflow whereas, 89
- overflow, 21
- Padé approximation, 246
- parasitic solution, 350, 366–367
- Pascal, 29–30
- Peano estimate, 313
- Peano kernel, 313
- Peano
 - Giuseppe, 312
- piecewise linear interpolation, 57
- Pisano
 - Leonardo, 143
- pivots, 409, 434
- Prenter
 - Paddy, 214
- QUADPACK, 30, 307, 317
- Raphson
 - Joseph, 93
- Rayleigh quotient iteration, 487–492
- Rayleigh quotient, 485–487
- Rayleigh shift, 499–501
- relative error, 15
- Richardson
 - Lewis Fry, 290
- Riemann sum, 251, 253
- Riemann
 - Georg Friedrich Bernhard, 251
- Rolle's Theorem, 58, 175
- Rolle
 - Michel, 58
- root conditions, 367–368
- Runge example, 238–240
- Runge
 - Carl, 234, 352
 - Runge-Kutta method, 234, 352–357
 - Runge-Kutta methods, 352
 - Runge-Kutta-Fehlberg scheme, 378–382, 384–388
 - scaled pivoting, 410
 - search direction, 529
 - secant line, 119
 - secant method, 119–121
 - Seidel
 - Ludwig Philipp von, 449
 - shooting methods, 395–397
 - Simpson's rule, 257
 - Simpson
 - Thomas, 257
 - single precision, 24, 28
 - SOR (Successive Over-Relaxation) iteration, 450
 - splines, 198
 - splitting methods, 446–451, 527, 530
 - stability polynomial, 367
 - Steffenson's iteration, 135
 - Stirling's formula, 273–274
 - Stirling
 - James, 273
 - Strong root condition, 367
 - superlinear convergence, 109, 121
 - systems of equations, 74, 406, 452
 - Taylor's theorem, 2
 - Taylor
 - Brook, 2
 - Trapezoid rule, 64
 - triangular matrix, 76, 405
 - tridiagonal systems, 74
 - Two-point boundary value problem (BVP), 80
 - unconditional stability, 511
 - vector norm, 426
 - virtual memory, 444
 - von Leibniz
 - Wilhelm Gottfried, 93
 - weak stability, 346–350
 - Wilkinson shift, 499, 501
 - Wilkinson
 - James Hardy, 431