

## Index

- ABCD Rule and scoring system, 459, 460  
 abciximab, ischemic stroke, 470  
 abdominal aortic aneurysm, 581–2  
 abdominal pain  
   appendicitis, 383  
   ectopic pregnancy, 392–3  
   pelvic inflammatory disease, 422–8  
 abdominal trauma, blunt, ultrasound, 579, 582–3  
 abortion  
   spontaneous (miscarriage)  
     prediction, 436, 437–8  
     threatened, bed rest, 438  
   therapeutic, ectopic pregnancy, 400–1  
 abscess  
   peritonsillar, 523, 528–31  
   soft tissue, 571–7  
 abuse (of persons)  
   elderly, 635–43  
   intimate partner violence, 590, 594, 600–5  
 acetaminophen (paracetamol)  
   overdose, 296–300, 631  
 N-acetylcysteine (NAC), acetaminophen poisoning, 299–300  
 acetylsalicylic acid *see* aspirin  
 acid (gastric)  
   agents suppressing secretion *see* antisecretory agents  
   reflux, 161–2  
 acromioclavicular joint, 350  
   injuries/separations, 352, 355–6  
 activated charcoal *see* charcoal  
 activated protein C, 239, 245  
 acute coronary syndrome (ACS), 47, 153–7, 165–76  
   syncope and, 274–6  
 Acute Physiology and Chronic Health Evaluation (APACHE) II score  
   in sepsis and septic shock, 239, 240  
 adamantanes, influenza, 138  
 adenosine, supraventricular tachycardia, 216–18  
 adherence *see* compliance  
 adhesives, tissue, 563  
 admission to hospital  
   agitated patient, 517  
   anaphylaxis, reducing, 146–7  
   asthma  
     predicting need for, 80–1  
     reducing, 81–3  
   COPD, reducing, 93  
   elderly patients  
     comprehensive geriatric assessment and readmission rates, 265–6  
     “hospital at home” scheme as alternative, 261, 263–5  
     identifying patient subsets on discharge at risk for readmission, 266  
   heart failure (acute decompensated), observation unit vs, 196  
   pelvic inflammatory disease, 431–2  
   pneumonia, determining need for, 103–4  
   seizures, 508  
   syncope, 278–9  
 adolescents and firearms, 595  
 adrenaline *see* epinephrine  
 adrenoceptors, drugs acting at *see*  $\alpha$ -antagonists;  $\beta$ -agonists;  $\beta$ -antagonists  
 agitation (undifferentiated acute), 254–6, 512–19  
 air-stirrup ankle brace, 331  
 airway management  
   cardiac arrest, 228  
   procedural sedation and analgesia, 554  
 albumin administration  
   hemorrhagic shock, 375  
   sepsis/septic shock, 243  
 albuterol *see* salbutamol  
 alcohol (ethanol) consumption, 590, 591–2, 624–9  
   elder abuse and, 638  
   interventions, behavioral change, 591–2  
   seizures related to, 286–8  
 alfentanil, procedural sedation and analgesia, 558  
 allergic reactions, 143  
   biphasic (in anaphylaxis), 147–8  
    $\alpha$ -antagonists ( $\alpha$ -blockers)  
     rhinosinusitis, 540  
     ureteric calculi, 409  
 alteplase, acute MI, 186  
 amantadine, 138  
 ambulance diversion, 68  
 American College of Emergency Physicians (ACEP) guidelines and clinical policies  
   abuse screening, 639  
   agitation, 512  
   altered mental status, 255, 256  
   smoking cessation, 610  
   syncope, 279  
   American College of Emergency Psychiatry (AAEP) guidelines, agitation, 512  
   American College of Obstetrics and Gynecology, screening for abuse, 639  
   American Medical Association, screening for abuse, 639  
   American Psychiatric Association guidelines, delirium, 255–6  
    $\epsilon$ -aminocaproic acid, hemorrhagic shock, 377  
 aminophylline  
   cardiac arrest, 229  
   COPD, 91–2  
 amiodarone  
   atrial fibrillation, 204  
   cardiac arrest, 229  
   ventricular arrhythmias, 218–19, 221–2  
 amitriptyline, post-concussive headache, 313  
 amoxicillin  
   pelvic inflammatory disease, 431  
   rhinosinusitis, 537–8, 539  
 ampicillin, pelvic inflammatory disease, 431  
 analgesia (pain management)  
   chest pain, 162  
   hip fracture, 68  
   knee injury, 340–1  
   migraine, 495–7, 498  
   misuse of analgesic, 629–32  
   radial fracture (distal), 347  
   sedation and, procedural, 553–60  
   streptococcal pharyngitis, 528  
   ureteric calculi, 406–9  
   wounds, 565–7  
 anaphylaxis, 143–9  
 anemia, stool occult blood testing, 447–8  
 anesthetic  
   liquid, atypical chest pain, 161–2  
   local, traumatic wounds, 565–7  
 aneurysm  
   abdominal aortic, 581–2  
   subarachnoid hemorrhage due to, 477, 478, 479, 480, 481, 482

## Index

- angina, unstable (UA), 153, 155, 157, 165–6
- angiography  
 CT pulmonary, 124, 126–8  
 lower GI bleeding, 451, 452  
 pelvic, in multiple trauma, 379
- angioplasty, coronary *see* percutaneous interventions
- animal bites *see* bite wounds
- ankle injury, 13, 325–34
- antacids, atypical chest pain, 161–2
- anterior drawer test, knee injury, 336, 337
- anthrax vaccination, 620
- anti-arrhythmics  
 atrial fibrillation, 203–4, 208–10  
 cardiac arrest, 229  
 supraventricular tachycardia, 216–18  
 ventricular arrhythmia, 218–23
- antibiotics  
 appendicitis, 383–4  
 bacteriuria in pregnancy, 441–2  
 chest tube insertion, prophylactic, 368–9  
 conjunctivitis, 535–8, 543  
 COPD, 92, 94–5  
 endocarditis prophylaxis, 574–5  
*H. pylori* eradication, 447  
 meningitis, 489  
 pelvic inflammatory disease, 429–31  
   vs hospitalization, 431–2  
 pneumonia, 104–10  
   avoidance of classes prescribed within 3 months, 110  
   first-generation vs second and third line, 104–6  
   intravenous vs oral, 104  
   outpatient vs inpatient, 106–7  
 rhinosinusitis, 537–9  
 sepsis/septic shock, 239–40  
 soft tissue abscess, 572–3, 574–6  
   community-acquired MRSA, 573–4  
 streptococcal pharyngitis, 526–8  
 urinary tract infection, 415–16, 418  
 wounds, 567–8
- anticholinergics  
 in asthma, 81–2  
 in COPD, 90–1
- anticoagulation  
 acute coronary syndrome, 170–2  
 atrial fibrillation, oral agents, 205–8  
 DVT, 117–19, 120  
 PE, 128–30  
 transient ischemic attack, 462–3  
   oral agents, 461–2
- anticonvulsants  
 alcohol-related seizures, 277–8  
 eclampsia, 438–9  
 refractoriness to, 507  
 seizures, 504, 505–8
- antidepressants, smoking cessation, 608
- anti-emetics  
 migraine, 497  
 pregnancy, 439–41
- anti-epileptics *see* anticonvulsants
- antifibrinolytics, hemorrhagic shock, 377
- antigen tests, influenza, 136–8
- antihistamines *see* histamine H<sub>1</sub> or H<sub>2</sub> antagonists
- antimuscarinics, ureteric calculi, 407–8
- antiplatelet agents  
 acute coronary syndrome, 167–9, 172–3  
 acute ischemic stroke, 470–1  
 atrial fibrillation, 205–8  
 transient ischemic attack, 460–1  
   vs oral anticoagulants, 461
- antipsychotics, agitation and/or delirium, 254–7, 515–17
- antisecretory (acid suppressing) agents  
 in upper GI bleeding, 447  
 prophylaxis in NSAID use, 448–9
- antiviral drugs
- influenza, 138–9  
 prophylactic use, 139
- aorta  
 abdominal, aneurysm, 581–2  
 thoracic, dissection, 158  
 traumatic rupture, 366–7
- appendicitis, acute, 383–90
- arrhythmias/dysrhythmias, 200–24  
 antipsychotic-induced, 256–7  
 supraventricular, 215, 215–18  
 ventricular, 214–15, 216–23
- arterial oxygen saturation (SaO<sub>2</sub>)  
 in asthma, 80–1
- arthrography, MR, rotator cuff tears, 354
- arthroscopy in knee injury, MRI predicting need for, 339–40
- aspiration  
 nasogastric, upper GI bleeding diagnosis, 449–50  
 needle, peritonsillar abscess *see* needle aspiration
- aspirin (acetylsalicylic acid; ASA)  
 in acute coronary syndrome, 167–9  
 in acute MI, 181  
   fibrinolysis and heparin and, 185, 187  
 in atrial fibrillation, 205–8  
 in chest pain, 162  
 in ischemic stroke, 470–1  
 in transient ischemic attack, 460–1, 461  
*see also* salicylate poisoning
- asthma  
 exacerbations, 79–88  
 medication adherence, 49, 53, 55  
 pediatric, 55
- atenolol  
 acute MI, 182  
 atrial fibrillation, 200–13  
 atrial fibrillation, 200–13
- atropine in cardiac arrest, 229
- audit, physician, 45
- AUDIT survey, 591, 592, 627
- automated defibrillators, 227
- avian influenza, 134, 139
- azithromycin  
 pelvic inflammatory disease, 430, 432  
 soft tissue abscess, 575  
 streptococcal pharyngitis, 528
- bacitracin/polymyxin ointment  
 conjunctivitis, 546  
 wounds, with neomycin, 567
- bacterial endocarditis prophylaxis, 574–5
- bacteriuria in pregnancy, 441–2
- bag mask ventilation in cardiac arrest, 228
- bandages, elastic, ankle injury, 330
- Banff guideline, 356
- barbiturates, seizures, 507  
 alcohol-related, 288
- bed management strategies, trials of variations in, 71
- bed rest, threatened abortion, 438
- bedside evidence-based medicine, 18–19
- bedside ultrasonography, abdominal aortic aneurysm, 581–2
- behavior  
 as barrier to evidence uptake, 15  
 health-related, change, 589–90  
 safety belts, 594, 595  
 smoking cessation, 610
- beneficial intervention, evidence for, 18–19
- benzodiazepines  
 agitation, 254–6, 515–17  
 seizures, 505–7  
   alcohol-related, 288, 290
- benzylamine hydrochloride, streptococcal pharyngitis, 528
- best evidence, 4
- β-agonists  
 asthma, inhaled, 81–2, 82  
 COPD, inhaled, 90–1  
   with anticholinergics, 91  
 long-acting, 85–6
- β-antagonists (β-blockers)  
 acute coronary syndrome, 169–70  
 acute MI, 181–2  
 atrial fibrillation, 202, 204
- bicarbonate (sodium)  
 IV  
   cardiac arrest, 229–30  
   salicylate poisoning, 301  
 local anesthetic buffering with, for wounds, 566–7
- bi-level positive airway pressure, acute decompensated heart failure, 193–4
- biochemical tests and biomarkers *see* laboratory tests
- bite wounds (animals incl. dogs), 561, 562–3, 567–8  
 rabies immunization, 618
- bleeding *see* hemorrhage
- blood cultures, pneumonia, 102–3
- blood pressure  
 abnormal *see* hypertension; hypotension  
 management in subarachnoid hemorrhage, 480
- blood testing  
 peripheral, in bacterial meningitis, 486–7  
 stool occult, in anemia, 447–8
- blood transfusion  
 sepsis/septic shock, 241, 245  
 upper GI bleeding, 450
- bone scan, scaphoid fracture, 345–6
- brace, ankle, 331, 332
- brain  
 alcohol-related changes, 286  
 imaging *see* neuroimaging  
 trauma, mild, 307–15
- brain-derived natriuretic peptide (BNP) in acute decompensated heart failure  
 diagnostic value, 191–2  
 recombinant, administration, 194–5

- breathing in procedural sedation and analgesia  
 spontaneous, 554  
 supplemental O<sub>2</sub> vs room air, 556–7  
 breathlessness *see* dyspnea  
 brief interventions (BIs) on alcohol use, 627, 628–9  
 bronchodilators  
 asthma, 81–2  
 COPD, 90–1  
 Brudzinski sign, 487  
 bundle branch block, left, 160  
 bundled therapy in sepsis/septic shock, 241–3  
 bupropion, 608, 609  
 Buscopan, ureteric calculi, 408
- C-reactive protein (CRP)  
 appendicitis, 385, 386  
 rhinosinusitis, 535  
 CAGE questionnaire, 591, 625, 626, 627  
 calcium channel blockers  
 atrial fibrillation, 202  
 subarachnoid hemorrhage, 480–1  
 supraventricular tachycardia, 216–18  
 ureteric calculi, 409  
 calculi, ureteric, 404–11  
 calf veins and thrombosis risk, 113–14, 460  
 California Rule, 459  
 California Smokers' Quitline, 610  
 Canadian Agency for Drugs and Technologies in Health (CADTH) study, procedural sedation and analgesia, 557–8, 558  
 Canadian C-spine Rule (CCR), 318, 319, 320  
 Canadian CT Head Rule, 308, 309, 311  
 Canadian Sinusitis Symposium, 539  
 Canadian Task Force on Preventive Health Care, elder abuse screening, 639  
 Canadian Triage and Acuity Scale (CTAS) *see* triage scales  
 capnography, procedural sedation and analgesia, 557  
 carbamazepine, alcohol-related seizures, 277–8  
 carbon dioxide, end-tidal, in procedural sedation and analgesia, 557  
 carbon monoxide poisoning, 288–93  
 carboxyhemoglobin, 288, 290, 291–3  
 carbuncle, 575  
 carcinogenic risk of CT to fetus, 388  
 cardiology *see* heart  
 cardiopulmonary resuscitation (CPR), 225  
 bystander, 226  
 futile efforts, patient subset, 230–1  
 cardioversion  
 electrical  
 in atrial fibrillation, 201–2, 204–5, 208–10  
 in ventricular tachycardia, 221–3  
 pharmacological  
 in atrial fibrillation, 203–4, 208–10  
 in supraventricular tachycardia, 216–18  
 care (patient)  
 clinical practice guidelines and their implementation  
 improving, 15–16  
 knowledge translation methods and innovative continuing medical education improving, 16  
 elderly *see* elderly  
 overcrowding and its impact on, 67–72
- caregiver (of elder)  
 abuse by, 637, 638, 639  
 screening, 639  
 satisfaction scores with "hospital at home" scheme, 263–5  
 carotid stenosis  
 carotid endarterectomy following TIA, 461  
 ultrasound detection, 463  
 CASCADE, 221  
 case management with medication non-adherence, 54–5  
 case scenario/vignette (in chapter organization), 6  
 cast(s)  
 ankle injury, 329–30, 331  
 wrist  
 distal radial fracture, 347–8  
 scaphoid fracture, 345  
 CAST, 470  
 causal factor charting in root cause analysis, 46  
 CDC *see* Centers for Disease Control and Prevention  
 CDP-choline, post-concussive headache, 312, 313  
 Centers for Disease Control and Prevention (CDC)  
 gun control and, 596  
 rhinosinusitis guidelines, 536  
 Centor (and modified Centor) score, 525  
 Central Register of Controlled Trials (CENTRAL), 4  
 central venous cannulation, 578  
 ultrasound-guided, 579, 580–1  
 central venous pressure monitoring in sepsis/septic shock, 241  
 cephalosporins  
 pelvic inflammatory disease, 430, 432  
 pneumonia, 106  
 rhinosinusitis, 538–9  
 streptococcal pharyngitis, 528  
 cerebrospinal fluid  
 analysis  
 bacterial meningitis, 486–7  
 subarachnoid hemorrhage, 478–80  
 culture, bacterial meningitis, 486–8  
*see also* lumbar puncture  
 cerebrovascular accident *see* stroke  
 cerebrovascular disease, 457–74  
 cervical spine injury, 316–24  
 CHADS<sub>2</sub> (Cardiac Failure, Hypertension, Age, Diabetes, Stroke [Doubled]) scoring system, 206  
 charcoal, activated, 283, 283–6  
 acetaminophen, 298–9  
 salicylate, 300–1  
 CHARISMA, 471  
 chemical restraint, agitation, 513–14, 515–17  
 chest compression ± ventilation, 226  
 chest pain, 153–64  
 atypical, 161–2  
 repeat, reduction in acute MI, 180–5  
 chest pain unit, admission in acute coronary syndrome, 47  
 chest trauma, 363–72  
 chest tube insertion, pneumothorax, 367–9  
 chest X-ray/radiography  
 blunt chest trauma, 366–7, 369  
 pneumonia, 102, 109–10  
 thoracic aortic dissection, 158  
 chickenpox (varicella) immunization schedules, 616  
 Chief Emergency Officer (CEO) and overcrowding, 65
- children  
 asthma medication adherence, 55  
 firearms and, 595  
 immunization  
 intramuscular, pain reduction, 620–1  
 schedules, 615  
 soft tissue abscess, 573  
 tobacco smoke exposure, 606  
 Chinese Acute Stroke Trial, 470  
*Chlamydomphila* and *Chlamydia*, 100, 414, 421  
*C. pneumoniae*, 105  
*C. trachomatis*, 415, 421, 430  
 chloramphenicol  
 conjunctivitis, 546  
 meningitis, 489  
 chlorpromazine, migraine, 496  
 chronic conditions, medical adherence, 49–57  
 chronic obstructive pulmonary disease (COPD), 89–99  
 cigarettes *see* smoking  
 ciprofloxacin  
 conjunctivitis, 546  
 pelvic inflammatory disease, 430, 432  
 urinary tract infection, 416, 417  
 circulation (spontaneous) in cardiac arrest, return to rate of  
 CPR alone and, 226  
 CPR and defibrillation vs pharmacotherapy and, 228–30  
 therapeutic hypothermia following, 230  
 clarithromycin, *H. pylori* eradication, 447  
 clavulanic acid/clavulanate  
 pelvic inflammatory disease, 431  
 rhinosinusitis, with amoxicillin, 538–9  
 clindamycin  
 pelvic inflammatory disease, 430, 432  
 soft tissue abscess, 575  
 streptococcal pharyngitis, 528  
 clinical epidemiology, terminology, 9–10  
 clinical examination *see* examination  
 clinical gestalt *see* gestalt  
 clinical practice guidelines and their implementation  
 strategies, 15–16  
 clinical prediction rules  
 bacterial meningitis, 486, 488  
 cardiac arrest, 231  
 deep venous thrombosis, 115  
 knee injury, 338  
 sepsis and septic shock, 239, 240  
 sore throat, 524–6  
 peritonsillar abscess, 528–9  
 syncope, 276–8, 278, 279  
 thoracic aortic dissection, 158  
 transient ischemic attack, 459  
 clopidogrel  
 acute coronary syndrome, 167–9  
 acute ischemic stroke, 471  
 acute MI, 186  
 atrial fibrillation, 208  
 closed-ended questions in assessment of medication adherence, 52  
 closed reduction  
 distal radial fracture, 347, 347–8  
 shoulder dislocation, 351–2  
 clonidine, agitation, 516

## Index

- co-amoxiclav (amoxicillin/clavulanate), rhinosinusitis, 538–9
- cocaine in topical anesthetics for wounds, 566
- Cochrane Central Register of Controlled Trials (CENTRAL), 4
- Cochrane Collaboration, 5
- systematic reviews and, 5, 6, 14, 17
- Cochrane Database of Systematic Reviews, 6, 14
- Cochrane Library, 6
- colic, Acute ureteric, 404–11
- colloids
- hemorrhagic shock, 374–5
- sepsis/septic shock, 243–4
- colonoscopy in lower GI bleeding, 450–2
- combitube in cardiac arrest, 228
- commercial entities, funding of CME, 34, 35–7
- COMMIT, 169, 182
- Committee for Procedural Quality and Evidence-Based Practice (CPQE), 19
- community-dwelling older adults, 261
- falls prevention program, 593
- “hospital at home”, 263–5
- comparisons (in clinical questions), 7
- compliance/adherence
- patient (to medication), 49–57
- physician, aids/initiatives, 19, 20
- complications *see* morbidity
- Comprehensive Geriatric Assessment model, 261, 265–6
- compression, chest,  $\pm$  ventilation, 226
- compression stockings, DVT, 119
- compression ultrasonography, DVT, 117
- computerized tomographic pulmonary angiography in PE, 124, 126–8
- computerized tomographic venography in suspected PE, 126–8
- computerized tomography (CT)
- abdominal injury (blunt), 579, 583
- appendicitis, 386–9, 389
- chest trauma
- diaphragmatic injury, 369
- pulmonary contusions, 367
- head
- bacterial meningitis, 488
- ischemic stroke, 468–9
- mild injury, 308–9, 310–12
- peritonsillar abscess, 529–30
- rhinosinusitis, 536–7
- seizure (first), 505
- subarachnoid hemorrhage, 478–9
- transient ischemic attack, 457, 458
- hemoperitoneum, 582
- multi-trauma patients, 377–8
- neck injury, 319–20, 321
- pelvic inflammatory disease, 426–7
- ureteric calculi, 405–6
- concealment of allocation, 25–6
- conclusions (in chapter organization), 6
- concussion (mild brain trauma), 307–15
- Confusion Assessment Method, 253–4
- conjunctivitis
- bacterial conjunctivitis, 545–8
- viral, 543
- connective tissue disorders and subarachnoid hemorrhage, 476–7
- contact time (CME programs), 39–40
- mandatory vs voluntary, 40
- continuing medical education/CME (and professional development), 34–42, 45
- active vs passive formats, 38
- innovative, 16
- continuing quality improvement (CQI), 19
- continuous positive airway pressure, acute decompensated heart failure, 193–4
- contraceptive pill overdose, 282–6
- contrast-enhanced CT, acute appendicitis, 387
- control (in clinical questions), 7, 8
- see also* randomised controlled trials
- contusions
- cardiac, 364–6
- pulmonary, 367
- coronary artery bypass surgery, clopidogrel and, 169
- coronary artery disease, 153–84
- smoking and, 611
- see also* acute coronary syndrome
- corticosteroids
- anaphylaxis, intravenous, 147
- asthma
- inhaled, 84–6
- with long-acting  $\beta$ -agonists, 85–6
- systemic/intravenous, 82–3
- bacterial meningitis, adjuvant use, 490–1
- COPD, systemic/intravenous, 93, 95–6
- migraine, 499
- nausea and vomiting in pregnancy, 440
- hyperemesis gravidarum, 441
- rhinosinusitis, inhaled, 539–40
- sepsis/septic shock, 246–7
- streptococcal pharyngitis, 528
- cosmetic outcome with wounds, 567
- cost (incl. cost-effectiveness) of management
- ankle injury, 329
- ectopic pregnancy, 398–9
- migraine, 498
- NSAID gastrointestinal side-effect prevention, 449
- seizures, 507–8
- wounds, 567
- co-trimoxazole *see* trimethoprim/sulfamethoxazole
- COX inhibitors *see* non-steroidal anti-inflammatory drugs
- cranberry juice, urinary tract infection, 416
- creatine phosphokinase-MB
- cardiac contusion, 364, 365
- critical appraisal/review, 14–15, 22–32, 44–7
- in chapter organization, 6
- in 5S levels of organization of research evidence, 16–17
- crowding *see* overcrowding
- cruciate ligament injury
- anterior (ACL), 335, 337, 338, 339
- posterior (PCL), 337, 339
- crystalloids
- hemorrhagic shock, 374, 375, 375–7, 378–9
- sepsis/septic shock, 243–4
- cultures
- blood, pneumonia, 102–3
- CSF, bacterial meningitis, 486–8
- CURB-65 criteria, 103
- cianoacrylate tissue adhesives, 563
- cyclo-oxygenase inhibitors *see* non-steroidal anti-inflammatory drugs
- cytidine diphosphocholine (CDP-choline), post-concussive headache, 312, 313
- D-dimer assay
- DVT, 116–17
- PE, 125
- data collection process in root cause analysis, 46
- database(s), electronic, 8, 14
- DDAVP, hemorrhagic shock, 377
- death *see* mortality; suicide
- decision (clinical)
- aids, 19
- rules
- ankle injury, 329
- DVT, 114–15
- head trauma (mild), 309–10
- neck injury, 318–19
- pneumonia, 103–4
- wrist injury, 346
- decongestants (nasal), rhinosinusitis, 540
- DEED II (Discharge of Elderly from the ED), 265
- deep vein thrombosis (DVT), 113–22, 123
- defibrillator shock
- in cardiac arrest, 227–8
- automated, 227
- in ventricular tachycardia, 221, 227–8
- biphasic vs monophasic, 227–8
- resistance to, 218–21
- delirium, 251–9
- differential diagnosis of etiologies, 252
- hallmarks/key features, 251, 252
- physical assessment, 252, 253–4
- dementia, 261
- screening/diagnosis, 262–3
- depression, 514, 517–18
- agitation and, 514, 517–18
- CME on, 40
- screening, 517–18
- desmopressin, hemorrhagic shock, 377
- dexamethasone
- asthma, 82
- bacterial meningitis, 491
- migraine, 499
- diabetes (in CHADS<sub>2</sub> scoring system), 206
- diagnostic assessment (in clinical questions), 7
- dialysis, salicylate poisoning, 301–2
- diaphragmatic injury, 369
- diastolic murmur, thoracic aortic dissection, 159
- diazepam, seizures, 506, 507
- eclampsia, 438–9
- diclofenac
- knee injury, 340–1
- ureteric calculi, 407
- diffusion-weighted MRI of head
- ischemic stroke, 469–70
- mild head injury, 312
- transient ischemic attack, 462
- digoxin, atrial fibrillation, 204
- dihydroergotamine (DHE)
- migraine, 495, 496, 496–7
- post-concussive headache, 313

- diltiazem  
atrial fibrillation, 202, 204  
supraventricular tachycardia, 217–18
- diphtheria *see* tetanus immunization, with  
diphtheria/pertussis
- dipstick test for urinary tract infection, 414–15
- dipyridamole, transient ischemic attack, 460–1, 461
- disability following distal radial fracture, reducing, 347–8
- discharge from hospital/ED  
acute decompensated heart failure, safety predictors, 195–6  
asthma, relapse following, 83–6  
cardiac arrest and survival to, 226–7, 228–30  
elderly prior to  
comprehensive geriatric assessment, 265–6  
polypharmacy screening and impact on adverse effects, 265  
prognostic screening tool, 266  
follow-up after *see* follow-up  
head injury (mild), safety criteria, 310–12  
headache following, in acute migraine, 499  
seizures, safety considerations, 507–8  
syncope, 278–9  
*see also* home
- dislocations, shoulder, 350–2, 356–7
- diuretics  
acute decompensated heart failure, 192–3  
ureteric calculi, 408–9
- divalproex sodium, post-concussive headache, 313
- dobutamine, sepsis/septic shock, 241
- doctor *see* physician
- dofetilide, atrial fibrillation, 204
- dog bite *see* bite wounds
- domestic violence, 590, 594, 600–5
- dopamine antagonists  
migraine, 497  
nausea and vomiting in pregnancy, 440
- Doppler sonography, pelvic inflammatory disease, 424–6
- doxycycline, pelvic inflammatory disease, 430, 431, 432
- drainage *see* incision and drainage
- dramamine, nausea and vomiting in pregnancy, 440
- drawer tests, knee injury, 336
- droperidol  
as anti-emetic  
migraine, 497  
in pregnancy, 440  
delirium/agitation, 256–7, 515–16
- drug misuse, 624–34
- drug overdose, 282–6, 296–303, 629–32
- drug therapy (medication; pharmacotherapy)  
adherence, 49–57  
in cardiac arrest, 228–30  
associated with shock-resistant ventricular arrhythmia, 218–21  
in delirium/agitation, 254–7  
elderly, polypharmacy screening prior to discharge and impact on adverse effects, 265  
*see also* specific (types of) drugs and disorders/diseases
- duodenal ulcer, 447, 449, 452
- dyspnea (breathlessness)  
in acute decompensated heart failure, 191  
in PE, 123, 130
- dysrhythmias *see* arrhythmias
- ear, nose and throat, 521–49
- early goal-directed therapy, sepsis/septic shock, 241–3
- ECG *see* electrocardiography
- echocardiography, cardiac contusion, 364–6
- eclampsia, 438–9
- ECST, 461
- ectopic pregnancy, 391–403
- education  
continuing medical *see* continuing medical education  
patient, 19  
asthma, 86  
COPD, 96–7  
medication adherence, 53–4  
soft tissue injury of neck and associated long-term pain, 322  
safety belt wearing in pregnancy, 595  
*see also* information
- effusion, pericardial, 369–70
- elastic bandages, ankle injury, 330
- elderly/older patients, 260–70, 635–43  
abuse, 635–43  
delirium  
diagnostic tests, 253–4  
in hospitalized patients, prevention, 257–8  
falls *see* falls  
immunization  
pneumococcus, 616, 618–19  
schedules, 616  
tetanus with contaminated wound, 617–18
- electrical cardioversion *see* cardioversion
- electrical shock in ventricular arrhythmia *see* defibrillator  
shock
- electrocardiography (ECG)  
acute MI, 179–80  
chest pain, 158–60  
chest trauma, 364–6  
syncope, 274, 276, 277  
ventricular/supraventricular tachycardias with aberrant conduction, 215–16
- electrolyte assessment, first seizure, 504–5
- electronic databases, 8, 14
- electronic medication monitors, 49
- electronic triage, 63
- embolism  
peripheral, in atrial fibrillation, reducing rate of, 208–10  
pulmonary, 114, 123–32
- embolization, pelvic bleeding in multiple trauma, 379
- emergency (emergencies), definition, 44
- emergency department  
overcrowding *see* overcrowding  
triage *see* triage
- emergency medical services  
definition, 44  
response (in cardiac arrest) vs public access automated defibrillators, 227
- Emergency Severity Index (ESI), 59, 60, 62
- emergency treatment, definition, 44
- emesis *see* anti-emetics; vomiting
- EMLA, wounds, 565
- end-tidal CO<sub>2</sub> in procedural sedation and analgesia, 557
- endocarditis (infective), prophylaxis, 574–5
- endoscopy in GI bleeding  
lower, 450–2  
upper, immediate vs delayed, 448
- endotracheal intubation *see* intubation
- enoxaparin  
acute coronary syndrome, 80, 170, 171, 174  
in acute MI, 180  
as adjunct to fibrinolysis, 185–7  
in VTE (DVT/PE), 120, 129
- ENT, 521–49
- epileptic vs non-epileptic seizures, 503  
*see also* seizures
- epinephrine (adrenaline)  
in anaphylaxis, 144–6  
self-injectable, 148  
cardiac arrest, 229  
in local anesthetics for wounds, 566
- eptifibatid, ischemic stroke, 470
- erythrocytes (red cells)  
sedimentation rate (ESR), rhinosinusitis, 535  
tagged cell test in lower GI bleeding, 451
- erythromycin, streptococcal pharyngitis, 528
- ethanol *see* alcohol
- etomidate in procedural sedation and analgesia, 557–8
- European Society of Cardiology guidelines, syncope, 279
- eutectic mixture of local anesthetics (EMLA), wounds, 565
- evidence  
barriers to uptake of, 14–15  
for beneficial or harmful intervention, 18–19  
best, 4  
collecting and interpreting, 10–11  
high-quality *see* high-quality evidence  
improving sensibility and use of, 16–17  
improving uptake of, 14–17  
levels *see* levels of evidence  
locating, 8  
pyramidal organization of, 16–18
- evidence-based (emergency) medicine (EBM; EBEM)  
definition/use of term, 3  
reasons for using/value of, 4
- examination, clinical/physical  
abuse  
elderly, 636–8  
intimate partner, 601
- appendicitis, 385
- bacterial meningitis, 487
- chest pain, 155–6, 158
- conjunctivitis, 544–5
- delirium, 252, 253–4
- heart failure (acute decompensated), 191–2
- injury  
knee, 336–8  
neck, 318, 321  
rotator cuff, 353–4
- pelvic inflammatory disease, 423
- in pregnancy  
ectopic pregnancy possibility, 392–3  
fetal loss prediction, 437–8
- rhinosinusitis, 535–6
- soft tissue abscess, 572
- sore throat, 524–6  
peritonsillar abscess, 528–9
- subarachnoid hemorrhage, 477

## Index

- exercise (physical) and second impact syndrome, 312–13  
 exposure (in clinical questions), 7  
 external rotation (in shoulder dislocation), 357, 359  
 immobilization in, 358  
 eye emergencies, 543–9
- facilitated management, distal radial fracture, 348  
 factor VIIA, recombinant, hemorrhagic shock, 376–7  
 factor Xa inhibitors, PE, 130  
 falls  
 elderly, 260, 266, 589, 590, 592–3, 596  
 clinical scenarios, 266, 269, 589, 596  
 prevention, 263, 592–3, 596  
 fasting in procedural sedation and analgesia, 554–6  
 fast-track systems, 69–70  
 fatal flaws in scientific papers, missing, 28–9, 29–30  
 fecal occult blood testing in anemia, 447–8  
 feedback, physician, 45  
 females *see* women  
 femoral veins and thrombosis risk, 113–14  
 fertility problems, pelvic inflammatory disease, 431–2  
 fetus  
 CT risk to, 388  
 loss *see* abortion  
 fibrillation  
 atrial, 200–13  
 ventricular (VF), 214–15, 216–21  
 fibrinolysis inhibitors, hemorrhagic shock, 377  
 fibrinolytics *see* thrombolytics  
 financial abuse of elderly, 636, 638  
 firearm injury prevention, 590, 595–6  
 5S levels of organization of research evidence, 16–18  
 flecainide, atrial fibrillation, 204  
 fludrocortisone, sepsis/septic shock, 246  
 fluids  
 bacterial meningitis, administration vs restriction, 489–90  
 resuscitation with  
 hemorrhagic shock, 373, 374–7, 378–9  
 sepsis/septic shock, 241, 243–4  
 ureteric calculi passage and administration of, 408–9  
 fluoroquinolones  
 pelvic inflammatory disease, 430  
 pneumonia, 106, 107  
 urinary tract infection, 416  
 focused assessment with sonography (FAST)  
 blunt abdominal trauma, 579, 582–3  
 multi-trauma patients, 377–8  
 folate inhibitors, rhinosinusitis, 537–8  
 follow-up (outpatient/discharge patients)  
 agitated patient, 517  
 in community-acquired pneumonia, 108–10  
 older patients, 266  
 fondaparinux, PE, 130  
 forced expiratory volume in one second (FEV<sub>1</sub>) in  
 asthma, 80–1  
 foreign body, rectum, 553, 559  
 fosphenytoin, 507–8  
 4-hour rule (UK), 71  
 fracture  
 ankle, 325, 326–9  
 C-spine (possibility of), 316, 317, 318, 319, 321, 322  
 hip, analgesia, 68  
 shoulder region, 356–7  
 iatrogenic, 356–7  
 proximal humerus, 352–3, 357–60  
 wrist, 345–8  
 distal radius *see* radius  
 full-capacity protocol, 71–2  
 functional outcome  
 in elderly patients following discharge, identifying  
 patients at risk of decline, 266  
 migraine, 499  
 shoulder dislocation, 357–60  
 wrist injury, 346–7  
 funding of CME by commercial vs non-commercial  
 entities, 34, 35–7  
 furosemide, acute decompensated heart failure, 193  
 furuncle, 575  
 fusidic acid, conjunctivitis, 546
- GABA receptors and alcohol, 286  
 gastric problems/medicine *see* stomach  
 gastro-esophageal reflux, 161–2  
 gastrointestinal decontamination, 282–6  
 acetaminophen, 298–300  
 salicylates, 300–1  
 gastroscopy in upper GI bleeding, immediate vs delayed,  
 448  
 general practice *see* primary care  
 genetic thrombophilias, 115, 116  
 genitourinary problems, 391–444  
 gentamicin  
 conjunctivitis, 546  
 pelvic inflammatory disease, 432  
 gestalt, clinical, 30  
 PE, 130  
 glenohumeral joint, 350  
 Global Registry of Acute Coronary Events (GRACE) risk  
 scores, 156–7, 167  
 gloves, sterile vs non-sterile, wound management, 564–5  
 glucose, blood  
 control (glycemic control), sepsis/septic shock, 246  
 first seizure, 504–5  
 glyceryl trinitrate *see* nitroglycerin  
 glycoprotein IIb/IIIa inhibitors  
 acute coronary syndrome, 172–3  
 acute ischemic stroke, 470  
 gonorrhea (*N. gonorrhoeae* infection), 414, 415, 420, 421,  
 430  
 GRACE risk scores, 156–7, 176  
 Grades of Recommendation Assessment, Development  
 and Evaluation (GRADE) system, 4  
 granisetron, migraine, 497  
 gun injury prevention, 590, 595–6
- H<sub>1</sub> and H<sub>2</sub> antagonists *see* histamine H<sub>1</sub> antagonists;  
 histamine H<sub>2</sub> antagonists  
*Haemophilus influenzae*, 100  
 in COPD, 95  
 type b, pediatric immunization schedules, 615  
 haloperidol, delirium/agitation, 255, 256, 257, 515, 516  
 hand lacerations (dog bite), 562–3
- hangman's fracture, 317  
 harmful intervention, evidence for, 18–19  
 head  
 imaging *see* neuroimaging  
 trauma, mild, 307–15  
 headache, 493–502  
 primary, 493–502  
 secondary  
 bacterial meningitis, 487  
 post-concussive, 312  
 subarachnoid hemorrhage, 475  
 healing by secondary intention, 562–3  
 health  
 patient, CME effects on outcome, 39  
 public, 587–643  
 heart, 151–233  
 arrest (out-of-hospital), 225–33  
 associated with shock-resistant ventricular  
 arrhythmia, 218–21  
 therapeutic hypothermia following, 22, 30,  
 230  
 contusion, diagnosis, 364–6  
 disease, 151–233  
 coronary *see* coronary artery disease  
 smoking and, 611  
 failure, acute decompensated, 190–9  
*see also* CHADS<sub>2</sub>  
 rhythm disturbances, 200–24  
*see also* myocardium  
 helical CT, acute appendicitis, 386–9  
*Helicobacter pylori* eradication therapy, 447  
 hemagglutinin and influenza subtype, 133  
 hemodialysis, salicylate poisoning, 301  
 hemodynamic parameters  
 anaphylaxis, 147  
 sepsis/septic shock, optimization, 241–3  
 hemopericardium, 370  
 hemoperitoneum, 582  
 hemo(pneumo)thorax, 368–9  
 hemorrhage (bleeding)  
 atrial fibrillation therapy and risk of, 210  
 GI *see* gastrointestinal bleeding  
 major, shock due to, 373–80  
 subarachnoid, 475–84  
 vaginal, first trimester, 392–3, 436, 437–8  
 hemostatic agents, hemorrhagic shock, 377  
 Hendey algorithm, 356  
 heparin  
 acute coronary syndrome, 170–2  
 acute MI, as adjunct to fibrinolysis, 185–7  
 DVT, 117–18  
 low molecular weight (LMWH), 117–18, 120  
 unfractionated (UFH), 117–18  
 PE, 128–30  
 transient ischemic attack, 462–3  
 hepatitis A immunization, 620  
 schedules, 616  
 children, 615  
 hepatitis B immunization schedules, 616  
 children, 615  
 hepatotoxicity (and need for transplantation),  
 acetaminophen, 298–9  
 herpes zoster immunization schedules, 616  
 heterogeneity among pooled statistics, 9–10

- high-quality evidence, identifying  
in systematic review, 26–8  
in therapeutic scientific paper, 25–6
- high-quality reports, criteria used by editors for ensuring, 29–30
- hip fracture, analgesia, 68
- histamine H<sub>1</sub> antagonists (antihistamines)  
anaphylaxis, 146–7, 147  
nausea and vomiting in pregnancy, 440
- histamine H<sub>2</sub> antagonists  
anaphylaxis, 146–7  
NSAID-induced gastroduodenal ulcer prevention, 449
- history-taking  
abuse  
elderly, 636–8  
intimate partner, 602  
alcohol problems, 627–9  
bacterial meningitis, 487  
chest pain, 155–6, 158  
conjunctivitis, 544–5  
delirium, 253–4  
heart failure (acute decompensated), 191–2  
neck injury, 321  
pelvic inflammatory disease, 423  
pneumonia, 101–3  
in pregnancy  
ectopic pregnancy possibility, 392–3  
fetal loss prediction, 437–8  
shoulder injury, 353–4  
sore throat, 524–6  
peritonsillar abscess, 528–9  
subarachnoid hemorrhage, 476–7
- holding chambers, inhalers with, in asthma, 81
- home  
anticoagulation therapy at  
DVT, 118–19  
PE, 128–9  
discharge from hospital to *see* discharge  
elderly and hospital in home (HIH) care model, 261, 263–5  
medication adherence and visits to, 54  
polypharmacy screening in elderly and impact on adverse events at, 265  
pregnancy test, 435, 437
- Hopkins' Verbal Learning Test, 262
- hospital(s)  
admission to *see* admission  
discharge from *see* discharge  
elderly patients in  
comprehensive geriatric assessment and subsequent hospitalization rates, 265–6  
delirium prevention, 257–8  
length of stay *see* length of stay  
overcrowding *see* overcrowding  
resources, conservation in acute decompensated heart failure, 196
- hospital emergency services, definition, 44
- hospital in home (HIH) care model, 261, 263–5
- β-human chorionic gonadotrophin and ectopic pregnancy, 393–5
- human papillomavirus (HBV) vaccination, 616  
pediatric, 615
- humeral fracture, proximal, 352–3, 357–60
- Hwalek—Sengstock elder abuse screening test, 639
- hydrocortisone  
COPD, 92  
sepsis/septic shock, 246, 247
- 5-hydroxytryptamine (serotonin) 1B/1D antagonist, migraine, 496
- hydroxyzine, nausea and vomiting in pregnancy, 440
- hyoscine butylbromide, ureteric calculi, 408
- hyoscyamine sulfate, ureteric calculi, 408
- hyperbaric oxygen, CO poisoning, 291, 291–3
- hyperemesis gravidarum, 440, 441
- hypertension  
chronic thromboembolic pulmonary (CTPH), 123  
systemic  
aortic dissection and history of, 159  
subarachnoid hemorrhage and, 480  
*see also* CHADS<sub>2</sub>
- hypertonic saline in hemorrhagic shock, 375–6
- hypotension in anaphylaxis, 144–5
- hypothermia, therapeutic, post-cardiac arrest, 22, 30, 230
- iatrogenic shoulder fracture (due to attempted reduction), 356–7
- ibutilide, atrial fibrillation, 204
- Identification of Seniors at Risk tool (ISAR), 266, 268
- imaging  
abdominal aortic aneurysm, 581–2  
appendicitis, 386–9, 389  
carotid, 463  
chest trauma, 366–7, 369–70  
head injury (mild), 308–9, 310–12  
knee injury, 338–40  
neck injury, 318–21  
PE, 126–8  
pelvic inflammatory disease, 422–8  
peritonsillar abscess, 529–30  
pregnancy (ectopic vs intrauterine), 395–400  
rhinosinusitis, 536–7  
rotator cuff tears, 354  
soft tissue abscess, 572  
thoracic aortic dissection, 158  
transient ischemic attack, 457, 458  
ureteric calculi, 405–7  
wrist injury, 345–6  
*see also specific modalities*
- immobilization  
ankle injury, 329–30  
knee injury, 341–2  
shoulder dislocation, 357–8  
wrist  
distal radial fracture, 347  
scaphoid fracture, 345, 346–7
- immunization, 614–23  
active *see* vaccination  
passive *see* immunoglobulin
- immunodiagnosis (antigen tests), influenza, 136–8
- immunoglobulin, IV (passive immunization with)  
hepatitis A, 620  
rabies, 618
- incision and drainage  
peritonsillar abscess, 530–1  
antibiotic pre-treatment prior to, 574–5  
soft tissue abscess, 472
- indomethacin  
knee injury, 340–1  
ureteric calculi, 407
- industrial/commercial entities, funding of CME, 34, 35–7
- infections  
chest tube-related, 368–9  
conjunctival, 543–9  
immunization *see* immunization  
lung, 100–12  
meningeal, 485–92  
nasal/paranasal sinus, 524–42  
pelvic (of upper genital tract), 420–34  
renal pelvis, in pregnancy, 441–2  
soft tissue, 571–7  
systemic *see* sepsis  
throat, 523–33  
urinary tract, 412–19  
in pregnancy, 441–2  
wounds, and its prevention, 562–5, 567–8
- infertility, pelvic inflammatory disease, 431–2
- inflammation, laboratory markers (in appendicitis) of, 385–7
- influenza, 133–45  
vaccination, 139–40, 616  
in COPD, 96  
pediatric, 615
- influenza-like illness, 134
- informant questionnaire (dementia), 262
- information (for patients)  
on alcohol use, brief interventions, 627, 628–9  
on medication adherence, 53, 55  
on smoking cessation, 607–8  
*see also* education
- inhalers in asthma, 81–2
- inherited thrombophilias, 115, 116
- injury (traumatic), 305–80, 587–643  
alcohol and risk of, 624, 625–7  
ankle *see* ankle injury  
brain, mild, 307–15  
chest, 363–72  
knee, 335–43  
multiple/major, hemorrhagic shock, 373–80  
neck, 316–24  
non-accidental  
of elderly *see* physical abuse  
firearms, 595  
by intimate partner, 590, 594, 600–5  
older person, accidental, 637  
fall-related *see* fall  
older person, non-accidental, 635–43  
pelvic, 378–9  
prevention, 587–643  
shoulder, 350–62  
wounds *see* wounds  
wrist, 344–9  
*see also* fracture
- inotropes, sepsis/septic shock, 241
- instability, ankle  
chronic, 332–3  
reducing, 329–31
- internal rotation, shoulder immobilization, 358
- International Headache Society's Classification of Headache Disorders, 493, 494
- internet (web)-based CME, 38–9

## Index

- intervention (in clinical questions), 7, 8  
intimate partner violence, 590, 594, 600–5  
intracranial abnormalities (in mild brain trauma) on CT requiring surgery, 308–9  
intramuscular vaccine injection in children, pain reduction, 620–1  
intravenous pyelography (IVU), calculi, 405, 405–6  
intravenous urography (IVU), calculi, 405, 405–6  
intubation (endotracheal)  
in acute decompensated heart failure, decreasing rates of, 193–4  
in cardiac arrest, 228  
in salicylate poisoning, 300  
invasive revascularization procedures *see* percutaneous interventions  
ipratropium bromide, inhaled  
asthma, 81–2  
COPD, 90–1, 92  
with  $\beta$ -agonists, 91  
irrigation, wound, with tap water vs other solutions, 563–4  
ISAR (Identification of Seniors at Risk tool), 266, 268  
ischemia, myocardial/cardiac  
chest pain in, 155–7, 158–61  
syncope and, 274–6  
ischemic attack, transient (TIA), 457–65  
ischemic stroke, acute, 468–72  
isorbide dinitrate, acute decompensated heart failure, 193  
Jadad scale, 25–6  
Jefferson's fracture, 317  
 $\kappa$ -statistic, triage systems, 60  
Kernig sign, 487  
ketamine, procedural sedation and analgesia, 558–9  
ketofol, procedural sedation and analgesia, 558  
ketorolac  
migraine, 496, 498  
ureteric calculi, 407  
kidney function in acute decompensated heart failure, nesiritide effects, 194–5  
*see also* pyelography; pyelonephritis  
knee injury, 335–43  
knowledge  
and barriers to uptake of evidence, 15  
CME effects on uptake and retention of, 37–9  
translation, 13–21  
models for understanding of, 16  
Kocher reduction, shoulder dislocation, 356, 357, 359  
laboratory tests (incl. biochemistry and biomarkers)  
appendicitis, 385–7  
bacterial meningitis, 486–7  
brain trauma, 212–13  
cardiac contusion, 364–6  
pelvic inflammatory disease, 422, 423–4  
rhinosinusitis, 535  
seizure (first), 504–5  
sepsis/septic shock, 239  
lacerations, repair, 561–70  
Lachman test, 336–7  
 $\beta$ -lactam antibiotics  
pneumonia, 106  
wound infection prophylaxis, 568  
lactate levels in sepsis/septic shock, 239  
laparoscopic salpingostomy, ectopic pregnancy, 400–1  
laparoscopy, pelvic inflammatory disease, 421, 422  
laparotomy, blunt abdominal trauma, 582–3  
laryngeal mask airway in cardiac arrest, 228  
law on abuse  
elderly, 640–1  
intimate partner violence, 603–4  
Leach classification, ankle sprain, 325  
learning (physician), CME producing changes in, 39  
left bundle branch block, 160  
legal issues *see* law  
*Legionella pneumophila*, 105–6, 106, 107  
length of stay (hospitalization time; LOS)  
COPD, reducing, 93  
delirium, 254  
ectopic pregnancy, 398–401  
elderly in “hospital at home” scheme and its impact on, 263–5  
overcrowding and, 66–7  
torso trauma, 583  
leukocytes *see* white blood cells  
leukotriene antagonists in asthma, 86  
levels of evidence, 5–6  
systematic reviews and, 5  
lidocaine  
cardiac arrest, 229  
ventricular arrhythmias  
shock-resistant, 218–21  
stable, 221–2  
wound management, 566  
buffered vs non-buffered, 566–7  
ligaments, shoulder, 350  
literacy and medication adherence, 52–3  
literature (research)  
critical review *see* critical appraisal  
high-quality in *see* high-quality  
missing “fatal flaws” in scientific papers, 28–9, 29–30  
search, 6, 8, 14  
liver toxicity and transplantation need, acetaminophen, 298–9  
local anesthetics, traumatic wounds, 565–7  
lomefloxacin, conjunctivitis, 546  
long QT interval, 214, 223  
lorazepam  
delirium/agitation, 255, 256, 257, 515, 516  
seizures, 506, 507  
alcohol-related, 288, 290  
low birth weight and maternal bacteriuria, 441–2  
lower extremity  
imaging in suspected PE, 126–8  
venous anatomy and thrombosis risk, 113–14  
lumbar puncture/LP (for CSF analysis)  
in bacterial meningitis, 488–9  
in subarachnoid hemorrhage, 478, 479  
*see also* cerebrospinal fluid  
lung (pulmonary organ)  
chronic obstructive disease (COPD), 89–99  
contusion, 367  
function  
in asthma, assessment, 80–1  
in COPD, treatment improving, 90–3  
infections, 100–12  
opioid overdose-induced acute injury, 632  
McMurray test, 337, 338  
macrolides, pneumonia, 106, 107  
magnesium sulfate, intravenous/parenteral  
asthma, 79, 82  
eclampsia, 438–9  
migraine, 498  
torsades de pointes, 222–3  
magnetic resonance arthrography, rotator cuff tears, 354  
magnetic resonance imaging  
appendicitis in pregnancy, 389  
head  
injury (mild), 312  
ischemic stroke, 468, 469, 469–70  
seizure (first), 505  
transient ischemic attack, 457, 458, 462  
knee injury, 339–40  
neck injury, 321  
pelvic inflammatory disease, 427–8  
rotator cuff tears, 354  
scaphoid fracture, 345–6  
Manchester Triage Scale (MTS) *see* triage scales  
measles/mumps/rubella immunization schedules, 616  
pediatric, 616  
mechanical ventilation in sepsis/septic shock, 247  
meclizine, nausea and vomiting in pregnancy, 440  
medical record reviews, 45–6  
medication *see* drug therapy  
MEDS (Mortality in Emergency Department Sepsis)  
score, 239, 240  
meningitis  
bacterial, 485–92  
viral,  
meningococcal immunization schedules,  
adult, 616  
pediatric, 615  
meniscal injury, 336, 337, 338, 339  
menstruation, first missed, 437  
mental illness  
abusers of elderly, 638  
firearm use and, adolescents, 595  
*see also* psychiatric assessment  
mental status  
altered, 251–9  
in bacterial meningitis, 487  
assessment, 253–4  
meperidine  
migraine, 495–6  
ureteric calculi, 406–7  
meropenem, pelvic inflammatory disease, 432  
methicillin-resistant *S. aureus*, community-acquired,  
571–2, 573–4  
methohexital, procedural sedation and analgesia,  
558  
methotrexate, ectopic pregnancy, 400–1  
methotrimeprazine, migraine, 497  
methylprednisolone  
asthma, 79  
COPD, 92  
methylxanthines,  
COPD, 91–2  
metoclopramide  
migraine, 497, 498  
post-concussive headache, 313  
in pregnancy, 440

- metoprolol  
acute coronary syndrome, 165, 169  
acute MI, 182  
atrial fibrillation, 202
- metronidazole, pelvic inflammatory disease, 430
- midazolam  
delirium/agitation, 255, 257  
procedural sedation and analgesia, 558
- migraine, 3, 495–8, 499
- Milch reduction, shoulder dislocation, 356, 357, 359
- Mini Mental Status Examination (MMSE), 253  
elderly, 262
- minor procedures, 551–85
- miscarriage *see* abortion
- misoprostol, NSAID-induced gastroduodenal ulcer prevention, 449
- montelukast, 86
- Moraxella catarrhalis*, 100
- morbidity/complications  
abdominal aortic aneurysm, 581–2  
appendicitis, 384–5  
atrial fibrillation, 200  
prevention, 205–10  
central venous cannulation, 580–1
- cerebrovascular disorders/disease  
acute ischemic stroke, 470–2  
subarachnoid hemorrhage, 480–1  
transient ischemic attack, 460–1, 462–3
- coronary disease  
acute coronary syndrome, 167–74  
acute MI, reduction, 185–7
- delirium, 254
- DVT and anticoagulants and, 118–19
- knee injury, reduction, 342–3
- overcrowding and its impact on, 67–8
- PE and  
and diagnostic algorithms, 130  
and heparin, 128  
pelvic inflammatory disease, 431–2  
pneumothorax (traumatic), 367–8  
in procedural sedation and analgesia, 554–7  
rhinosinusitis, 537–40  
streptococcal pharyngitis, 526–8
- mortality (death)  
abdominal aortic aneurysm, 581–2  
acetaminophen poisoning, 299–300  
agitated patient, 517–18  
anaphylaxis, 145–6  
appendicitis (acute), 384–5  
atrial fibrillation, 208–10  
bacterial meningitis, 489–91  
cerebrovascular disorders  
acute ischemic stroke, 470–2  
subarachnoid hemorrhage, 480–2  
transient ischemic attack, 460–1, 462–3
- coronary disease  
acute coronary syndrome, 167–74  
acute MI, 179–87
- delirium, 254
- eclampsia, 438–9
- elderly  
from abuse, 640  
in “hospital at home” scheme, 263–5  
screening for risk after discharge, 266
- heart failure (acute decompensated), 193–5
- intimate partner violence, reducing, 602–4
- multi-trauma/hemorrhagic shock, 374–7, 378–9
- overcrowding and its impact on, 67–8
- PE and  
and anticoagulants, 128, 129–30  
and diagnostic algorithms, 130
- pneumonia, 108
- salicylate poisoning, 300–1
- sepsis and septic shock, 238, 238–47  
interventions affecting, 239–47  
risk assignment, 238–9
- streptococcal pharyngitis, 526–8  
*see also* suicide
- moxifloxacin,  
pelvic inflammatory disease, 431, 432
- multiple trauma, hemorrhagic shock, 373–80
- mumps *see* measles/mumps/rubella immunization
- murmur, diastolic, thoracic aortic dissection, 159
- musculoskeletal surgeon or specialist, referral in knee injury to, 342
- mutagenic risk to fetus of CT, 388
- Mycoplasma pneumoniae*, 100, 105
- myocardium  
infarction, 158–60, 162, 166–70, 177–89  
aspirin prophylaxis, 162  
ECG in diagnosis of, 158–60  
non-ST elevation (NSTEMI), 153, 155, 157, 165–6, 166–70  
smoking continuance vs cessation and risk of, 611  
ST elevation (STEMI), 45, 155, 177–89  
syncope and detection of, 274–6  
ischemia *see* ischemia
- nadroparin, acute coronary syndrome, 171
- naloxone, 631–2
- naltrexone, post-concussive headache, 313
- NAPQI (N-acetyl-p-benzoquinoneimine), 296
- narcotic analgesics (opiates), 629–32  
knee injury, 340–1  
migraine, 495–6  
overdose/misuse, 629–32  
antidotes, 631–2  
ureteric calculi, 406–8
- nasal decongestants, rhinosinusitis, 540
- nasogastric aspiration, upper GI bleeding diagnosis, 449–50
- National (Australasian) Triage Scale (ATS), 59
- National Emergency X-ray Utilization Study *see* NEXUS
- National Institute for Health and Clinical Excellence (NICE) guidelines, mild head injury, 308–9
- National Institute of Alcohol Abuse and Alcoholism guidelines, 625, 626
- National Institute of Neurological Disorders and Stroke’s (NINDS) tPA trial, 471, 472
- National Registry of Myocardial Infarction, Second, 185
- natriuretic peptide, B-type *see* brain-derived natriuretic peptide
- nausea in pregnancy, 439–41
- nebulized drugs in asthma, 81–2, 82–3
- neck  
injury, 316–24  
stiffness, bacterial meningitis, 487
- needle(s)  
for lumbar puncture in bacterial meningitis, 489  
pediatric pain reduction with procedures involving, 620–1
- needle aspiration of peritonsillar abscess  
diagnostic, 529–30  
therapeutic, 530–1
- neglect, elderly, 636, 637–8
- Neisseria gonorrhoeae* infection (gonorrhoea), 414, 415, 420, 421, 430
- neomycin/bacitracin/polymyxin ointment, wounds, 567
- nesiritide, acute decompensated heart failure, 194–5
- neuraminidase (influenza virus)  
inhibitors, 138–9  
subtyping, 133
- neuroimaging  
bacterial meningitis, 488  
head injury (mild), 308–9, 310–12  
ischemic stroke, 468–70  
seizure (first), 505  
subarachnoid hemorrhage, 478  
transient ischemic attack, 462  
*see also specific modalities*
- neuroleptics (antipsychotics), agitation and/or delirium, 254–7, 515–17
- neurological disorders, 455–519  
alcohol-induced, 286–8  
CO poisoning, delayed (DNS; delayed neurological sequelae), 291, 293  
syncope due to, 277
- neurological outcome (and possible deficits)  
bacterial meningitis, 489–91  
subarachnoid hemorrhage, 481  
ventricular arrhythmias, 218–21
- neuropsychiatric problems, 251–9  
CO poisoning, 291
- New Orleans Criteria, 308, 309, 311
- NEXUS (National Emergency X-ray Utilization Study)  
brain trauma (mild), 309, 310  
neck trauma, 318–19, 319, 320, 321
- NICE guidelines, mild head injury, 308–9
- nicotine replacement therapy, 608, 609, 611
- nicotinic acetylcholine receptor antagonist, smoking cessation, 608
- nimodipine, subarachnoid hemorrhage, 480, 481
- nitrites, urinary tract infection, 414–15
- nitroglycerin  
acute decompensated heart failure, 192–3  
coronary disease, 160–1, 169
- NMDA receptors and alcohol, 286
- non-accidental injury *see* injury
- non-invasive ventilation (NIV)  
acute decompensated heart failure, 193–4  
COPD, 92, 93–4
- non-ST elevation myocardial infarction (NSTEMI), 153, 155, 157, 165–6, 166–70

## Index

- non-steroidal anti-inflammatory drugs (NSAIDs; cyclo-oxygenase/prostaglandin synthase inhibitors)  
 DVT, 117–18  
 GI side-effect prevention, 448–9  
 knee injury, 340–1  
 migraine, 498  
 post-concussive headache, 312  
 streptococcal pharyngitis, 528  
 ureteric calculi, 406–7, 408  
 norepinephrine, sepsis/septic shock, 244, 245  
 norfloxacin, conjunctivitis, 546  
 nurses, triage and reliability scores with, 61
- observation unit  
 acute decompensated heart failure, 196  
 occupation, time to return to *see* return  
 octylcyanoacrylate, 563  
 ocular emergencies, 543–9  
 ofloxacin  
 pelvic inflammatory disease, 431, 432  
 urinary tract infection, 417  
 older patients *see* elderly  
 omeprazole, 447, 450  
 ondansetron, nausea and vomiting in pregnancy, 440  
 online (web)-based CME, 38–9  
 open-ended questions in assessment of medication adherence, 52  
 open reduction, shoulder dislocation, 352  
 ophthalmological emergencies, 543–9  
 opiates *see* narcotic analgesics  
 oral contraceptive pill overdose, 282–6  
 orthopedic clinic, displaced distal radial fracture, 348  
 orthopedic surgeon or specialist, referral in knee injury to, 342  
 oseltamivir, 138  
 osteoporosis prevention, distal radial fracture, 348  
 Ottawa Ankle Rule, 327–8  
 Ottawa Knee Rule, 338–9  
 outcome  
 in clinical questions, 7, 8  
 patient health, CME effects, 39  
 scores, in triage, 61  
 outpatient follow-up *see* follow-up  
 outpatient management, pelvic inflammatory disease, 431–2  
 overcrowding, 65–75  
 defining, 66–7  
 interventions reducing, 68–72  
 triage and, 62–3, 69  
 overdose, drug, 282–6, 296–303, 629–32  
 Overview Quality Assessment Questionnaire (OQQAQ), 26–8  
 oxycodone, knee injury, 341  
 oxygen  
 administration  
 acute decompensated heart failure, 192–3  
 chest pain, 162  
 CO poisoning, 291, 291–3  
 COPD, 92  
 procedural sedation and analgesia, 556–7  
 saturation (SaO<sub>2</sub>) in asthma, 80–1
- pain  
 abdominal *see* abdominal pain  
 in acromioclavicular joint separation, reducing duration and intensity, 355–6  
 in ankle injury, reducing, 329–31  
 chest *see* chest pain  
 head *see* headache  
 of intramuscular vaccine injection in children, reduction, 620–1  
 in knee injury, reducing, 340–1  
 in neck injury, long-term, 321–2  
 pelvic, chronic, following pelvic inflammatory disease, 431–2  
 pharmacological management *see* analgesia  
 in rotator cuff injury, reducing duration, 354–5  
 ureteric, 404–11  
 in wrist injury, 345, 345–7, 347  
 paper-based triage, 63  
 paracetamol *see* acetaminophen  
 patient  
 care *see* care  
 death *see* mortality  
 education *see* education  
 flow, impact of interventions reducing overcrowding on, 68–9  
 getting evidence to, 18–19  
 health outcome, CME effects, 39  
 medication adherence, 49–57  
 satisfaction *see* satisfaction  
 suffering in migraine, decreasing, 498  
*see also* morbidity  
 payment incentives, smoking cessation, 610–11  
 PDSA (plan, do, study, act), 44–5  
 peak expiratory flow (PEF) in asthma, 79  
 pediatrics *see* children  
 pelvic inflammatory disease, 420–34  
 pelvic trauma, 378–9  
 penetrating chest trauma, 370  
 penicillins  
 meningitis, 489  
 pneumonia, 106  
 streptococcal pharyngitis, 528  
 pentobarbital, 507  
 peptic (gastroduodenal) ulcer, 447, 448, 449, 450, 452  
 percutaneous interventions (PCI; angioplasty; invasive/revascularization procedures), 183–4  
 in acute coronary syndrome, 173–4  
 in acute MI, 177–8, 183–4  
 primary, vs fibrinolytics, 183–5  
 reducing, 181–7  
 performance (physician), changes/improvements in CME producing, 39  
 quality improvements producing, 44–5  
 pericardial effusion and tamponade, 369–70  
 peripheral blood biomarkers, bacterial meningitis, 486–7  
 peritoneal cavity in blunt abdominal trauma  
 blood in, 582  
 diagnostic lavage, 579, 583  
 free fluid, 582–3  
 peritonsillar abscess, 523, 528–31  
 pertussis *see* tetanus immunization, with diphtheria/pertussis
- pethidine, ureteric calculi, 406–7  
 pharmacotherapy *see* drug therapy  
 pharyngitis, 523–33  
 phenazopyridine, urinary tract infection, 418  
 phenobarbital, 507  
 phenothiazines  
 agitation, 515–17  
 nausea and vomiting in pregnancy, 440  
 phenytoin, 507, 507–8  
 alcohol-related seizures, 277–8, 289, 290  
 eclampsia, 438–9  
 physical abuse  
 elderly  
 definition, 536  
 history and physical signs, 636–7  
 intimate partner, 590, 594, 600–5  
 physical activity/play/sport  
 causing knee injury, 335  
 causing shoulder injury, 350  
 return to *see* return  
 physical examination *see* examination  
 physician  
 audit and feedback, 45  
 clinical judgment and experience  
 ankle injury, 327–8, 329  
 syncope, 278  
 compliance/adherence, aids/initiatives, 19, 20  
 decisions *see* decision  
 learning, CME producing changes in, 39  
 medication adherence assessed by, 51–2  
 performance *see* performance  
 primary care *see* primary care  
 smoking cessation advice, 607–8  
 physiotherapy  
 ankle injury, 331  
 humeral fracture (proximal), early vs delayed, 358–60  
 knee injury, 341–2  
 rotator cuff tears, 354–5  
 piracetam, post-concussive headache, 312, 313  
 pivot shift test, 337  
 placebo  
 acute coronary syndrome, 167, 169–70  
 atrial fibrillation, 203–4, 205–8  
 DVT, 117–18  
 GI bleeding (upper), 450  
 migraine, 495–8, 499  
 pyelonephritis in pregnancy, 441–2  
 rhinosinusitis, 537–8  
 seizures, 505–7  
 urinary tract infection, 416  
 ventricular arrhythmia, 219–21  
 plain films *see* X-ray  
 plaster lower-leg cast, ankle, 331  
 platelet inhibitors *see* antiplatelet agents  
 pneumococcus *see* *Streptococcus pneumoniae*  
 pneumonia  
 community-acquired, 100–12  
 elderly, pneumococcal vaccine and risk of, 618–19  
 severity index (PSI), 103  
 pneumothorax, traumatic, 367–9  
 poisoning *see* toxicology  
 poliovirus immunization schedules, 616

- polymyxin/bacitracin ointment *see* bacitracin  
 polypharmacy screening in elderly prior to discharge and impact on adverse effects, 265  
 pooled statistics, heterogeneity among, 9–10  
 population (in clinical questions), 7, 8  
 positive pressure ventilation (non-invasive), acute decompensated heart failure, 193–4  
 post-concussive symptoms, 312–14  
 posterior drawer test, knee injury, 336  
 post-phlebotic (post-thrombotic) syndrome, 119  
 prediction rules *see* clinical prediction rules  
 prednisone  
   asthma, 82  
   COPD, 92  
 pregnancy, 391–403, 435–44  
   CT in appendicitis, hazards and alternatives, 388–9  
   ectopic, 391–403  
   loss *see* abortion  
   missed, 435  
   test, 435, 437  
 prehospital fibrinolysis, acute MI, 180–1  
 prescription instructions and medication adherence, 53  
 preterm delivery and bacteriuria, 441–2  
 primary care (incl. physician/general practitioner)  
   displaced distal radial fracture, 348  
   ED care vs, 70  
   gatekeeping, 70  
   referral back with medication non-adherence, 53–5  
 probenecid, pelvic inflammatory disease, 430  
 procainamide  
   atrial fibrillation, 204  
   ventricular tachycardia, 221–2  
 prochlorperazine  
   migraine, 497, 498  
   in pregnancy, 440  
 prognostic screening tool, elderly prior to discharge, 266  
 prolactin levels, seizures vs pseudoseizures or syncope, 508–9  
 promethazine, nausea and vomiting in pregnancy, 440  
 pronator fat stripes, 345  
 propafenone, atrial fibrillation, 204  
 propofol  
   procedural sedation and analgesia, 558, 559  
   status epilepticus, 507  
 propranolol, post-concussive headache, 313  
 prostaglandin synthase inhibitors *see* non-steroidal anti-inflammatory drugs  
 protein C, recombinant activated, in sepsis/septic shock, 239, 245  
 proton pump inhibitors, upper GI bleeding, 448, 450  
   prophylactic use, 449  
*Pseudomonas aeruginosa* in COPD, 95  
 pseudo-seizures, 503, 508–9  
 psychiatric assessment, agitation, 517  
 psychiatric illness *see* mental illness  
 psychological abuse of elderly, 636  
 psychometric principles and properties, 23–4  
 psychosis and agitation, 515  
 psychotherapy, supportive, smoking cessation, 610  
 public access defibrillation, 227  
 public health, 587–643  
 pulmonary embolism, 114, 123–32  
 pulmonary hypertension, chronic thromboembolic (CTPH), 123  
 pulmonary non-vascular tissue *see* entries under lung  
 pulse deficits, thoracic aortic dissection, 158, 159  
 pulseless ventricular tachycardias, defibrillation, 227–8  
 PURSUIT (Platelet Glycoprotein IIb/IIIa in Unstable Angina: Receptor Suppression Using Integrilin Therapy) model, 166, 167  
 pyelography, intravenous (IVP), calculi, 405, 405–6  
 pyelonephritis in pregnancy, 441–2  
 pyramidal organization of research evidence, 16–18  
 pyrexia (fever), bacterial meningitis, 487  
 pyridium, urinary tract infection, 418  
 pyridoxine, nausea and vomiting in pregnancy, 440  
  
 QT interval, long, 214, 223  
 qualitative rating tools, 24  
 quality  
   assessment  
     of published articles (in general), 23, 25–6  
     randomised controlled trials, 25–6  
   of care, improvement, 43–8  
     continuing, 19  
   of life, 9  
     acromioclavicular joint injury and, postoperative, 355–6  
     community-acquired pneumonia, 108–10  
     neck (soft tissue) injury-related pain, 322  
     pneumothorax (traumatic) treatment and, 367–8  
     rotator cuff tear treatment and, 354–5  
     scaphoid fracture imaging and, 345  
   quantitative rating tools, 24–5  
   questions  
     arising from case (in chapter organization), 6  
     development, 7–8  
     improving efficiency, 8  
     open vs closed-ended, in assessment of medication adherence, 52  
   quinidine, atrial fibrillation, 203  
   quinsy (peritonsillar abscess), 523, 528–31  
  
 rabies immunization, dog bite, 618  
 radiography, chest *see* chest X-ray  
 radiology *see* imaging and specific modalities  
 radionuclide scans *see* scintigraphy  
 radius, distal, fracture  
   clinical scenario, 533, 559  
   displaced, 347–8  
 randomised controlled trials (RCTs), 8, 14  
   moderate evidence from, 4  
   quality assessment, 25–6  
   rating tools, 23, 24–5  
 Rational Clinical Examination, streptococcal pharyngitis, 525  
 recommendation generation and implementation in root cause analysis, 46  
 rectum, foreign body, 553, 559  
 red cells *see* erythrocytes  
 red eye, 543–9  
 reduction  
   distal radial fracture, 347, 347–8  
   shoulder dislocation, 351–2, 356–7  
 referral  
   back to primary care physician with medication non-adherence, 53–5  
   to orthopedic surgeon or specialist, knee injury, 342  
 renal colic *see* ureteric colic  
 renal function/problems *see* kidney function;  
   pyelography; pyelonephritis  
 repair, wound, 561–70  
 research  
   closing gap between practice and, 19–20  
   evidence *see* evidence  
   respiratory medicine, 77–149  
   respiratory syndrome, severe acute (SARS), 81, 134  
 restraint  
   physical, in agitation, 513–14, 515–17  
 resuscitation *see* cardiopulmonary resuscitation; fluids  
 reteplase, acute MI, 186  
 return  
   to activities (including daily activities/play/sport/work)  
     ankle injury and time to, 329–30  
     knee injury and time to, 341–2  
     rotator cuff tears and time to, 354–5  
     second impact syndrome risk, 312–13  
     streptococcal pharyngitis, 528  
   to ED  
     asthma management strategies decreasing,  
     COPD management strategies decreasing,  
     medication non-adherence as cause of, 52–3  
     migraine management strategies decreasing,  
     498  
   revascularization, percutaneous *see* percutaneous  
     interventions  
   reviparin in PE, 129  
   rheumatic fever, 527  
   rhinosinusitis, 524–42  
   rimantadine, 138  
   risk stratification and risk factor profiling  
     alcohol problems, 627–9  
     chest pain, 156  
     elder abuse, 638–9  
     intimate partner violence, 602  
     syncope, 277  
     unstable angina/NSTEMI, 166–7  
   root cause analysis, 46–7  
   rotator cuff, 350  
     injuries, 352, 353–5  
   rotavirus, pediatric immunization schedule, 616  
   rubella *see* measles/mumps/rubella immunization  
  
 safety belts, 590, 594–5  
 salbutamol  
   asthma, 79, 82–3  
   COPD, 92  
 salicylate poisoning, 296–8, 300–2  
 saline  
   in hemorrhagic shock, hypertonic, 375–6  
   in sepsis/septic shock, vs colloids, 243–4  
   wound irrigation with, vs tap water, 563–4  
 salpingitis, 420, 422  
 salpingostomy, laparoscopic, ectopic pregnancy, 400–1  
 San Francisco Syncope Rule (SFSR), 276–7

## Index

- satisfaction  
 care-giver, elderly in "hospital at home" scheme, 263–5  
 patient  
 ankle injury X-ray and, 328–9  
 elderly in "hospital at home" scheme, 263–5
- scaphoid fracture, 345–7
- scapular manipulation, 357, 359
- scintigraphy (radionuclide scans)  
 appendicitis in pregnancy, 389  
 pelvic inflammatory disease, 428  
 scaphoid fracture, 345–6
- screening  
 abuse of person  
 elderly, 639  
 intimate partner, 601  
 alcohol abuse, 591, 625  
 dementia, 262–3  
 depression, 517–18  
 mortality risk in elderly following discharge, 266  
 polypharmacy, in elderly prior to discharge and impact on adverse effects, 265  
 toxicology, 630–1
- seat (safety) belts, 590, 594–5
- second impact syndrome, 312–13
- second intention healing, 562–3
- sedation and analgesia, procedural, 553–60
- seizures/fits, 503–11  
 alcohol-related, 286–8  
 drug use, *see also* anticonvulsants  
 eclamptic, 438–9  
 epileptic vs non-epileptic, 503  
 first, 504–5  
 partial vs generalized, 503  
 refractory, 507  
 simple vs complex, 503
- self-injectable epinephrine, 148
- self-neglect, elderly, 636
- self-reporting in assessing medication adherence, 51
- semi-rigid ankle support, 330
- sepsis (systemic infection), 237–50  
 definition, 237  
 severe, 237–50  
 definition, 238
- septic shock, 237–50  
 definition, 238
- serotonin 1B/1D antagonist, migraine, 496
- severe acute respiratory syndrome (SARS), 81, 134
- sexual abuse of elderly, 636
- sexual intercourse, unintentional, 589, 596
- sexually-transmitted disease vs simple urinary tract infection, 414–15
- Sgarbossa criteria, 160
- shingles (zoster) immunization schedules, 616
- shock  
 electrical, in ventricular arrhythmia *see* defibrillator shock  
 hemorrhagic, 373–80  
 septic, 237–50
- shoulder injury, 350–62
- silver sulfadiazine, wounds, 567
- simple agreement (SA), triage systems, 60
- single photon emission computed tomography (SPECT), mild head injury, 312
- sinus rhythm, cardioversion *see* cardioversion
- sinusitis/rhinosinusitis, 524–42
- smallpox immunization, 619
- smoking, 606–13  
 cessation, 606–13  
 in COPD, 96–7  
 epidemiology, 606–7  
 heart disease risk, 611
- sodium bicarbonate *see* bicarbonate
- sodium valproate, 507  
 and valproic acid (divalproex sodium), post-concussive headache, 313
- soft tissue abscess, 571–7
- soft tissue injury  
 neck, 321–2  
 shoulder, 352
- sonography *see* ultrasonography
- sore throat, 523–33
- sotalol  
 atrial fibrillation, 204  
 ventricular tachycardia, 221–2
- Soterion Rapid Triage System, 63
- Spaso reduction method, 357
- spectrophotometry, xanthochromia in CSF, 479–80
- spinal injury, cervical, 316–24
- spiral (helical) CT, acute appendicitis, 386–9
- sport *see* physical activity
- sprain, ankle, 325, 331–3  
 repeat, with chronic instability, prevention, 331–3
- ST elevation MI (STEMI), 45, 155, 177–89
- staffing levels,  
 overcrowding due to, 69
- Staphylococcus aureus*, community-acquired methicillin-resistant, 571–2, 573–4
- status epilepticus, 504  
 anticonvulsants, 506, 507
- sterile vs non-sterile gloves, wound management, 564–5
- steroids *see* corticosteroids
- stomach (gastric)  
 acid *see* acid  
 emptying, drug overdose, 282, 283, 284  
 endoscopy in upper GI bleeding, immediate vs delayed, 448  
 lavage in upper GI bleeding diagnosis, 449–50  
 ulcer, 449
- stone, ureteric, 404–11
- stool occult blood testing in anemia, 447–8
- streptococcal sore throat, 523, 524–8  
 group A  $\beta$ -hemolytic streptococcus (GABHS), 523, 526, 527, 528, 529
- Streptococcus pneumoniae* (pneumococcus), 100, 106, 110  
 COPD, 95  
 immunization schedules, 616  
 children, 615  
 elderly, 616, 618–19
- streptokinase (SK) in acute MI, 181, 183, 186
- stroke (cerebrovascular accident), 466–73  
 risk following TIA, 457–8, 458–60, 462  
 risk in atrial fibrillation, 200  
 prevention, 205–10
- studies (in the 5S levels of organization of research evidence), 16
- subarachnoid hemorrhage, 475–84
- substance misuse, 624–34
- suicide risk  
 adolescents (mentally ill) and firearms, 595  
 agitated patient, 514–15, 517
- sulbactam, pelvic inflammatory disease, 431
- sulfamethoxazole *see* trimethoprim/sulfamethoxazole
- sumatriptan  
 migraine, 496, 497  
 post-concussive headache, 313
- summaries (in 5S levels of organization of research evidence), 18
- support (external), ankle, 330, 331–2
- supraventricular arrhythmias (SVT), 215, 215–18
- surgery  
 ankle instability (chronic), 332–3  
 appendicitis, prompt vs delayed, 383–4  
 coronary artery bypass, clopidogrel and, 169  
 ectopic pregnancy, 400–1  
 intracranial abnormalities (in mild brain trauma) on CT requiring, 308–9  
 knee injury and referral to orthopedic surgeon, 342  
 minor procedures, 551–85
- shoulder  
 acromioclavicular joint separation, 355–6  
 rotator cuff tears, 354–5
- torso trauma, 583  
*see also* incision and drainage
- swelling, reducing  
 knee injury, 340–1  
 soft tissue abscess, 572–3
- syncope, 271–81  
 designated syncope unit, 279–80  
 seizures and, differentiating, 508–9
- synopses (in 5S levels of organization of research evidence), 16–17
- syntheses (in 5S levels of organization of research evidence), 16
- system(s)  
 in 5S levels of organization of research evidence, 18  
 of work, overcrowding related to, 69
- system-wide interventions, 70–1
- systematic reviews  
 Cochrane Collaboration and, 5, 6, 14, 17  
 in 5S levels of organization of research evidence, 16  
 identifying high-quality evidence in, 26–8  
 levels of evidence and, 5
- systemic inflammatory response syndrome, 237
- tachycardias  
 supraventricular (SVT), 215, 215–18  
 ventricular (VT), 214–15, 215–23  
 defibrillator shock in, 221, 227–8  
 defibrillator shock-resistant, 218–21  
 polymorphic, 214, 222–3  
 pulseless, 227–8  
 stable, 221–2
- tactile stimulus, response in sedation and analgesia, 554

## Index

- tamponade, pericardial, 369–70  
tamulosin, ureteric calculi, 409  
tap water, wound irrigation, 563–4  
taping, ankle injury, 330–1  
technetium-99 tagged WBCs/leucocytes  
  appendicitis in pregnancy, 389  
  pelvic inflammatory disease, 428  
telephone quitlines (smoking), 610  
tenecteplase  
  acute MI, 186  
  cardiac arrest, 229  
tetanus immunization, 617–18  
  contaminated wound, 617–18  
  with diphtheria/pertussis, 616  
  pediatric, 615  
tetracaine, topical, wounds, 566  
tetracycline, pelvic inflammatory disease,  
  431  
thiopental, 507  
thoracic aorta *see* aorta  
thoracic trauma, 363–72  
throat, sore, 523–33  
thrombocytopenia, heparin-induced, 170  
thromboembolic disease, 113–32  
  in atrial fibrillation  
  prophylaxis, 201–2, 205–8  
  risk, 200, 201–2  
thrombolytics/fibrinolytics  
  in acute ischemic stroke, 471–2  
  in acute MI, 160, 177, 183–6  
  adjunctive heparin, 185–7  
  prehospital administration, 180–1  
  primary angioplasty vs, 183–4  
  in cardiac arrest, 229  
  in PE, 129–30  
thrombophilias, inherited, 115, 116  
thrombosis, deep vein (DVT), 113–22,  
  123  
TIMI risk score, 156–7, 167  
tinzaparin  
  acute coronary syndrome, 171  
  PE, 129  
tirofiban, ischemic stroke, 470  
tissue adhesives, 563  
tissue-plasminogen activator (t-PA)  
  acute ischemic stroke, 471–2  
  acute MI, 183  
  cardiac arrest, 229  
tobacco *see* smoking  
tobramycin, conjunctivitis, 546  
torsades de pointes  
  in atrial fibrillation,  
  in cardiac arrest, 214, 222–3  
toxicology, 282–303  
  screen, 630–1  
tracheal intubation *see* intubation  
tranexamic acid, hemorrhagic shock, 377  
transfusion *see* blood transfusion  
transient ischemic attack (TIA), 457–65  
transvaginal ultrasound (TVUS)  
  ectopic vs intrauterine pregnancy, 396  
  pelvic inflammatory disease, 424–6  
trauma *see* fracture; injury  
triage scales  
  ATS, 59  
  CTAS, 59, 60, 62, 63  
  MTS, 59, 62  
triage, 58–64  
  delirium, 251  
  overcrowding and, 62–3, 69  
  reliability of systems, 60–1  
  systems and their differences, 59–60  
  validity of systems, 60–1, 61–2  
Triage Risk Screening Tool, 266, 268  
trimethoprim/sulfamethoxazole (co-trimoxazole),  
  539  
  urinary tract infection, 416  
troponin  
  cardiac contusion, 364, 365–6  
ulcer, peptic/gastroduodenal, 447, 448, 449, 450,  
  452  
ultrasonography, 578–85  
  appendicitis, 387–8, 389  
  carotid disease, 463  
  DVT, 117  
  PE, 126–8  
  pelvic inflammatory disease, 424–6  
  pericardial effusion and tamponade,  
  369–70  
  peritonsillar abscess, 529–30  
  pregnancy, intrauterine, 436–7  
  ectopic vs, 395–400  
  rhinosinusitis, 536–7  
  rotator cuff tears, 354  
  soft tissue abscess, 572  
  trauma patients  
  abdominal trauma, 579, 582–3  
  multi-trauma patients, 377–8  
  *see also* echocardiography  
ureteric colic, 404–11  
urine test  
  for infection, dipstick, 414–15  
  for pregnancy, home kit, 437  
urogenital problems, 391–444  
urography, intravenous, ureteric calculi (IVU), 405,  
  405–6  
US Preventive Services Task Force, abuse screening,  
  639  
vaccination, 614–23  
  in COPD, 96–7  
  influenza *see* influenza  
vaginal bleeding  
  ectopic pregnancy,  
  first trimester, 392–3, 436, 437–8  
  threatened abortion,  
valproic acid and sodium valproate (divalproex sodium),  
  post-concussive headache, 313  
varenicline, 608, 609  
varicella immunization schedules, 616  
vasopressin  
  analog (DDAVP), hemorrhagic shock, 377  
  cardiac arrest, 229  
  sepsis/septic shock, 244–5  
vasopressors  
  cardiac arrest, 229  
  sepsis/septic shock, 241, 244–5  
vasospasm prevention in subarachnoid hemorrhage,  
  480  
venography, CT, in suspected PE, 126–8  
venous access and monitoring, central, *see entries under*  
  central venous  
venous thromboembolic disease *see* thromboembolic  
  disease  
venous thrombosis, deep (DVT), 113–22, 123  
ventilation  
  in cardiac arrest, 228  
  chest compression  $\pm$ , 226  
  mechanical, in sepsis/septic shock, 247  
  non-invasive *see* non-invasive ventilation  
ventilation-perfusion (V/Q) scan in PE, 124, 126  
ventricular arrhythmias, 214–15, 216–23  
verapamil  
  atrial fibrillation, 204  
  post-concussive headache, 313  
  supraventricular tachycardia, 217–18  
verbal command, response in sedation and analgesia,  
  554  
verbal instructions and medication adherence, 53  
Verbal Learning Test, Hopkins', 262  
violence, intimate partner, 590, 594, 600–5  
viral conjunctivitis, 543  
Virchow's triad, 113, 115  
virological diagnosis of influenza, 135–8  
vital signs  
  blunt abdominal trauma, 582–3  
  normal, hemorrhagic shock resuscitation to,  
  374  
vitamin B<sub>6</sub> (pyridoxine), nausea and vomiting in  
  pregnancy, 440  
vomiting in pregnancy, 439–41  
  *see also* anti-emetics  
warfarin  
  atrial fibrillation, 205–8  
  transient ischemic attack, 461  
water  
  intake, increasing  
  ureteric calculi, 409–10  
  urinary tract infection, 416  
tap, wound irrigation, 563–4  
web-based,  
  CME, 38–9  
  patient education,  
Wells' prediction rule, deep venous thrombosis,  
  115  
whiplash-associated disorder, 321–2  
white blood cells (leukocytes; WBCs)  
  count in appendicitis, 385–6  
  esterase test for urinary tract infection, 414–15  
  radionuclide scans with  
  appendicitis in pregnancy, 389  
  pelvic inflammatory disease, 428  
wide complex tachycardia, 215–16  
women/females  
  intimate partner violence *see* intimate partner violence  
  urinary tract infection, 414–16  
work, time to return to *see* return  
wounds  
  contaminated, tetanus risk, 617–18  
  repair, 561–70

## Index

wrist injury, 344–9  
written action plan  
asthma,  
medication adherence, 53

X-ray (plain films/radiographs)  
ankle injury, 325, 327, 328–9  
chest *see* chest X-ray

knee injury, 338–9  
neck injury, 219–20  
peritonsillar abscess,  
529–30  
rhinosinusitis, 536–7  
wrist injury, 345  
*see also* NEXUS  
xanthochromia in CSF, 479–80

xylometazoline, rhinosinusitis,  
540

Zahiri reduction method, 357  
zanamivir, 138  
zatosetron, migraine, 497  
ziprasidone, agitation, 257  
zoster immunization schedules, 616