

Index

- 3TC see lamivudine
- abdominal
- decompression 140–1
 - incisions 343–4, 348–9
 - nerves block 355
 - stretch marks 35–6
- abnormal presentation see presentation
- abortions see termination
- absorbable synthetic
- sutures 319–20
- acquired immunodeficiency syndrome (AIDS) 102
- activated charcoal 120
- active management 263, 268–9, 289
- acupressure 280–1
- acupuncture
- induction 225, 244–5
 - pain during labour 280–1
 - symptoms during pregnancy 34–5, 37–8
- acyclovir 203–4
- advanced extrauterine pregnancies 357
- advice during pregnancy 13–22
- see also counselling
- aerobic exercise
- advice during pregnancy 21–2
 - diabetes mellitus 90–1
 - hypertension 59–60
 - weight retention 366–7
- AFV see amniotic fluid volume
- age assessment 128–9
- AIDS see acquired immunodeficiency syndrome
- albuterol 179
- alcohol 17–20
- alloimmune thrombocytopenia 95
- alloimmunisation 147–9
- aloe vera 353–4
- ambulatory blood pressure
- monitoring 69–70
- amino acid supplements 144
- amniocentesis 129–30
- amnioinfusion 155, 310, 313–15
- amnionitis 153
- amniotic fluid volume (AFV) 138, 146
- amniotomy
- induction 229–31, 238, 240–1
 - slow progress during labour 287–9
- amoxicillin 200, 201–2
- ampicillin 211, 338–9
- anaemia 85–9
- malaria 114–18
 - maternal 29–30
 - prevention 86–7
 - treatment 87–9
- anaesthesia
- caesarean section 328–33, 355
 - general 328–9
 - labour 261, 263–4, 266–7
 - local 355
 - postnatal care 371–2
 - regional 328–33
 - topical 371–2

A Cochrane Pocketbook: Pregnancy and Childbirth G.J. Hofmeyr et al.
Copyright © 2008, Z. Alfiervic, C. A. Crowther, L. Duley, A. M. Gulmezoglu, G. ML. Gyte, E. D. Hodnett, G. J. Hofmeyr, J. P. Neilson

- anal sphincter injuries 321–2
- antacids 264
- antenatal
 - clinics 157–8, 189–90
 - corticosteroids 167–8
 - depression 123–5
 - education 254
 - lower genital tract
 - infections 165–6
 - psychosocial screening 378–9
 - records 3
- antenatal care 7–54
 - behaviour/advice during pregnancy 13–22
 - general 7–13
 - medicines during pregnancy 40–1
 - miscarriage 41–54
 - nutrition during pregnancy 22–33
 - pre-pregnancy evaluation 7
 - symptoms during pregnancy 33–40
 - weight gain 23–5, 33
- antepartum fetal assessments 136–7
- antepartum haemorrhage 197–8
- anti-D immunoglobulin 147–9
- anti-emetics 34–5
- antibiotics
 - breastfeeding 392–3
 - caesarean section 337–9
 - genital tract infections 181–2, 200–7
 - medical problems during pregnancy 68–9, 99, 118
 - operative vaginal births 326
 - perineal trauma 322
 - postnatal care 377–8
 - prelabour rupture of the membranes 153–4, 155–6, 181, 199–200
 - preterm births 157, 160–1, 181–2, 199
 - retained placenta 361
- anticoagulants 42–3, 46–7
- anticonvulsants 77, 80–5
- antidepressants 125, 381–3
- antiemetics 331
- antiepileptics 77, 84–5, 100–1
- antigens, dietary 367–8
- antihistamines 34–5
- antihypertensive agents 70–1, 72–6
- antimalarials 115–16
- antioxidants 64–5
- antiphospholipid (APL) antibody/syndrome 42–3, 46–7
- antiplatelet agents 65–6
- antipsychotics 123–4, 384–5
- antiretroviral (ARV) drugs 101–11
- antiseptics 333–4, 336–7
- antithrombotic therapy 69
- anxiety 378–9
- Apgar scores
 - epidural analgesia 283
 - nutrition 27, 28
 - screening 135
 - shoulder dystocia 293
- APL see antiphospholipid
- aromatherapy 280–1
- artesunate 116
- ARV see antiretroviral
- Ascaris lumbricoides* 217
- aspiration pneumonitis 331
- aspirin 35, 42–3, 46–7, 65–6
- assisted vaginal delivery see operative vaginal births
- asymptomatic bacteriuria 207–9
- atopic eczema 367–8
- atosiban 180–1, 185
- audioanalgesia 280–1, 333
- audit 3–4
- auscultation, intermittent 308
- autoimmune thrombocytopenia 94
- azithromycin 201–2
- Baby Friendly Hospitals Initiative 390–2
- back pain 37–8
- bacterial vaginosis 118, 160–1, 167, 199
- bed rest

- hypertension 58–9, 72–3
- impaired fetal growth 141–2
- miscarriage 43–4
- multiple pregnancies 190–2
- preterm births 158–9, 177–8
- behaviour/advice during pregnancy 13–22
- Bendectin 34
- benign ovarian tumours 122–3
- beta-agonists 50
- beta-blockers 71, 73–4
- betamethasone 79–80, 170
- betamimetics
 - compromised fetal condition 142, 310–12
 - multiple pregnancies 194
 - preterm births 164–5, 174–5, 180–1, 184–5
- beta-stimulants 298–9
- biochemical tests of placental function 134–5
- biofeedback 282
- biophysical profiles (BPP) 135–6
- bladder catheterization 347
- blood group incompatibilities 147–51
- blood transfusions 88–9
- BPP *see* biophysical profiles
- bradycardia 73
- breast engorgement 395, 397
- breast stimulation 225, 245–6
- breastfeeding
 - antenatal care 40
 - antenatal preparation 386–8
 - augmenting medications 391–2
 - breast engorgement 394–5, 397
 - breast examinations 388
 - continuity of care 3
 - duration 387–9, 391, 395–7
 - education 387–9
 - expression 391–2
 - human immunodeficiency virus 104–6, 112–13
 - hypertension 75–6
 - initiation rates 387
 - mastitis 392–3
 - nipple pain 389–90
 - nutrition 393–4
 - postnatal care 363–5, 367–8, 374, 377, 386–98
 - support/interventions 388–94
 - suppression of lactation 397–8
 - workplace interventions 390–1
- breech births
 - caesarean section 300–2
 - expedited vaginal breech delivery 302
 - external cephalic version 294–9
 - fetopelvic disproportion 291–2
 - see also* presentation
- bromocriptine 397
- bupivacaine 329
- caesarean section 327–57
 - anaesthesia 328–33, 355
 - antenatal care 14–16
 - compromised fetal condition 304, 306
 - human immunodeficiency virus 106–9
 - hypertension 77
 - induction 223, 234–5
 - infections 204, 213, 333–41, 347–9
 - morbidly adherent placenta 361
 - multiple pregnancies 195–6
 - non-clinical interventions 16
 - non-medical reasons at term 262–3
 - operative vaginal births 326–7
 - perioperative measures 333–41
 - post-operative care 352–4
 - pregnancies following 354–7
 - presentation 296–7, 300–2
 - preterm births 186–7
 - repeat 223, 354–5
 - screening 135–6
 - slow progress during labour 289, 290–1, 292
 - surgical techniques 341–51, 399
- caffeine 17, 41

- calcium 36–7, 62–3
- calcium channel blockers
 - compromised fetal condition 143
 - hypertension 80–1
 - preterm births 175–6, 181, 183–4
- candidiasis 205
- cannulae 130–1
- carbamazepine 100
- carbetocin 273–4
- carbohydrate supplements 144
- cardiac disease 98–9
- cardiotocography (CTG) 136–8, 139, 303–4, 306, 307–9
- caregivers, labour 251–3, 255–6
- carnite 144
- castor oil 243–4
- catgut sutures 319–20
- caulophyllum 219–20, 247–8
- cefixime 200
- ceftriaxone 200
- cefuroxime 210
- cephalhaematoma 324
- cephalic version 294–9
- cephalosporins 338–9, 378
- cephradine 210
- cerebral palsy 172
- cervical
 - assessment 9–10
 - cerclage 157, 159
 - ripening 224–9, 234, 240–4, 245–7, 248–9
- CHAMOCA regimen 53–4
- chemotherapy 53–4
- childbirth *see* labour
- Chinese herbal medicines 74–5, 297–8
- chlamydia trachomatis 201–2
- chlorhexidine 205–7, 336–7
- chloroquine 115
- chlorpheniramine 35
- cholestasis 119–20
- chorioamnionitis 118, 154, 207
- chorionic villus sampling (CVS) 129–31
- chromosomal miscarriage 41, 43
- chronic hypertension 55, 57
- clindamycin 157, 201–2, 211, 377–8
- clipping (hair) 262, 334–6
- Clostridium tetani* 216–17
- closure of the peritoneum 345–6, 399
- clotrimazole 205
- CM *see* contingency management
- co-amoxiclav 153–4
- coagulation defects 95
- combination chemotherapy 53–4
- combined spinal-epidural (CSE)
 - analgesia 284–5, 286–7
- complementary therapies
 - acupressure 280–1
 - audioanalgesia 280–1, 333
 - caesarean section 331
 - Chinese herbal medicines 74–5, 297–8
 - homoeopathic remedies 219–20, 225, 247–8
 - hypnotherapy 280–1
 - pain during labour 280–1
 - presentation 297–8
 - see also* acupuncture
- complications in pregnancy 153–217
 - antepartum haemorrhage 197–8
 - prelabour rupture of the membranes 153–6, 160, 181, 199–200, 219–20
 - prolonged pregnancy 187–8
 - see also* infections; multiple pregnancies; preterm births
- compression stockings 40
- compromised fetal
 - condition 302–17
 - conservative management 140–6, 315
 - diagnosis 302–9
 - prevention 309–11
 - treatment 311–17
- congenital syphilis 212
- congenital toxoplasmosis 214–16
- conservative management 140–6, 315
- constipation 38–9

- context of care 1–5
- contingency management (CM) 18–19
- continuity of care 1–3
- continuous
 - cardiotocography 303–4, 306, 307–9
 - subcutaneous insulin injection 93–4
 - sutures 320–1
- contraception 372–6
- contractions
 - ineffective 287–90
 - see also* tocolytic therapy
- convulsions *see* eclampsia
- cooling treatments 370–1
- cord prolapse 317
- cord traction 274–5
- corticosteroids 79–80, 150
 - caesarean section 356–7
 - induction 241
 - preterm births 167–71, 399
- counselling
 - postnatal care 376, 379–80, 383, 393
 - prenatal diagnosis 132
- COX *see* cyclo-oxygenase
- critical incident audit/feedback 3–4
- cross-infection 339–40
- CSE *see* combined spinal-epidural
- CSII *see* continuous subcutaneous insulin injection
- CTG *see* cardiotocography
- curettage 52–3
- CVS *see* chorionic villus sampling
- cyclo-oxygenase (COX)
 - inhibitors 176–8
- cystitis 207
- cytomegalovirus 114

- d4T *see* stavudine
- ddl *see* didanosine
- Debendox 34
- deep vein thrombosis 98
- delayed delivery 151
- depilatory creams 334–6

- depression
 - antenatal 123–5
 - postnatal 378–83
- dexamethasone 79–80, 150, 170
- diabetes mellitus 89–94, 221–2
- diarrhoea 113
- diazepam 34, 77, 82–3
- diazoxide 80–1
- didanosine (ddl) 106
- diet *see* nutrition
- dietary antigens 367–8
- diethylstilbestrol 40–1, 45, 399
- digital cervical assessment 9–10
- dilatation and curettage 52–3
- dinoprostone 51–2, 232–3
- disadvantaged mothers 10–12
- disinfectants 109–11
- disposable face masks 340–1
- dopamine 85
- dopamine antagonists 264
- Doppler ultrasound 139–40
- double gloving 339–40
- dressings 353–4
- drug misuse 17–21
- dyspareunia 369–70

- early fetal death 51–2
- early skin-to-skin contact (SSC) 364–5
- ECG *see* electrocardiograms
- eclampsia
 - antenatal care 28
 - definition 55, 57
 - prevention 62–3, 69
 - severe 76–85
 - treatment 76–85
- ECV *see* external cephalic version
- eczema, atopic 367–8
- educational interventions 19–20
 - antenatal care 14, 254
 - breastfeeding 387–9
 - postnatal care 363–4, 372–4, 387–9
- EFM *see* electronic fetal monitoring
- electrocardiograms (ECG) 304–5

- electronic fetal monitoring (EFM)
 - 303–5, 306, 307–9
- EMA-CO/EMA-EP regimen 53
- encephalopathy, neonatal 304–5, 306, 314
- end-to-end repair 321–2
- endometritis
 - caesarean section 337–8
 - postpartum 199–200, 207, 377–8
- endotracheal intubation 316–17
- enemas 256–7
- energy intake 23–5
- ephedrine 332
- epidural anaesthesia 266–7, 330–1
- epidural analgesia
 - combined spinal-epidural
 - analgesia 284–5, 286–7
 - discontinuation 285–6
 - external cephalic version 298–9
 - pain during labour 282–7
 - preloading for regional
 - analgesia 286–7
- epilepsy 99–101
- episiotomy 264–8, 317–19, 399
- ergometrine 268–9, 270–2, 276
- erythromycin 153–4, 201–2
- erythropoietin 88–9
- estriol estimations 134–5
- exercise
 - advice during pregnancy 21–2
 - diabetes mellitus 90–1
 - hypertension 59–60
 - weight retention 366–7
- expectant management
 - death of the unborn baby 151
 - hypertension 77–8
 - induction 219–20, 228
 - labour 268–9
 - miscarriage 50–1
 - multiple pregnancies 195
 - preterm births 186–7
- expedited vaginal breech
 - delivery 302
- external cephalic version (ECV)
 - 294–9
- extra-abdominal repair 346–7
- extra-amniotic infusion 235, 236–7
- extrauterine pregnancies 357
- face masks 340–1
- family planning 372–6
- feedback 3–4
- fetal
 - alcohol syndrome 20
 - death 51–2, 91, 151, 220–1
 - distress 311–12, 315–16
 - electrocardiograms 304–5
 - fibronectin 161–2
 - growth retardation 26–7
 - lung maturation 169–70
 - macrosomia 221–3
 - manipulation 137–8
 - movement counting 133–4
 - pulse oximetry 305–6
 - scalp lactate sampling 308–9
 - surgery 132
 - vibroacoustic stimulation 136–7
 - wellbeing 140–6, 302–17
- feto-placental function 134–5
- fetomaternal alloimmune
 - thrombocytopenia 149–50
- fetopelvic disproportion 290–2
- fibre supplements 38–9
- fibronectin 161–2
- first stage labour 257–8
- fluid supplementation 393
- fluoxetine 383
- flushing 82
- folate/folic acid
 - anaemia 86–7, 88–9
 - epilepsy 100
 - nutrition 22–3, 25, 32–3
- forceps 130–1, 310–11, 323–4, 327
- fortified food stuffs 23
- fourth-degree perineal trauma 322
- fragile X syndrome 131–2
- fundal height measurements 189
- fundal pressure 267–8, 274–5
- furosemide 75

- GA see general anaesthesia
- garlic 60–1
- Gaskin manoeuvre 292
- gastric aspiration 263–4
- gauze dressings 353–4
- GBS see group B streptococcal
- GDM see gestational diabetes mellitus
- gemeprost 51–2
- general anaesthesia (GA) 328–9
- general antenatal care 7–13
- genetic disorders 131–2
- genital herpes see herpes simplex virus
- genital tract infections 165–6, 181–2, 200–7
- gentamycin 211, 377–8
- gestational age assessment 128–9
- gestational diabetes mellitus (GDM) 91–3
- gestational hypertension see pregnancy-induced hypertension
- gestational trophoblastic disease (GTD) 53–4
- ginger root extract 34–5
- glove protection 339–40
- glucose 138–9
- glyceryl trinitrate 67
- gonorrhoea 118, 200
- group antenatal education 14
- group B beta-haemolytic streptococcus 118
- group B streptococcal (GBS) infections 204, 205–6
- GTD see gestational trophoblastic disease
- guar gum 120
- H₂ receptor antagonists 264
- HAART see highly active antiretroviral treatment
- haemoconcentration 87
- haemoglobinopathies 97
- haemorrhoids 121–3
- hair removal, preoperative 334–6
- hands and knees position 299–300
- HCV see hepatitis C virus
- HELLP syndrome 57, 79–80
- helminthic infections 217
- heparin 42–3, 46–7, 95–7
- hepatitis C virus (HCV) 213
- heroin 18
- herpes simplex virus (HSV) 114, 203–4
- Hibiscrub 336–7
- high-risk pregnancies compromised fetal condition 134–6, 139–40 preterm births 157–8, 168–71
- higher order multiple pregnancies 192
- highly active antiretroviral treatment (HAART) 101–2
- HIV see human immunodeficiency virus
- HIVNET 012 regimen 105
- home births 251, 252
- home uterine monitoring 164
- home-based support 10–12, 17–18
- home-like birth settings 252–3
- homoeopathic remedies 219–20, 225, 247–8
- hospital births 251, 252
- hospitalisation hypertension 72–3 multiple pregnancies 190–2 preterm births 158–9
- HSV see herpes simplex virus
- human immunodeficiency virus (HIV) 101–13 antiretroviral drugs 101–11 caesarean section 106–9 micronutrients 112–13 mother-to-child transmission 101–12 vaginal disinfection 109–11 vitamin A 111–13
- human papilloma virus 114
- hyaluronidase 243

- hydatidiform moles 53–4
- hydralazine 74, 76
- hydrargaphen 205
- hydration
 - breastfeeding 393
 - caesarean section 352
 - compromised fetal condition 146
 - labour 261
 - preterm births 177–8
- hyperbaric bupivacaine 329
- hyperglycaemia 90–2, 94
- hypertension 55–85
 - categorisation 57
 - detection 69–70
 - generic protocol 56–8
 - HELLP syndrome 57, 79–80
 - mild/moderate 70–6
 - postpartum 75–6
 - prevention 56–69, 75–6
 - screening 139
 - severe 76–85
 - tight/very tight control 71–2
 - treatment 70–85
 - see also* eclampsia; pre-eclampsia
- hypnotherapy 280–1
- hypoglycaemia 92, 94
- hypotension 330–3
- hypotonic hydration 146
- hypoxaemia 144–5
- hypoxia 136
- hysterectomy 361
- idiopathic thrombocytopenia 94
- IGT *see* impaired glucose tolerance
- imidazoles 205
- immersion in water 258–9
- immunoglobulins 44–5, 46, 150
- immunotherapy 44–5
- impaired fetal growth
 - screening 132–3
 - treatment 141–6
- impaired glucose tolerance (IGT) 92–3
- impetigo herpeticiformis 125
- incomplete abortions 52–3
- indicator gloves 340
- individual antenatal education 14
- indomethacin 372
- induction 219–50
 - acupuncture 225, 244–5
 - amniotomy 229–31, 238, 240–1
 - caesarean section 223, 234–5
 - cervical ripening 224–9, 234, 240–4, 245–7, 248–9
 - generic protocols 224–5, 249–50
 - homoeopathic remedies 219–20, 225, 247–8
 - hypertension 77
 - indications 219–22
 - late pregnancy 224–5
 - mechanical methods 233–5
 - membrane sweeping 235–6
 - misoprostol 220–1, 231–3, 248–9
 - multiple pregnancies 195
 - oxytocin 219–20, 224, 227–31, 233, 236–9
 - primary reviews 223–49
 - prolonged pregnancy 187–8
 - prostaglandins 219–20, 224–33, 236–9, 246–7
 - screening 128–9, 135–6, 139, 151
 - secondary reviews 249–50
 - sexual intercourse 246–7
 - smooth muscle stimulation 224–5, 245–6
 - techniques 223–50
- indwelling bladder
 - catheterization 347
- ineffective uterine
 - contractions 287–90
- infections 114–18, 199–217
 - caesarean section 204, 213, 333–41, 347–9
 - congenital 212, 214–16
 - genital tract infections 165–6, 181–2, 200–7
 - helminthic 217
 - intraamniotic infection 210–11
 - labour 261–2, 279–80, 399
 - neonatal 203–4, 205–6

- parasitic 214–16
- preterm births 160–1, 165–7, 181–2, 199, 201
- screening 207, 216
- urinary tract infections 207–10
- vaccination 213–17
- see also human immunodeficiency virus
- influenza 114
- inherited coagulation defects 95
- insecticide-treated nets (ITNs) 117–18
- instrumental births see operative vaginal births
- insulin 90, 93–4, 222
- intermittent auscultation 308
- interrupted sutures 320–1
- intra-abdominal repair 346–7
- intra-amniotic infection 210–11
- intracervical prostaglandins 226–7, 228, 231
- intracranial haemorrhage 70
- intrapartum
 - fetal distress 311–12, 315–16
 - fetal scalp lactate sampling 308–9
- intrauterine devices (IUDs) 374–6
- intrauterine fetal death 91, 220–1
- intrauterine growth restriction (IUGR)
 - antenatal care 13, 29–30
 - hypertension 63–4, 69
- intrauterine pressure catheters (IUPC) 289–90
- intravaginal prostaglandins 225–6, 228, 231
- intravenous immunoglobulin 44–5, 46
- intravenous prostaglandins 237–8
- inversion of the uterus 362
- iodine deficiency 120–1
- iron supplementation 32–3, 86–9
- iron-deficiency anaemia 85–9
- isobaric bupivacaine 329
- isoimmunisation 150–1
- isosorbide mononitrate 248
- isotonic hydration 146
- isradipine 74
- itching 35
- ITNs see insecticide-treated nets
- IUDs see intrauterine devices
- IUGR see intrauterine growth restriction
- IUPC see intrauterine pressure catheters
- jaundice 150–1
- Joel-Cohen incisions 342–4
- ketanserin 80–1
- ketosis 260–1
- Kristellar manoeuvre 291
- labetalol 80–1
- labour 251–362
 - anaesthesia 261, 263–4, 266–7
 - assessment programs 254–5
 - caregivers 251–3, 255–6
 - compromised fetal condition 302–17
 - cord prolapse 317
 - episiotomy 264–8, 399
 - extrauterine pregnancies 357
 - first stage 257–8
 - infections 261–2, 279–80, 399
 - inversion of the uterus 362
 - maternal positions/mobility 257–8, 266–7
 - normal birth 264–80
 - operative vaginal births 315, 322–7
 - pain 280–7
 - perineal trauma 317–22
 - place of birth 251, 252–3
 - postpartum haemorrhage 268–71, 272–4, 276–8, 357–61
 - presentation 290–2, 294–302
 - retained placenta 359–61
 - routine care 253–80
 - second stage 264–8
 - shoulder dystocia 292–4

- labour (*continued*)
 - slow progress 287–92
 - spontaneous 259–60, 278–9, 287–9
 - third stage 268–80
 - see also caesarean section; induction; preterm birth
- lactation see breastfeeding
- lactational amenorrhoea 396
- lamivudine (3TC) 103, 106
- laparoscopy 122–3
- laxatives 38–9
- leg cramps 36–7, 39
- leg oedema 39–40
- leukocytes 44–5
- lidocaine 372
- lifestyle interventions 13
- lignocaine 372
- liver disease 119–20
- LMWH see low molecular weight heparin
- local anaesthesia 355
- local cooling 370–1
- low birthweight
 - antenatal care 8–9, 13, 20–1
 - hypertension 64
 - malaria 117–18
 - multiple pregnancies 190–1
- low-dose dopamine 85
- low molecular weight heparin (LMWH) 46–7, 96
- low-resource populations
 - antenatal care 32–3
 - context of care 5
 - human immunodeficiency virus 101–2
- lower genital tract infections 165–6, 181–2, 200–7
- lupus anticoagulant 46–7
- lytic cocktail 83–4

- MAC regimen 53–4
- McRobert's manoeuvre 292–3
- macrosomia 221–3
- magnesium sulphate
 - hypertension 77, 80–5
 - myasthenia gravis 98
 - nutrition 26–7
 - preterm births 162–3, 173–4, 178–9, 183, 399
- maintenance therapy 18, 182–6
- malaria 114–18
- manual fetal manipulation 137–8
- marine oil 63–4
- masculinisation 68
- mastitis 392–3
- maternal
 - control 3
 - hydration 146
 - oxygen administration 312, 329–30
 - positions/mobility 257–8, 266–7
- maternity waiting facilities 5
- Maylard muscle-cutting incisions 344
- MBUs see mother and baby units
- MDI see multiple daily injections
- measles 114
- mechanical induction 233–5
- meconium-staining 314–15, 316–17
- medicines during pregnancy 40–1
- mefloquine 116
- membrane sweeping 235–6
- Mendelson's syndrome 261, 264
- methotrexate 51–2, 53–4
- methyl dopa 71, 74, 76
- metronidazole 202–3
- MI see motivational interviewing
- micronutrients
 - human immunodeficiency virus 112–13
 - impaired fetal growth 144
- midwifery 2–3
- mifepristone 51–2, 239–40
- minerals 22–3, 25–33, 37
- miscarriage 41–54
 - chromosomal 41, 43
 - expectant care vs surgery 50–1
 - prevention 42–8, 399
 - recurrent 41, 42–4, 47–8

- spontaneous 43
- threatened 49–50
- treatment 49–54
- Misgav-Ladach incisions 342–3
- misoprostol 51–2, 220–1, 231–3, 248–9, 272–3, 358–9
- MMS *see* multiple-micronutrient supplements
- morbidity audit/feedback 3–4
- morbidly adherent placenta 361
- morning sickness 34–5
- mortality audit/feedback 3–4
- mother and baby units (MBUs) 385–6
- mother to infant HCV transmission 213
- mother to infant HIV transmission 101–12
- mother-infant rooming-in 391
- motivational interviewing (MI) 19
- Mouchel incisions 344
- moxibustion 297–8
- multidisciplinary teams 9
- multiple daily injections (MDI) 93–4
- multiple pregnancies 188–96
 - antenatal clinics 189–90
 - caesarean section 195–6
 - expectant management 195
 - higher order 192
 - induction 195
 - miscarriage 48
 - pregnancy reduction 192
 - screening 128–9
 - tocolytic therapy 189, 194
 - twin-twin transfusion syndrome 193
- multiple-micronutrient supplements (MMS) 32–3
- multivitamins 22–3, 37, 112–13
- music 280–1, 333
- myasthenia gravis 98

- nausea 34–5, 331–3
- near-infrared spectroscopy 306–7
- necrotising enterocolitis 154

- Neisseria gonorrhoeae see* gonorrhoea
- neonatal
 - bradycardia 73
 - encephalopathy 304–5, 306, 314
 - infections 203–4, 205–6
 - jaundice 150–1
 - mortality 21, 24
 - necrotising enterocolitis 154
 - periventricular
 - haemorrhage 171–3
 - respiratory disease 168–70, 399
 - tetanus 216–17
- neural tube defects 22–3, 25
- neurodevelopmental disorders 171–4
- nevirapine (NVP) 104–5, 106, 108–9
- nicardipine 74
- nifedipine 67, 75–6, 181, 184
- night blindness 28–9
- nimodipine 80–1, 82
- nipple pain 389–90
- nipple stimulation 225
- nitric oxide 66–7
- nitric oxide donors 179–80, 248, 299
- nitroglycerine 179, 248, 299
- non-affective psychosis 384–5
- non-clinical interventions 16
- non-closure of the
 - peritoneum 345–6, 399
- non-proteinuric gestational hypertension 71–2
- non-reactive
 - cardiotocography 137–8
- nonreassuring fetal heart rate 307–9
- non-steroidal anti-inflammatory drugs (NSAIDs) 369
- non-viable pregnancies 51–3
- NSAIDs *see* non-steroidal anti-inflammatory drugs
- nulliparity 267
- nutrients *see* micronutrients

- nutrition 22–33
 - breastfeeding 393–4
 - caesarean section 352
 - diabetes mellitus 91–2
 - energy/protein intake 23–5
 - labour 261
 - pre/periconceptual 22–3
 - vitamins/minerals 25–33
 - weight gain 23–5, 33
 - weight retention 366–7
- NVP *see* nevirapine
- nystatin 205
- obstetric anal sphincter injuries
 - (OASIS) 321–2
- occipito-posterior position 299–300
- occipito-transverse position 299–300
- oedema 39–40
- oestriol estimations 134–5
- oestrogens
 - breastfeeding 398
 - compromised fetal condition 143
 - induction 240–1
 - miscarriage 40–1, 45, 399
 - postnatal depression 380–1
- oligohydramnios 138, 146, 310, 314–15
- operative vaginal births 322–7
 - antibiotics 326
 - caesarean section 326–7
 - compromised fetal condition 304, 310–11, 315
 - forceps 310–11, 323–4, 327
- opiate dependence 18
- oral fluids 352
- oral prostaglandins 238–9
- overlap repair 321–2
- oxygen administration 144–5, 312, 329–30
- oxytocin
 - breastfeeding 377
 - induction 219–20, 224, 227–31, 233, 236–9
- labour 263, 269–71, 276
 - postpartum haemorrhage 358–9, 360
 - presentation 298
 - slow progress during labour 288–9
- oxytocin agonists 273–4
- oxytocin receptor antagonists 180–1, 185
- Ozzlo pillows 37–8
- P6 acupressure 34–5
- pain during labour 280–7
 - biofeedback 282
 - complementary therapies 280–1
 - epidural analgesia 282–7
 - TENS therapy 281–2
- parasitic infections 214–16
- parenthood 14
- partograms 259–60
- parvovirus B19 114
- paternal cell immunization 44–5
- pelvic pain 37–8
- pelvimetry 290–1, 399
- pemphigoid gestationis 125
- penicillin 212, 378
- periconceptual nutrition 22–3
- perinatal mortality 3–5, 20–1, 26, 376
- perineal
 - pain 368–72
 - shaving 261–2, 334–6, 399
 - trauma 265–7, 317–22, 368–72
- periodontal disease 166
- peritoneum, closure/non-closure 345–6, 399
- periventricular haemorrhage 171–3
- PET *see* pulsed electromagnetic energy
- PETRA regimens 103, 106
- Pfannenstiel incisions 342–4
- phenobarbital 150–1, 171–2
- phenobarbitone 100
- phenylephrine 332
- phenytoin 77, 84–5, 100

- physical activity
 - advice during pregnancy 21–2
 - diabetes mellitus 90–1
 - hypertension 59–60
 - weight retention 366–7
- physiotherapy 38
- PIH see pregnancy-induced hypertension
- piles 121–3
- pillows 37–8
- piracetam 315–16
- place of birth 251, 252–3
- placental
 - abruption 197, 198
 - cord drainage 278–9
 - delivery 347–8
 - function 134–5
 - insufficiency 145–6
 - retention 359–61
- placental-site trophoblastic tumour (PSTT) 54
- placenta praevia 197–8
- planned early birth see induction
- plasma volume expansion 78–9, 145
- Plasmodium vivax* 118
- pneumococcal infections 213–14
- postnatal care 1–2, 363–98
 - basic care of mother and baby 363–8
 - breastfeeding 363–5, 367–8, 374, 377, 386–98
 - discharge 365–6
 - education 363–4, 372–4, 387–9
 - family planning 372–6
 - medical conditions 377–8
 - perinatal death 376
 - perineal pain 368–72
 - psychiatric illness 378–86
- postnatal depression 378–83
- post-operative care 352–4
- postpartum
 - endometritis 199–200, 207, 377–8
 - hypertension 75–6
 - iron-deficiency anaemia 88–9
 - morbidity 107–8
- postpartum haemorrhage (PPH)
 - labour 268–71, 272–4, 276–8, 357–61
 - primary 358–9
 - retained placenta 359–61
 - secondary 359
- post-term labour 128–9, 138, 187–8
- postural management 294–5, 299–300
- PPH see postpartum haemorrhage
- PPROM see preterm prelabour rupture of the membranes
- preconceptional nutrition 22–3
- prednisone 46
- pre-eclampsia
 - antenatal care 24, 26–7, 28–30, 31–3, 48
 - Chinese herbal medicines 74–5
 - definition 55, 57
 - generic protocol 56–8
 - HELLP syndrome 57, 79–80
 - multiple pregnancies 189
 - prevention 56–69
 - screening 58, 140–1
 - severe 76–85
 - treatment 76–85
- pre-existing gestational hypertension 71–2
- pregnancy reduction 192
- pregnancy-induced hypertension (PIH) 24, 55, 57
- prelabour rupture of the membranes (PROM) 153–6, 199–200
 - induction 219–20
 - preterm births 160, 181
- prenatal
 - alcohol exposure 19–20
 - corticosteroids 168–9
 - diagnosis 7, 129–32
- preoperative hair removal 261–2, 334–6, 399

- presentation 294–302
 - breech births 291–2, 294–9, 300–2
 - external cephalic version 294–9
 - fetal malposition 299–300
 - fetopelvic disproportion 291–2
 - multiple pregnancies 196
 - pelvimetry 290–1, 399
 - postural management 294–5, 299–300
- preterm births 156–87, 399
 - antenatal care 9–10, 13, 20–1, 25–6, 29–30
 - antenatal clinics 157–8
 - high-risk pregnancies 157–8, 168–71
 - home uterine monitoring 164
 - infections 118, 160–1, 165–7, 199, 201
 - labour inhibition/treatment 174–82
 - maintenance therapy 182–6
 - mode of birth 186–7
 - nutrition 27
 - prediction 157–8, 161, 164, 165–6
 - prenatal treatment for
 - babies 167–74
 - prevention 157–67
 - recurrent 182–6
 - risk scoring systems 161
 - screening 165–6
 - tocolytic therapy 164–5, 174–81, 182–6
 - umbilical cord clamping 275–6, 278, 399
- preterm prelabour rupture of the membranes (PPROM) 153–5, 160, 220
- primary postpartum haemorrhage (PPH) 358–9
- probenecid 200
- probiotics 166–7
- progesterone
 - induction 239
 - preterm births 157, 163–4, 182
- progestogens
 - breastfeeding 398
 - hypertension 67–8
 - miscarriage 47–8, 49
 - postnatal depression 380–1
- proguanil 115
- prolonged pregnancy 128–9, 138, 187–8
- PROM *see* prelabour rupture of the membranes
- prostaglandins
 - induction 219–20, 224–33, 236–9, 246–7
 - labour 272–3, 360
- protein intake 23–5
- proteinuria 55, 57, 58, 71
- prurigo gestationis 125
- pruritic urticarial papules 125
- PSTT *see* placental-site trophoblastic tumour
- psychiatric disorders 123–5, 378–86
- psychological interventions 379–80
- psychosis, non-affective 384–5
- psychosocial interventions 18–20, 123–4, 378–80
- pulmonary embolism 98
- pulse oximetry 305–6
- pulsed electromagnetic energy (PET) 370–1
- pyelonephritis 207–9
- pyrexia 210
- pyroxidine *see* vitamin B6
- quinine 116
- RA *see* regional anaesthesia
- rapid negative pressure 324–5

- rashes 35
- RDS see respiratory distress syndrome
- rectal analgesia 355–6, 368–9
- rectal mucosa injury 322
- recurrent
 - miscarriage 41, 42–4, 47–8
 - preterm births 182–6
- red cell isoimmunisation 150–1
- reflexology 40
- regional anaesthesia (RA) 328–33
- relaxin 242
- renal disease 99
- repeat
 - caesarean section 223, 354–5
 - digital cervical assessment 9–10
- respiratory disease
 - maternal 119
 - neonatal 168–70
- respiratory distress syndrome (RDS)
 - caesarean section 356–7
 - preterm births 167–8, 170
- retained placenta 359–61
- retinal haemorrhages 324
- Rhesus alloimmunisation 147–9
- rigid vacuum extractors 325
- risk scoring systems 161
- ritodrine 179, 183
- ROM see rupture of the membranes
- rooming-in 391
- routine assessment see screening
- rubella 114
- rupture of the membranes (ROM)
 - induction 219–20, 229–31, 238, 240–1
 - prelabour 153–6, 160, 181, 199–200, 219–20
- rutosides 39–40, 122
- S-adenosylmethionine (SAdMe) 120
- salt intake 61–2, 359–61
- SAdMe see S-adenosylmethionine
- scalp lactate sampling 308–9
- SCD see sickle cell disease
- schizophrenia 385–6
- screening 127–9
 - blood group
 - incompatibilities 147–51
 - conservative management 140–6
 - counselling 132
 - fetal death 151
 - fetal wellbeing 133–41
 - genetic disorders 131–2
 - gestational age assessment 128–9
 - impaired fetal growth 132–3, 141–6
 - infections 207, 216
 - prenatal diagnosis 129–32
 - preterm births 165–6
 - psychosocial 378–9
 - symphysis-fundal height 127–8
- secondary postpartum haemorrhage (PPH) 359
- secondary thrombocytopenia 94–5
- selective termination 192
- self-diagnosis of active labour
 - onset 254
- semen 246–7
- sexual intercourse 246–7, 369–70
- SGA see small-for-gestational-age
- shoulder dystocia 292–4
- sickle cell disease (SCD) 97
- single deepest vertical pockets 138
- skin diseases 125
- skin staples 349–50
- skin-to-skin contact (SSC) 364–5
- slow progress during labour 287–92
 - adverse outcomes 289–90
 - amniotomy 287–9
 - fetopelvic disproportion 290–2
 - ineffective uterine
 - contractions 287–90
 - oxytocin 288–9
- small-for-gestational-age (SGA) births
 - hypertension 64, 68, 73–4
 - nutrition 24, 27
- smoking 2, 20–1

- smooth muscle stimulation 224–5, 245–6
- social factors 8–9, 10–13
- sodium nitroprusside 248
- sodium valproate 100
- soft vacuum extractors 325
- spectinomycin 200
- spinal anaesthesia 329, 330–1
- spontaneous
 - labour 259–60, 278–9, 287–9
 - miscarriage 43
- SSC *see* skin-to-skin contact
- SSIs *see* surgical site infections
- staples 349–50
- stavudine (d4T) 106
- stepwise negative pressure 324–5
- stillbirths 21, 24, 48
- stretch marks 35–6
- striae gravidarum 35–6
- suicide 378–9
- sulfadoxine-pyrimethamine 115
- supplementation *see* nutrition
- surgical
 - face masks 340–1
 - gloves 339–40
 - techniques 341–51
- surgical site infections (SSIs) 333–41
- sutures 319–21, 348–50
- sweeping of the membranes 235–6
- symphysiotomy 291–2
- symphysis-fundal height 127–8
- symptomatic urinary tract
 - infections 209–10
- symptoms during pregnancy 33–40
- syntometrine 273–4, 358–9
- syphilis 118, 212

- TBAs *see* traditional birth attendants
- TED *see* venous thromboembolic disease
- teenage mothers 10–11
- TENS *see* transcutaneous electrical nerve stimulation
- terbutaline 183, 184–6
- terconazole 205

- term prelabour rupture of the membranes 155–6
- termination
 - induction 220–1
 - miscarriage 42, 52–3
 - multiple pregnancies 192
- tetanus 216–17
- Thai-CDC regimen 104, 105
- third party donor leukocytes 44–5
- threatened miscarriage 49–50
- thrombocytopenia 94–5
- thromboembolic disease (TED) 72, 96–8
- thrombophilias 95–7
- thrush *see* candidiasis
- thyroid disease 120–1
- thyrotropin-releasing hormone (TRH) 170–1, 399
- timolol 76
- tocodynamometry 289–90
- tocolytic therapy
 - caesarean section 351
 - calcium channel blockers 175–6, 181, 183–4
 - compromised fetal condition 310–12
 - COX inhibitors 176–8
 - magnesium sulphate 183
 - maintenance therapy 182–6
 - multiple pregnancies 189, 194
 - nitric oxide donors 179–80
 - oxytocin receptor
 - antagonists 180–1, 185
 - preterm births 164–5, 174–81, 182–6
 - terbutaline 183, 184–6
 - topical anaesthesia 371–2
- toxaemia *see* pre-eclampsia
- toxoplasmosis 118, 214–16
- traditional birth attendants (TBAs) 4–5
- transcutaneous electrical nerve stimulation (TENS) 281–2

- transcutaneous
 - electrostimulation 145–6
- TRH *see* thyrotropin-releasing hormone
- trichomoniasis 202–3
- Trichuris trichura* 217
- Trofolastin 36
- trophoblast membranes 44–5
- twin-twin transfusion syndrome 193

- UDCA *see* ursodeoxycholic acid
- ultrasound
 - morbidly adherent placenta 361
 - postnatal care 369–70
 - screening 127, 128–9, 138, 139–40
- umbilical cord
 - clamping 275–6, 278, 399
 - compression 313–14
 - prolapse 317
 - topical care 279–80
 - traction 274–5
- umbilical vein injection (UVI) 276–7, 359–61
- unfractionated heparin 46–7, 96
- ureaplasma 201
- urinary tract infections (UTIs) 207–10, 337
- ursodeoxycholic acid (UDCA) 120
- uterine
 - contractions 287–90
 - cramping 377
 - hyperstimulation 232, 234–5, 238, 242
 - incisions 344–6
 - inversion 362
 - involution 377
 - massage 277–8
 - muscle relaxants 49–50
 - see also* tocolytic therapy
- UTIs *see* urinary tract infections
- UVI *see* umbilical vein injection

- vaccination 213–17
- vacuum aspiration 52–3
- vacuum extraction 323–7
- vaginal birth after caesarean (VBAC) 14, 16
- vaginal candidiasis 205
- vaginal disinfection 109–11
- vaginal trichomonas 118
- valacyclovir 203–4
- valproate 100
- varicella 114
- varicose veins 39–40
- VBAC *see* vaginal birth after caesarean
- venous thromboembolic disease (TED) 72, 96–8
- ventouses *see* vacuum extraction
- Verum ointment 36
- vibroacoustic stimulation 136–7, 307–8
- viral infections 114
 - see also* human immunodeficiency virus
- vitamin A
 - breastfeeding 393–4
 - human immunodeficiency virus 111–13
 - infections 217
 - nutrition 28–9
- vitamin B6 27–8, 34–5
- vitamin C 29–30, 64–5
- vitamin D 30–1
- vitamin E 31–2, 64–5
- vitamin K 172–3
- vitamins
 - antenatal care 22–3, 25–33, 37, 48
 - human immunodeficiency virus 111–13
- vomiting 34–5, 331–3
- vulvovaginal candidiasis 118

- waiting facilities 5
- water gymnastics 37–8
- water immersion 258–9

weanling's dilemma 395
weight gain 23–5, 33
weight retention 366–7
WHO/UNICEF initiatives
388–92
Wood's method 292
work during pregnancy 21–2
workplace interventions 390–1

wound closure 348–50
wound drainage 350–1

x-ray pelvimetry 290–1, 399

Zanavelli's method 292
zidovudine (ZDV) 103–6, 107–9
zinc 25–6