

Index

Note: SFBT = solution focused brief therapy

- A&E Departments 19–20, 189–90
- ABC model 58–9
- abuse survivors 7, 14, 27–8
 - see also* trauma survivors
- acceptance
 - dialectical behaviour therapy 56, 57
 - therapeutic relationship 10, 113
- access points to services 26, 189–91
- Accident & Emergency Departments 19–20, 189–90
- acknowledgment 91, 92, 98, 100–1, 108, 117
- addiction 73
- adolescents
 - dialectical behaviour therapy 57–8
 - wise old you question 143
- affirming questions 71, 73, 97, 137–8
- aggression 55
- alcohol abuse 14, 17, 81, 145
- Alex, case study 180–2
- anger 54–5, 146, 147–8
- antidepressants 36, 40–2, 194–5
- anxiety, practitioners' 9
- assertiveness skills 55
- assessment of suicide risk 20–1, 31–7, 123–30, 198
- attempted suicide
 - definition 24
 - intervention effectiveness 11–12, 190
 - lessons to be learned from 18, 52
 - mental health policy 15
 - World Health Organization 16
- awfulisation 58
- baby (small) steps, SFBT 88–9, 148–9
- Barry, case study 178–9
- Beck Hopelessness Scale (BHS) 51–2
- behaviour therapy *see* cognitive behavioural therapy
- biological model, depression 25, 64–5
 - see also* medical model
- blinkered thinking 49, 56, 118, 132
- bridging statements 101–2
- brief therapy 67–8
 - solution focused *see* solution focused brief therapy

- care-givers *see* practitioners
- care plans, Culverhay CMHC 39–40
- case studies 151–88
- cat and mouse game 145
- catastrophisation 58
- catharsis 108
- change
 - dialectical behaviour therapy 56, 57
 - solution focused theory of 68–71
- child abuse survivors 7, 27–8
- child neglect
 - author's experience 7
 - case study of Reg 151–76
- client-centred therapy *see* person-centred counselling
- clients
 - case studies 151–88
 - collaboration in therapy 39, 50–1
 - encounters with practitioners *see* therapeutic encounters
 - lead in SFBT 103, 104, 106
 - 'radar' 9, 10, 110–11, 112
 - relationship with practitioner *see* therapeutic relationship
 - SFBT *see* solution focused brief therapy
- clinicians *see* practitioners
- clues, SFBT 96–7, 102–3, 130–8
- cognitive behavioural therapy (CBT)
 - 51–5
 - dialectical behaviour therapy 55–8
 - rational emotive behaviour therapy 58–9
- cognitive constriction 49, 56
- cognitive restructuring 53
- collaboration, clinician–client 39, 50–1
 - see also* therapeutic relationship
- communication 6–7
 - dialectical behaviour therapy 57
 - of hope 139–40
 - non-verbal 6–7, 111–12, 113, 123–4, 126–8
 - suicidal intent 123–8
 - training 193
 - transactional analysis 62–3
- community institutionalisation 46
- Community Mental Health Centres (CMHCs) 39–40
- competence, practitioners' 8–9, 193
- compliments 90–1, 100
- confidence
 - communication of 118–19
 - scaling 88
- congruence, suicide intent 123–8, 129–30
- contracts, therapy 54
- control
 - loss of 128, 138
 - self-harm and 27
- conversational flow 107, 187
- coping questions 84
- cost effectiveness, suicide prevention 12–14, 175
- counselling, person-centred *see* person-centred counselling
- counselling skills training 193
- counsellors *see* practitioners
- crematorium scenario question 144–5, 181–2
- crisis interventionists 21
- crisis interventions
 - access points 26, 189–91
 - cognitive behavioural 53–4
 - rational emotive behaviour therapy 58
 - SFBT 122–3
- crucial first ten minutes 109–19, 122–3, 213
- cutting (self-harm) 27
- deathbed scenario (DBS) 145–6
- deep empathy 113–14, 122–3
- degrimming 99–100, 183
- deliberate self-harm 27–8
- depression 194–5
 - antidepressants 36, 40–2, 194–5
 - cognitive behavioural therapy 51–2, 56, 58
 - definition 25
 - dialectical behaviour therapy 56
 - endogenous 64–5

- rational emotive behaviour therapy 58
- research evidence for interventions 12, 41
- suicide risk assessment 20, 21, 31, 35, 36, 123–30
- trough of 127
- World Health Organization 16, 17
- despair, practitioners' 140
 - see also* hopelessness
- dialectical behaviour therapy (DBT) 55–8
- distress
 - defining suicide 23, 24
 - practitioners' 9
- doctors *see* general practitioners
- double blind trials 14, 196–7
- drug abuse 73, 145
- drug therapy 40–2, 194–5
 - self-poisoning with 18, 40, 194
 - suicide risk assessment 36
- dual awareness exercise 156
- eating disorders 14
- ego states 61–3
- emotional abuse 7
- emotional dysregulation 56
 - see also* pain, psychological
- empathy 10, 113–14, 122–3
- empowering questions, SFBT 71, 73, 97, 137–8
- empowerment 99, 140–1, 170, 175
- endogenous depression 64–5
- evidence-based practice *see* research evidence
- exceptions to problems, SFBT 84–5
- existential logotherapy 64–6
- eye contact 126
- facial expressions 128
- fast-forwarding the video technique 82, 83
- feelings tank 146–8
- financial costs, suicide prevention 12–14, 175
- 5 o'clock rule 80
- flashbacks 70–1, 156–7
- formula first session task (FFST) 69, 76–7, 97, 110
- free association 59
- functional analytic therapy 54
- game playing 63, 145
- general practitioners (GPs)
 - research evidence 12
 - suicide after contact with 18, 19, 20
 - suicide risk assessment 20, 37
 - training 190
- genuineness, practitioners' 9–10, 113
- goal development, SFBT 81–2, 138–9
- going with the client 98, 153, 175
- Gotland Study 12
- government policy, mental health 14–16, 195
- graveside scenario question 144–5, 181–2
- guaranteeing against suicide principle 130–1
- guilt 59, 65
- healing letters exercise 163–7, 172
- health professionals *see* practitioners
- healthier nation policy 14–16
- helpers *see* practitioners
- helping model 50–1
- helplessness
 - learned 54
 - practitioners' 8–9
- helplines 189–90, 192–3
- homework *see* tasks
- hope
 - cognitive behavioural therapy 52–3, 55, 57, 64
 - practitioners' 8–9, 139–40
 - SFBT 117, 138–40
- hope kits 53
- hopelessness
 - cognitive behavioural therapy 51–3, 56, 57, 58
 - practitioners' 140
 - SFBT 117, 118, 140

- hospital admissions 38–9, 190–2
- humour, degrimming 99–100, 183
- incongruence, suicide intent 123–8, 129–30
- interpretation, psychoanalysis 60
- interventions *see* preventive interventions
- Josephine, case study 182–5
- joyfulness, superficial 126–7
- keys, SFBT 96–7, 102–3, 130–8
- learned helplessness 54
- lethality, risk assessment 20–1, 32, 51
- life, meaning in 64–6
- logotherapy 64–6
- magic wand technique 82–3
- manualised approaches 197
- meaning in life 64–6
- medical model 42–6, 198
- medication 18, 36, 40–2, 194–5
- mental health policy 14–16, 195
- mental health practitioners *see* practitioners
- mental health services *see* service provision
- mental ill-health, medical model 43–6
- metaphors, SFBT 94–5
- mindfulness 55
- miracle question (MQ) technique 80–1, 82, 83, 84–5, 87, 141–3, 154, 184–5
- motivation, scaling 87–8
- multi-modal therapies 48–9, 56, 198
- national prevention targets 16, 195
- National Service Frameworks 15, 16
- negative beliefs, measurement 51–2
- negative thinking 118
 - see also* hopelessness
- non-possessive warmth 10
- non-verbal communication 6–7
 - crucial first ten minutes 111–12, 113
 - of hope 139
 - suicidal intent 123–4, 126–8
- noögenic neurosis 64, 65
- normalisation, SFBT 92, 108, 117, 136–7, 138
- optimism 118, 138
- outcomes, evidence base *see* research evidence
- overdoses, prescribed drugs 18, 40, 194
- pain, psychological
 - causal pathways 56
 - definition of suicide 23, 24
 - existential logotherapy 65
 - person-centred therapy 49–50
- paradox, small steps 88–9
- paradoxical intention 34–5, 54, 65, 92–3
- paraphrasing client's language 95–6, 193
- patients *see* clients
- person-centred counselling 48–51
 - therapeutic relationship 9–10, 48, 49–50
- personal hygiene 127
- personality disorders 57
- pessimism 118
- Peter, case study 179–80
- political suicides 28–9, 30
- possibility generation, SFBT 98–9
- postural clues, suicidal intent 126
- practitioners
 - competence 8–9, 193
 - encounters with clients *see* therapeutic encounters
 - qualities 9–10
 - relationship with clients *see* therapeutic relationship
 - skills 15, 50–1, 193
 - training 190, 192–4, 195
 - in empathy 114
 - mental health policy 15

- SFBT 107, 220
 - suicide risk assessment 37
 - therapeutic relationship 18–19, 119
 - use of book 1–2
- pre-session change, SFBT 69, 73, 134–5
- presuppositional language 71, 89–90, 97, 110, 130–2, 137–9, 143–4
- prevention, definition 25–7
- preventive interventions 37–40
 - pharmacological *see* medication
 - psychotherapeutic *see* psychotherapy
 - research evidence 11–12, 38, 41, 196–7
 - secondary prevention 26
 - World Health Organization 16–17
- primary care
 - access points 26, 189–90
 - mental health policy 15, 16
 - suicide after contact with 18, 19, 20, 21, 191
 - suicide risk assessment 20, 21, 36, 37 *see also* general practitioners
- primary prevention, definition 26
- problem focused therapy 67, 103, 177, 195–6, 198
 - solution focused alternative *see* solution focused brief therapy (SFBT)
- problem-free talk (PFT) 75, 77–9, 112
- problem land 79–80
- problem solving
 - CBT technique 53
 - dialectical behaviour therapy 57
 - SFBT compared 104
- psychache 24, 56
 - see also* pain, psychological
- psychiatrists 18, 20–1, 37
- psychoanalytic therapy 59–61
 - see also* transactional analysis
- psychotherapy 47–68
 - cognitive behavioural 51–9
 - dialectical behaviour 55–8
 - existential logotherapy 64–6
 - person-centred 48–51
 - psychoanalytic 59–61
 - rational emotive behaviour 58–9
 - research evidence 11–12, 196–7
 - solution focused *see* solution focused brief therapy (SFBT)
 - transactional analysis 61–3
- psychotropic medication 40, 41
- purpose in life (P-I-L) 65, 66
- ‘radar’, clients’ 9, 10, 110–11, 112
- randomised controlled trials 11
- rapport building 9–10, 113, 114, 115, 122–3, 213
- rational emotive behaviour therapy (REBT) 58–9
- Reg, case study 151–76
- research evidence 11–12, 38, 41, 196–7
- resources
 - cost of suicide prevention 12–14, 175
 - identification of personal 53
- risk assessment 20–1, 31–7, 123–30, 198
- risk factors 35–6
- risk management 37–40, 198
- Samaritans 24, 190
- Saving Lives: Our Healthier Nation* 14–16
- scaling tool, SFBT 85–8, 133–5, 154–5
- secondary care 189–92
 - mental health policy 15, 16
 - risk assessment 20–1, 36–7
 - risk management 38
 - suicide after contact with 18, 19–21, 191
- secondary prevention, definition 26–7
- self-harm, definition 27–8
- self-punishment 59
- service provision 31–46
 - access points 26, 189–91
 - medical model 42–6, 198
 - medication 40–2
 - mental health policy 16, 195
 - prevention targets 16, 195

- service provision (*continued*)
- quality 191–2
 - risk assessment 31–7
 - risk factors 35–6
 - risk management 37–40, 198
- service users *see* clients
- sexual abuse survivors 14, 27–8
- silence, use of 83
- Simon, case study 185–7
- skills
- clients' 53, 54, 75, 97, 140
 - practitioners' 15, 50–1, 192–3
- small steps, SFBT 88–9, 148–9
- social skills training 54
- solution focused brief therapy (SFBT)
- 67–108, 195–8
 - case studies 151–88
 - catharsis 108
 - clients' feelings 108, 146–8
 - conversational flow 107, 187
 - cost effectiveness 13–14, 175
 - crucial first ten minutes 109–19, 122–3, 213
 - evidence base 196–7
 - focus on the positives 105–6
 - guiding principles 72–5
 - key points 149–50
 - manualised approach 197
 - myths about 102–8
 - number of episodes of 104–5
 - origins 67–8
 - pre-session change 69, 73, 134–5
 - starting the first session 112–14, 122–3
 - sticking plaster criticism 102–3, 106–7
 - theory of change 68–71
 - tools and techniques 76–102, 107, 122–49, 213–19
 - acknowledgment 91, 92, 98, 100–1, 108, 117
 - assessing suicidal intent 123–30
 - bridging statements 101–2
 - clues 96–7, 102–3, 130–8
 - compliments 90–1, 100
 - deathbed scenario 145–6
 - degrimming 99–100, 183
 - eliciting suicidal ideation 128–30, 182–3, 186–7
 - end of session tasks 101–2, 148–9
 - exceptions to the problem 84–5
 - fast-forwarding the video 82, 83
 - feelings tank 146–8
 - formula first session task 69, 76–7, 97, 110
 - goal development 81–2, 138–9
 - hearing out the problem 79–80, 105
 - keys 96–7, 102–3, 130–8
 - magic wand 82–3
 - metaphors 94–5
 - miracle question 80–1, 82, 83, 84–5, 87, 141–3, 154, 184–5
 - normalisation 92, 108, 117, 136–7, 138
 - paradoxical intention 92–3
 - possibility generation 98–9
 - presuppositional language 71, 89–90, 97, 110, 130–2, 137–9, 143–4
 - problem-free talk 75, 77–9, 112
 - question examples 128–49
 - rapport building 122–3, 213
 - scaling 85–8, 133–5, 154–5
 - session ending 100–2
 - small steps 88–9, 148–9
 - stories 93–4
 - using the client's language 95–6
 - validation 91–2, 101, 108, 117
 - wise old you question 143–4
 - worst case scenario 144–5, 181–2
 - training 107, 220
- solution forced therapy 103
- solution land 79–80
- specialist mental health services *see* secondary care
- stigma 45
- 'Stop!' technique 70

- stories, SFBT 93–4
 stress, practitioners' 9
 substance abuse 16, 17, 81, 145
 suffering, existential logotherapy 65
 suicidal ideation
 continuum of 33–4
 questions to elicit 128–30, 182–3, 186–7
 scale of 51
 see also suicidal intent, assessment
 suicidal intent, assessment 20–1, 31–7, 123–30, 198
 suicidality, use of term 24
 suicide, definition 23–5
 suicide attempters *see* attempted suicide
 suicide bombers 29–30
 suicide encounters *see* therapeutic encounters
 Suicide Intent Scale (SIS) 51
 suicide mode 52
 suicide option 7–8, 24, 28, 52, 53, 129, 136–7, 170
 suicide prevention, definition 25–7
 suicide rates
 UK mental health policy 15, 16, 195
 World Health Organization 16–17
 suicide risk *see* risk assessment; risk management
 suicidology 2, 24

 task forces 16, 17
 tasks, SFBT 69, 76–7, 97, 101–2, 110, 148–9
 telephone helplines 189–90, 192–3
 terrorism, suicide bombers 29–30
 tertiary prevention, definition 26
 therapeutic alliance 114–15
 therapeutic conversations, flow 107, 187
 therapeutic empathy 114
 therapeutic encounters
 crucial first ten minutes 6–7, 8, 9, 10, 109–19, 122–3, 213
 practitioners' feelings 8–9
 risk assessment 20–1, 36–7, 123–30
 suicide committed after 17–21, 119, 191
 see also psychotherapy; solution focused brief therapy (SFBT); therapeutic relationship
 therapeutic relationship
 building 9–10, 113, 114–19, 122–3, 213
 importance 114–15
 person-centred counselling 9–10, 48, 49–50
 research evidence 11
 training for 18–19, 119, 193
 therapists *see* practitioners
 therapy contracts 54
 training 190, 192–4, 195
 in empathy 114
 mental health policy 15
 SFBT 107, 220
 suicide risk assessment 37
 therapeutic relationship 18–19, 119
 transactional analysis (TA) 61–3
 transference 60
 trauma survivors, solution focused therapy 14, 70–1
 see also abuse survivors; Reg, case study
 treatment, definition 25–7
 pharmacological *see* medication
 psychotherapeutic *see* psychotherapy
 see also preventive interventions
 trust, development 115–16, 213
 24-hour helplines 189–90, 192–3

 unconditional positive regard 10, 113
 unconscious, the 59, 60, 61

 validation
 SFBT 91–2, 101, 108, 117
 strategies 55
 very, very small steps 148–9
 video fast-forwarding technique 82, 83

warmth, practitioners' 10

will to meaning 64, 65

wise old you (WOY) question
143–4

workers *see* practitioners

working relationship *see* therapeutic
relationship

World Health Organization (WHO)

16–17, 193–4

worst case scenario question 99, 144–5,
181–2

youth suicide 17

see also adolescents