



Index

Figures are indicated by *italic numbers*

- acetonitrile, in CTV-based structures 141,
142, 143, 146–7, 147, 148
- acridine ligands 236–7
- active pharmaceutical ingredients (APIs) 25,
27
- co-crystals 27, 35
- with carbamazepine 36–45
- adamantane-type units 140, 141, 173–4, 283,
284
- adipic acid, adducts, with DABCO 8
- adsorption–desorption isotherms, methane in
dicopper paddlewheel complexes 215
- alkali-metal ions, in anionic metal-carbonate
networks 283, 287–8, 291, 294
- anamnox* bacteria, membrane lipids 59
- anionic metal-carbonate network
282–94
- body-centered cubic structures 289–91
- cadmium-based 289–90
- calcium-based 290–1
- cobalt-based 291–4
- copper-based 283–9
- face-centered cubic structures 291–4
- manganese-based 289–90
- antiferromagnetism 305
- aqua complexes, hydrogen-bonding interactions
71
- argentophilic interactions 249–55
- see also* silver. . .
- aromatic edge–edge C–H . . . N dimers 93–5,
98
- aromatic rings
- edge-to-face interactions 93, 220
- offset face-to-face interactions 92–93, 220
- aurophilic interactions 220, 256–8
- see also* gold. . .
- aza-1,3-*peri* aromatic hydrogen interactions
96, 97, 98
- ball-and-socket assemblies 137, 153
- ‘beads-on-thread’ structures 187
- 1,2,4,5-benzenetetracarboxylic acid, as linear
template 62–3
- 2,2'-biimidazole, in nickel complexes 226,
229
- binary bismuth xanthates 130–3
- binary mercury dithiocarbamates 128–30
- binary zinc xanthates 123–5
- biological complexes, hydrogen-bonding
interactions in 235–7
- 2,2'-bipyridine, in metal complexes 231,
232–3, 233
- 2,4'-bipyridine, in silver complexes 251, 253
- 4,4'-bipyridine
- co-crystals 31, 32, 38, 39
- in hydrogen-bonded supramolecules 31,
39, 220–1
- in metal coordination polymers 245, 247–8
- in silver complexes 250–1, 253
- as spacer ligand 214, 221, 223
- bipyridine adducts of zinc dithiophosphates
125–8

- 4,4'-bipyridylethane, in zinc coordination polymer 316
- 4,4'-bis(benzylammonium)stilbene
hexafluorophosphate, co-crystallization with crown ether 62
- bismuth xanthates, binary 130–2, 133
- bismuth(III) sulfide, morphology affected by precursors 131–3
- bis-*p*-phenylene[34]-crown[10], as linear template 62
- bis(phenylimino)isoindoline 68, 87
- trans,trans*-1,4-bis(4-pyridyl)-1,3-butadiene, co-crystallization with
5-methoxyresorcinol 59–60
- trans*-1,2-bis(2-pyridyl)ethene, co-crystallization with resorcinol 56, 56
- trans*-1,2-bis(4-pyridyl)ethene co-crystallization with
1,2,4,5-benzenetetracarboxylic acid 62–3
1,8-naphthalenedicarboxylic acid 61
resorcinol and derivatives 54, 55, 56, 56
tetrakis(4-iodoperfluorophenyl)erythritol 63–4
tricarballic acid 62–3
- 1,4-bis[2-(4-pyridyl)ethenyl]benzene, co-crystallization with resorcinol and derivatives 57–8
- trans,trans,trans*-1,6-bis(4-pyridyl)-1,3,5-hexatriene, co-crystallization with 5-methoxyresorcinol 59–60
- body-centered cubic structures, anionic metal-carbonate networks 289–91
- Borromean rings 178, 180
2-D → 2-D parallel interpenetration 179, 181
2-D → 3-D parallel interpenetration 178, 181
- 'brick-wall' structure 204
- 'bricks-and-mortar' inclusion systems 106–8
- brominated diphenyl diquinoline inclusion hosts, in bricks-and-mortar inclusion systems 107–8
- brominated diquinoline inclusion hosts 93, 94
edge–edge interactions 94
in layered inclusion compounds 106
- brominated diquinoxaline inclusion hosts 94
edge–edge interactions 94, 95
- p*-bromo-*N*-(*p*-dimethylaminobenzylidene)aniline 88
- N,N'*-bis-(4-bromophenyl)melamine/5,5'-diethylbarbituric acid polymorphic co-crystals 32, 33
- t*-butanol, hydrogen bonding with water molecules in cadmium cyanide networks 270, 271
- cadmium cyanide networks 269–71
effect of hydrogen bonding between water and *t*-butanol molecules 270, 271
- calcium aqua complex ion, in CTV-based structures 142
- calixarenes 136
back-to-back packing 140
- calixresorcinarene hydrogen-bonded networks 136
- Cambridge Structural Database (CSD)
AJAJEA–AJAJEA01 31–2, 33
analysis of data in 26
aza-1,3-*peri* aromatic hydrogen interactions 96
co-crystals 26, 28
copper(II) acetate 'paddlewheel' complexes 74
- EBIBEW 46
- ether-1,3-*peri* aromatic hydrogen interactions 95
- GASTRN10 43
- HADKUT–HADKUT01 29
- IHANIO 43
- JICTUK–JICTUK10 32, 33
- MACCID–MACCID01–MACCID02 31, 32
- MUROXA–MUROXA01 30–1
- OBIZII 43
- PIKLEA 46
- polymorphic co-crystals 28, 29–35
- QAWNAD 46
- QUIDON–QUIDON02 29–30
- thioether-1,3-*peri* aromatic hydrogen interactions 96
- VAPBAP 46
- VENLUV 46
- ZIGPAG–ZIGPAG01 32, 34, 34
- carballic acid, as linear template 62–3
- carbamazepine [5H-dibenz(b,f)azepine-5-carboxamide] 36
acetic acid solvate 43, 44
acetone solvate 39, 40
- 1,3,5,7-adamantanetetracarboxylic acid co-crystal 40–1, 42

- 4-aminobenzoic acid co-crystal 42, 43
benzoquinone co-crystal 37
4,4'-bipyridine co-crystal 38, 39
butyric acid solvate 43, 45
carboxylic acid co-crystals 39–44
dimer 36–7
dimethylsulfoxide solvate 39, 41
formamide solvate 39, 41
formic acid solvate 43, 44
nicotinamide co-crystal 38, 39
5-nitroisophthalic acid co-crystal 40, 42
pharmaceutical co-crystals 36–45
polymorphs 36
2,6-pyridinedicarboxylic acid co-crystal 41, 42
saccharin co-crystal 39, 39, 43, 45
 compared with Tegretol 45
terephthalaldehyde co-crystal 38
trimesic acid co-crystal 40, 42
carbonate complexes *see* anionic
 metal-carbonate networks
o-carborane
 molecular structure 136
 network structures
 with CTV 137, 138
 with CTV and fullerenes 137–8, 138
 with CTV and water 139, 140
carborane anions, in CTV-based structures 140–9
carboxylic acid groups
 coordination complexes with 222–4
 in paddlewheel complexes 195–217
catenane-like interactions 162
catenanes, interpenetrating networks as
 polymeric analogs 178, 183
cavitand hosts 135
CdSO₄ topology 178, 180, 257, 258
cerium(III) dihydroxybenzoquinone derivatives,
 hydrogen bonding with water molecules
 271–2, 273, 274
'Chinese blinds' network structure 187
chiral catalysts 301
chiral centers, helicity of coordination
 polymers affected by 306–8
chiral co-crystals, preparation by seeding 6
chiral ligands 239, 306
chloranilic acid 273
 lanthanide complexes 272–3, 276–7
chromium(II) acetate 'paddlewheel' complexes 197
chrysene 204
 interactions with dirhodium(II)
 tetra(trifluoroacetate) 204
clathrates
 active pharmaceutical ingredients 35
 applications 135–6
 edge–face (EF) interactions 93, 220
 halogenated heteroaromatic hosts 92–115
 meaning of term 91, 135
 offset face–face (OFF) interactions 92–93,
 97, 220
 pen structures 97–9
cobalt hexacyanide anion, as linker 201–2
cobalt saccharate 281
cobalt(III) zwitterionic sandwich complex
 [Co^{III}(η^5 -C₅H₄COOH)(η^5 -C₅H₄COO)]
 solvent-free reactions 13–18
 alkali-metal ions 17–18
 ammonia 14–15
 formic acid 16–17
 HBF₄ 15, 16
 hydrogen chloride 14
 volatile haloacids 15–16
co-crystal formers 35–6
co-crystals
 in Cambridge Structural Database 26
 meaning of term 26
 preparation of 4–6, 26–7
 grinding and milling 5–8, 27
 kneading 5–6, 27
 seeding 6, 7
 see also pharmaceutical co-crystals
collective properties 19
conductivity, mixed-valence rhodium-based
 paddlewheel complexes 206–7
conformational isomorphism 68, 87
conformational polymorphism 68, 87
coordination bonds, advantages and
 disadvantages 297
coordination networks, mechanochemical
 preparation of 10–13
coordination polymers
 cyclotrivertatrylene-based 144–52
 gold(I) cyanide-based 257–8
 hydrogen-bonded 297–317
 π – π interactions 239–58
 silver(I) cyanide-based 255
copper(I) imidazole complex 249, 252
copper(I) thiourea complex 72
 in crystalline array with crown ethers 72–3
copper(II) acetate hydrate, co-grinding with
 1,3-di(4-pyridyl)propane 12–13

- copper(II) acetate 'lantern'/'paddlewheel' complexes 73–4, 197
- copper(II) *N*-(2-hydroxybenzyl)-alanine complex 303–6
 helicity 306–7, 307
 magnetochemical properties 304–5
 packing of racemic crystals 307–8, 308
- copper(II)
N-(2-hydroxybenzyl)-1-aminocyclopentyl-1-carboxylic acid complex 312–13
- copper(II) *N*-(2-hydroxybenzyl)-glutamic acid complex, staircase helical coordination polymer 313–16
- copper(II) *N*-(2-hydroxybenzyl)-glycine complex, C=O $\cdots\pi$ interactions 308–10
- copper(II) *N*-(2-hydroxybenzyl)-valine complex, supramolecular isomerism 310–12
- corannulene 205
 interactions with dirhodium(II) tetra(trifluoroacetate) 205
- corrugated layers 244, 253
- covalent bonds, mechanochemical formation of 8–13
- covalent capture, and target-oriented organic synthesis 52
- [Cr(en)₃][Ni(CN)₅] \cdot 1.5H₂O 68
- crown ethers
 as hydrogen-bond acceptors 70
 as linear/planar templates 62, 136
 oxonium ions isolated in 85
- 15-crown-5 complexes 74–5, 76
- 18-crown-6 complexes 74, 75
 AgNO₃ complex 74
 effect of ion–dipole interactions on molecular shape 74, 75
 KNO₃ complex 74, 75
 mercury(II)-containing 81
 NaNO₃ complex 74, 75
 strong-acid hydrogen-bonded complex 75, 76, 77
 temperature-dependent distortion 76, 77
- crystal engineering
 concept first introduced 25
 meaning of term 1, 18, 25, 117, 135
- crystal packing interactions 69
 effect of disorder and 84
- CTV *see* cyclotrimeratrylene
- cuprophilic interactions 248–9
see also copper...
- Curie temperature 304
- cyanamide, interpenetrating hydrogen-bonded nets 172
- cyanuric acid/1,2-*bis*(4-pyridyl)ethene polymorphic co-crystals 29
- cyclodextrin inclusion compounds 135
 preparation by kneading 5
- cyclotrimeratrylene (CTV) 136–7
 as chelating ligand 136, 144–9
 coordination polymers 144–9
 Cs⁺ polymers 145–7, 147, 148
 K⁺ polymers 145, 146, 147–8, 148
 Na⁺ polymers 144, 144, 145, 145, 146, 146, 147, 148–9
 Rb⁺ polymers 145, 147, 148
 dimethoxy groups in 136, 137
 extended-arm derivatives 149, 150
 applications 137
 coordination polymers 150–2
 as hydrogen-bond acceptor 137–44
 hydrogen-bonded structures
o-carborane as H-bond donors 137–8
 with N–H and O–H donor groups 139–44
 molecular structure 136
- diamondoid networks 206–7, 207, 266, 267, 283
 hexagonal 266, 267–8, 312
 interpenetrating 161, 173–5, 271, 275
 in solid-state supramolecular transformations 306
- 1,4-diazabicyclo[2.2.2]octane (DABCO)
 adducts with
 adipic acid 7, 8
 ferrocenyl dicarboxylic acid complex 6–7
 malonic acid 7–8
 silver acetate 10–11, 12
 zinc chloride 11–12, 12
 molecular structure 7
 as pillar ligand, in dicopper paddlewheel complexes 214
- N,N'*-dibenzyl-4,13-diaza-18-crown-6, CH– π interactions 77
- dibromo diquinolines 93
 edge–edge interactions 94
 heteroatom-1,3-*peri* interactions 95–7
 in molecular pen structures 97–9
- 1,2-dicarbadodecaborane *see o*-carborane
- dicarboxylic acids, adducts, with DABCO 7–8

- 1,2-dichlorobenzene, in CTV-based structures 137–8, 138
- 1,4-diethynylcyclohexane-1,4-diol 87
- dihydroxybenzoquinone 272
- lanthanide complexes 271–2, 273–4, 275
- N,N'*-dimethylformamide (DMF), in CTV-based structures 139, 140, 145, 146, 148–9, 149
- dinitro diquinoline inclusion hosts, in molecular spheres 108–10
- diorganotin carboxylates 118–20
- diorganotin dichlorides 84
- diphenyl tin bis(xanthate) complex 81, 82, 83
- 4,4-diphenyl-2,5-cyclohexanedione 87
- dirhodium(II) tetra(trifluoroacetate) interactions with
- chrysene 204
 - corannulene 205
 - diiodine 202–3
 - elemental sulfur 203
 - π -ligands 203–4
 - polycyclic aromatic hydrocarbons 204–5
 - triphenylene 204
- see also* ‘paddlewheel’ dinuclear complexes
- diruthenium(II) complexes, magnetic properties based on 209–12
- diruthenium(II,III) tetracarboxylate complex cation, magnetic properties based on 208–9
- dithiocarbamate anion 123
- see also* mercury dithiocarbamates
- 1,1-dithiolate ligands
- structures containing 123–32
 - see also* bismuth xanthates;
 - dithiocarbamates; dithiophate anion;
 - mercury dithiocarbamates; xanthate anion; zinc dithiophosphates; zinc xanthates
- dithiophosphate anion 123
- see also* zinc dithiophosphates
- double-strand helicates 235–6, 239, 240
- Ag...Ag interactions 253
- double-stranded polymeric chains, gold(I) complexes 256, 258
- electrochemical artificial muscles 244
- entangled networks 185–7
- entangled rings 178, 180–1
- ether-1,3-*peri* aromatic hydrogen interactions 95, 96
- ethylenediammonium cinnamates 54, 55
- europium aqua complex ions, in CTV-based structures 142, 143
- extended supramolecular arrays, synthesis of 12–13
- face-centered cubic structures, anionic metal-carbonate networks 291–4
- ‘fake’ pharmaceutical co-crystals 45–6
- ferrocenyl dicarboxylic acid complex, adducts with nitrogen-containing bases 6–7
- flexible structures 182, 233–4, 248
- ‘gate-opening’ pressure for 182, 248
- four-connected networks 173–5, 266
- fullerenes
- network structure, with CTV and *o*-carborane 138, 138
 - see also* hemifullerene
- gas adsorption structures 182, 214–15
- ‘gate-opening’ pressure for 182, 215, 248
- gold(I) chain complexes, polymorphism in 256, 257
- gold(I) cyanide anion, coordination polymers with 257–8
- grinding and milling 5, 6–8, 27
- solvent-assisted 5, 9, 27
 - see also* kneading
- guanidinium cation 7, 283
- in CTV-based structures 139
- in metal-carbonate networks 283, 285–6
- halogenated heteroaromatic clathrate systems 92–115
- aromatic edge–edge C–H...N dimers 93–5, 98
 - bricks-and-mortar inclusion systems 106–8
 - edge–edge halogen interactions 99–101
 - edge–face (EF) interactions 93
 - general design of host molecules 92–3
 - heteroatom-1,3-*peri* interactions 95–7, 98
 - interlocking molecular grids 110–12
 - layer inclusion compounds 106
 - offset face–face (OFF) interactions 92, 93, 97
 - π -halogen dimer interactions 101–6
 - staircase inclusion compounds 101–5

- helical channels, in staircase coordination polymers 313–15
- helical coordination polymers
solid-state transformation to 3-D network structure 303–6
staircase topology 313, 314
- helicates 235–6
arc-shaped 237
double-strand 235–6, 239, 240, 253
multistrand 235–6
triple-strand 242
- helicity of coordination polymers, effect of chiral centers 306–8
- helix-with-helix structure 313–15
- hemifullerene 206
interactions with dirhodium(II) tetra(trifluoroacetate) 205–6
- heteromolecular seeding 6
- hexagonal channels
cadmium cyanide 269, 270
lanthanide mucates 281
see also honeycomb-like networks
- hexagonal diamondoid networks 266, 267–8, 312
- Hittorf's phosphorus 164–5
- homochiral chains, metal complexes 239, 242
- honeycomb-like networks 201–202, 206, 207, 211, 221, 224, 229, 266, 270, 299, 300
- hourglass-shaped channels 237, 238
- hydrated protons *see* oxonium ions
- hydrates 35
- hydrogen-bonded adducts, mechanochemical preparation of 6–8, 19
- hydrogen-bonded coordination polymeric structures 297–317
- hydrogen-bonding interactions
in anionic metal-carbonate networks 282–94
in cadmium cyanide network 269–71
in coordination complexes 220–37
with aquometal centers 220–2
biologically relevant complexes 235–7
with carboxylic acid groups 222–4
with CH \cdots X bonding sites 229–34
with imidazole groups 225–9
in coordination polymers 297
in crown ether complexes 70–7
factors affecting 117, 268
in halogenated heteroaromatic clathrate systems 94–101
in lanthanide chloranilate complexes 276–7
in lanthanide dihydroxybenzoquinone complexes 271–4, 275
in lanthanide mucates 282
strength of hydrogen bonds 220
in template-controlled solid-state reactions 54–64
in zinc saccharate 278–81
- hydroquinone/*p*-benzoquinone polymorphic co-crystals 29–30
- hydrothermal reaction conditions, coordination polymers affected by 233
- N*-(2-hydroxybenzyl)-aminoacids 298
C=O \cdots π interactions 308–10
as tridentate ligands in coordination complexes 298–316
- N*-(2-hydroxybenzyl)-1-aminocyclopentyl-1-carboxylic acid, molecular structure 313
- N*-(2-hydroxybenzyl)-glutamic acid, molecular structure 314
- N*-(2-hydroxybenzyl)-valine, molecular structure 311
- 2-[(*E*)-2-(2-hydroxy-5-methylphenyl)-1-diazenyl]-benzoic acid 77, 121
in triorganotin carboxylates 77–9, 120–3
- hypervalent compounds, crystal packing interactions 77
- imidazolate derivatives, coordination complexes with 249, 250
- imidazole groups, coordination complexes with 225–9, 249, 252
- in situ* building blocks for 'paddlewheel' dinuclear tetracarboxylate complexes 199, 199
porous properties based on 213–16
- inclined interpenetration 160
- inclusion compounds/systems *see*
bricks-and-mortar inclusion systems; layer inclusion compounds; staircase inclusion compounds
- intercalation 159
2-D \rightarrow 3-D architecture 246, 247–8
zipper-like structures 231, 232, 239, 241, 245
- interdigitating networks 157, 158, 185–187
- interlocking molecular grids, halogenated heteroaromatic clathrate systems 110–12
- interpenetrating networks 157–85
1-D nets 160, 162–4
1-D \rightarrow 1-D parallel interpenetration 162


- 1-D \rightarrow 2-D inclined interpenetration 162
- 1-D \rightarrow 2-D parallel interpenetration 162, 163
- 1-D \rightarrow 3-D inclined interpenetration 163, 164
- 1-D \rightarrow 3-D parallel interpenetration 162
- 1-D/2-D \rightarrow 2-D interpenetration 178
- 1-D/2-D \rightarrow 3-D interpenetration 178, 179
- 1-D/3-D interpenetration 178
- 2-D nets 160–1, 164–72
- 2-D \rightarrow 2-D parallel interpenetration 164–6, 179, 181
- 2-D \rightarrow 3-D parallel interpenetration 166–7, 178–9, 181
- 2-D inclined interpenetration 167–72
- triple interwoven 221–2, 224, 251, 253
- 2-D/3-D interpenetration 178, 180
- 3-D nets 161, 172–8
- 3-connected nets 172–3
- 3,4-connected nets 176, 178
- 3,5-connected nets 177, 178
- 4-connected nets 173–5
- 5-connected nets 175
- 6-connected nets 175–6
- cadmium cyanide 269
- diamondoid networks 161, 173–5, 271, 275
- triple interpenetrated 255, 256
- consequences of interpenetration 179, 181–3
- enclosed cavities formed by 182–3
- flexible structures 182
- microporous structures 182
- notation 159–61
- [Ru^{II,III}(O₂CR)₄]⁺-based 208, 209
- unusual interpenetration 178–9
- interpenetration
- inclined 160
- meaning of term 158
- parallel 160
- topology 160
- interwoven chains 186
- iron(II) aqua complex ion, in CTV-based structures 142, 143
- isomerism 310–12
- Kagomé lattice 211
- kneading 5–6, 19, 27
- examples of use 5, 17–18
- see also* solvent-drop grinding
- knots 3, 183
- ladder structure(s)
- CTV-based 139
- metal complexes 242–3
- [*n*]-ladderanes, target-oriented synthesis 59–60
- ladders, hydrogen-bonded 162, 163
- lanthanides
- chloranilic acid derivatives 272–3, 276–7
- dihydroxybenzoquinone derivatives 271–2, 273–4, 275
- muocate structures 281, 282
- lanthanoids *see* lanthanides
- lanthanum chloranilic acid derivative 272, 276
- lattice inclusion chemistry 135
- Laué diffraction method 71
- layer inclusion compounds 106
- layer structures 97–9, 130
- Lewis acid–base interactions 117
- linear tapes 32, 33, 63, 67
- linear templates
- reactivity controlled using 53–4
- resorcinol and derivatives as 52, 54–6
- Lonsdaleite network 266, 267–8, 312
- magnetic properties, diruthenium
- tetracarboxylate paddlewheel complexes 208–12
- magnetic susceptibility, temperature
- dependence, diruthenium tetracarboxylate paddlewheel complexes 209, 210, 211
- magnetochemical properties, Cu(II) complexes 304–5
- malonic acid, adducts, with DABCO 7–8, 8
- Mannich bases 298
- see also* *N*-(2-hydroxybenzyl)-aminoacids
- mercury compounds 79–81
- Hg(18-crown-6)(NO₃)₂ 81, 82
- Hg(18-crown-6)X₂ 81
- mercury(II) bis(dithiocarbamates) 79–81, 128–30
- compared with mercury(II) bis(xanthates) 79
- mercury(II) bis(xanthates), compared with mercury(II) bis(dithiocarbamates) 79
- metal-organic framework (MOF) structures 213, 233–4
- metallophilic interactions 220, 248–58
- argentophilic interactions 249–55
- aurophilic interactions 220, 256–8
- cuprophilic interactions 248–9

- methane, adsorption–desorption isotherms 215
- N*-methylurea/oxalic acid polymorphic co-crystals 30–1
- microporous structures
hydrogen-bonded networks 222, 224
interpenetrating networks 181–2
paddlewheel complexes 212–16
 π – π -bonded metal complexes 242, 243
- milling *see* grinding and milling
- molecular building-block approach 195, 196
- molecular conformation 87–8
- molecular grids
interlocking 110–12, 113–14
construction of 111, 113–14
- molecular hosts
halogenated heteroaromatic compounds 92–115
incorporation into network structures 136–53
meaning of term 135
- molecular magnets 304
magnetic ordering temperatures 304, 305
- molecular pen structures 97–9
interactions between layers 98, 99
interactions between pens 97, 100, 101
- molecular sieves 301
- molecular spheres 108–10
- molecular synthons 51
see also supramolecular synthons
- mucates 275
disadvantages as ligands 275, 277
positive aspects of use 277–8
see also lanthanide mucate structures
- multistrand helicates 235–6
see also double-strand...; triple-strand helicates
- muscle fibers 244
- nanotubes 242, 243
- 1,8-naphthalenedicarboxylic acid, as linear template 61
- NbO topology 255, 256
- nearest-neighbor interactions 69
- network structures
CTV-based 136–53
molecular hosts incorporated into 135–53
- networks
1-D nets
CTV-based structures 139
interpenetrating 160, 162–4
- 2-D nets
CTV-based structures 138, 139, 140, 144, 145
interpenetrating 160–1, 164–72
(6,3) nets 159, 266, 272, 276
- 3-D nets
CTV-based structures 140, 141, 143, 144
interpenetrating 161, 172–8
(10,3)-*a* nets 172, 266, 267
(10,3)-*b* nets 172
(4,4) nets/sheets 125, 158–9
interpenetrating 157–85
methods of maximizing packing efficiency 157
notation 159–61
open porous, applications for 157
shortest-circuit approach to analysis 183–5
3,12-connected net 140, 141
see also entangled networks; interpenetrating networks; self-penetrating networks
- nickel(II) nitrate, co-grinding with 1,10-phenanthroline 12
- non-covalent bonds, breaking and forming of 6–8, 19, 53
- non-covalent derivatization 26
- octagonal copper complex 249, 250
- octahedral geometry 140, 141
- olefins, solid-state [2+2] photodimerization of 54, 55, 58, 59
- oxa-bridged complexes, ruthenium complexes 86
- oxa-bridged diquinoline dibromide 95
- oxonium ions, isolation by crown ethers 85
- paddlewheel complexes
chromium(II) acetate complexes 197
copper(II) acetate complexes 73–4, 197
- paddlewheel dinuclear complexes
dicopper cores
[Cu₂(O₂CCH₃)₄] 201
electronic structure 197
in situ formation 199
porous properties 214–15
- dirhodium cores
electronic structure 197
preparation of preorganized building blocks 199
[Rh₂(O₂CCF₃)₄] 201, 202–6
[Rh₂(O₂CCH₃)₄] 202

- diruthenium cores
 electronic structure 197–8
 preparation of preorganized building blocks 199–200
- dizinc cores
 in situ formation 199
 porous properties 215–16
- high-dimensional frameworks 197, 201–2
- synthetic strategy 198–200
- paddlewheel dinuclear tetracarboxylate
 building blocks 195–217
 advantageous characteristics 195–6, 196
 coordination mode of axial positions 200
 electronic structures of dinuclear cores 197–8
 in situ 198–9, 199
 porous properties based on 213–16
 preorganized 199, 199
 conductivity of complexes with mixed-valance rhodium cores 206–7
 crystal structures based on 201–6
 magnetism based on paramagnetic ruthenium cores 208–12
 porous properties based on 212, 213
 preparation of 199–200
- palladium-catalyzed reactions 9
- [2.2]-paracyclophane 57
 target-oriented synthesis 52, 56–8, 59
- parallel interpenetration 160
- pen structures 97–9
 see also molecular pen structures
- pentagonal bipyramidal geometry 118
- pentagonal dodecahedron, $\text{Ce}_2(\text{H}_2\text{O})_{18}$ cage 271, 273
- periodical coordination chemistry 1
 compared with convergent/molecular coordination chemistry 1, 2
 knots and spacers 3
- pharmaceutical co-crystals 25, 27
 advantages 46
 case study (carbamazepine) 36–45
 ‘fake’ 45–6
 meaning of term 25–6
- 1,10-phenanthroline ligands 230–1, 239
- 4,7-phenanthroline ligands 242
- phenyl nitronyl nitroxide radical/
 phenylboronic acid polymorphic co-crystals 32, 34, 34
- [2+2] photodimerization, template-controlled 53, 54
- pi–halogen dimer (PHD) interactions 101–6
 typical motif 102
- pi–pi (π – π) interactions
 0-D metal complexes 237–9
 1-D coordination polymers 239–45
 2-D coordination polymers 245–8
 strength of interactions 220
- 2-picolinic acid 118
 in diorganotin carboxylates 118, 119, 120
- pillar ligands 213–14
- piperazine-2,5-dione/2,5-dihydroxy-*p*-benzoquinone polymorphic co-crystals 31–2, 33
- cis*-platinum complexes 9
- platinum(II) square-planar complex 223, 226
- α -Po networks 175, 176, 178, 255
- polycyclic aromatic hydrocarbons, interactions with, dirhodium(II) tetra(trifluoroacetate) 204–5
- polyimidazole ligands, in metal complexes 225–6, 228
- polymorphism 28, 68
 co-crystals 28–35
 gold(I) chain complexes 256, 257
- polyrotaxane-like structure 315–16, 316
- porous materials 212–16, 248, 299–300
 see also microporous structures
- porous properties
 paddlewheel complexes 212–16
 with *in situ* building blocks 213–16
 with preorganized building blocks 212
- porphyrin complexes 223–4, 227
- preorganized building blocks for paddlewheel dinuclear tetracarboxylate complexes 199, 199
 architectures based on 201–6
 conductivity based on 206–7
 magnetism based on 207–12
 porous properties based on 212, 213
 preparation of 199–200
- pseudo-polymorphs *see* solvates
- Pt_4O_4 network 176, 181
- pyridine ferrocenyl derivatives, solid-state synthesis 9, 10
- pyrimidine ferrocenyl derivatives, solid-state synthesis 9, 10
- 2-quinaldic acid 118
 in diorganotin carboxylates 118, 119, 120
- quinhydrone 29
 see also hydroquinone/*p*-benzoquinone polymorphic co-crystals

- 'railroad' polymers 178, 187
resorcinol and derivatives, as linear templates 52, 54–6, 57–60
retrosynthetic analysis 51
 examples 57, 59
rhodium complexes *see* paddlewheel dinuclear complexes
rhombohedral networks 224, 227
rings-and-rods interpenetration 162
rotaxane-like interactions, in interpenetrating networks 162, 166
rotaxane-like structures (beads on thread) 187
rotaxenes, interpenetrating networks as polymeric analogs 178, 183
'ruthenium blue solution', tetracarboxylate complexes formed from 200
ruthenium complexes 85–7
 see also paddlewheel dinuclear complexes
rutile topology 183, 184
- S-shaped chains, in 3-D porous networks 242, 243
saccharates 275, 277–81
 disadvantages as ligands 275, 277
 positive aspects of use 277–8
 see also zinc saccharate
saddle-like channels 233
Schlenk techniques 5
second-sphere interactions 69–77
seeding 4, 6
self-assembly 53, 69, 101, 106, 236
self-entanglement net 249, 251
self-penetrating networks 183–5
shortest-circuit approach (analysis of networks) 183–5
silver acetate/1,4-diazabicyclo[2.2.2]octane co-crystals 10–11, 12
silver(I) complexes
 argentophilic interactions 249–55
 intermolecular aromatic interactions 243, 244
silver(I) coordination polymers
 with tris(isonicotinyl)-substituted CTV cavitands 151–2
 with tris(2-pyridylmethyl)-substituted CTV cavitands 150–1
silver(I) cyanide anion, coordination polymers with 255
sixteen-membered rings 123, 125
S_N2 reactions, template-controlled 53, 54
¹¹⁹Sn NMR spectrometry 78, 120, 122
sodalite network 283, 285–6
 metal-carbonate complexes 287–8, 287, 288
 body-centered cubic structures 290–1
 face-centered cubic structures 291, 293–4
'soft' coordination environments 77–84
 effect of disorder 84
 solid-state structure compared with gas-phase calculations 81–3
solid-state supramolecular transformations, coordination polymeric structures 298–306
solvates 35
 see also hydrates
solvent-drop grinding 27
 see also kneading
solvent-free reactions 3–19
 advantages 3
 gas–solid reactions 3, 13–17, 202–3
 solid–solid reactions 3, 6–13, 17–18
spacer ligands 3, 67, 68, 214, 221, 223
spherical hexamers, dinitro diquinoline inclusion hosts 108–10
spin-crossover polymers 255
spiral arrays, silver(I) complexes 255
square grid networks 214, 221, 222, 223, 226
square planar coordination environment 128, 289
square pyramidal geometry 68, 244, 298
staircase helical coordination polymer 313–16
staircase inclusion compounds 101–5
star-like channels 312, 313
steric control 37, 77–81, 87, 117–33
 binary bismuth xanthates 130–2, 133
 binary mercury dithiocarbamates 128–30
 binary zinc xanthates 123–5
 bipyridine adducts of zinc dithiophosphates 125–8
 diorganotin carboxylates 118–20
 triorganotin carboxylates 77–9, 120–3
strong-acid hydrogen-bonded 18-crown-6 complex 75, 76, 77
strontium aqua complex ion, in CTV-based structures 140–1, 141
sulfathiazole, solvates 35
sulfonamides, co-crystals, with aromatic carboxylic acids 7
4,4'-sulfonyldiphenol/pyrazine co-crystal 162, 163
supramolecular chemistry, advantages 52
supramolecular isomerism 310–12

- supramolecular synthons 27, 52, 67
 carboxylic acids as 222–4
 listed for carbamazepine 36
 and polymorphism 34
supramolecular targets 52
Suzuki reactions, mechanically induced 9
syntactic structural chemistry 77
- target-oriented organic synthesis 51–2
 and covalent capture 52
 in solid state 56–64
 ladderanes 59–60
 [2.2]-paracyclophane 56–8
 see also template-controlled synthesis
- tectons 67–8, 117, 136
- template-controlled reactivity
 generality of method 56
 in liquid phase 53, 54
 modularity of method 56
 in solid phase 54
- template-controlled synthesis 53–4
 1,2,4,5-benzenetetracarboxylic acid as linear
 template 62–3
 bis-*p*-phenylene[34]-crown[10] as linear
 template 62
 1,8-naphthalenedicarboxylic acid as linear
 template 61
 resorcinol and derivatives as linear templates
 54–60
 tetrakis(4-iodoperfluorophenyl)erythritol as
 linear template 63–4
 tricarballic acid as linear template 62–3
- template switching 52, 58, 59–60
- tetracyanoethene, as linker in 2-D sheet
 structure 201
- tetrahedral geometry, tin complexes 78
- tetrahedral nodes, networks constructed from
 173–5, 266, 269
- tetrakis(4-iodoperfluorophenyl)erythritol, as
 linear template 63–4
- rcit*-tetrakis(4-pyridyl)cyclobutane (4,4'-tpcb),
 54, 55
- tetrakis(4-pyridyl)-1,2,9,10-diethano[2.2]-
 paracyclophane 57
- thermogravimetric analysis
 zinc *N*-(2-hydroxybenzyl)-*L*-alanine
 compounds 302
 zinc saccharate 279
- thia-bridged diquinoline tetrabromide,
 edge–edge halogen interactions
 99–101
- thioether-1,3-*peri* aromatic hydrogen
 interactions 96
- three-connected networks 159, 172, 266
- tin compounds 77–9, 81–4
 Me₃SnOH 83, 84
 see also diorganotin dichlorides; diphenyl tin
 bis(xanthate) complex; triorganotin
 carboxylates
- trapezoidal bipyramidal geometry 118
- triangular channels, in 3-D network 221
- triazine, coordinated to diruthenium
 paddlewheel units 211
- tricapped trigonal prismatic geometry,
 lanthanide complexes 272
- tricarballic acid, as linear template 62–3
- trifluoroethanol, in CTV-based structures 144,
 144, 145, 146, 147, 148
- trigonal bipyramidal geometry 68, 122
- trigonal modes, networks constructed from
 159, 172, 266
- α -trimesic acid, 2-D \rightarrow 3-D parallel
 interpenetration structure 166–7, 169
- γ -trimesic acid, 2-D inclined interpenetration
 structure 167–8, 170
- triorganotin carboxylates 77–9, 120–3
- triphenylene 204
 interactions with dirhodium(II)
 tetra(trifluoroacetate) 204
- triphenylsilanol/4,4'-bipyridine polymorphic
 co-crystals 31, 32
- triple-decker stacks 63
- triple-strand helicates 242
- triply interpenetrated networks
 2-D nets 221–2, 224, 251, 253
 3-D nets 255, 256
- tris(2-benzimidazolylmethyl)amine, in metal
 complexes 226, 228
- tryptamine/hydrocinnamic acid chiral
 co-crystals 6
- uranyl crown ether complexes 74–5, 76
- water
 cyclic tetramers 316
 as H-bond donor, in CTV-based structures
 139, 140, 140
- woven cloth-like layers 186
- xanthate anion 123
 see also bismuth xanthates; zinc xanthates



- zigzag polymeric chain
 paddlewheel dirhodium tetracarboxylate
 complexes 202
 straightening by steric control 126–7,
 127–8
- zinc dithiophosphates, bipyridine adducts
125–8
- zinc enzymes 225
- zinc *N*-(2-hydroxybenzyl)-alanine dimer
298–9
 chloro-/methyl-substituted, reversible
 dehydration 302–3
 hydrogen-bonded 3-D network 299
 thermal dehydration 299–301
- zinc saccharate 278–81
 absorption of guest molecules
 280–1
 dehydration/hydration study 280
 extended 3-D structure 278, 280
 preparation of 278
 stability 279–80
 thermogravimetric analysis of hydrated
 crystals 279
 ‘wall’ structure 278, 279
- zinc–vanadium phosphate complex 245
- zinc xanthates, binary 123–5
- zipper-like structures 231, 232, 239,
241, 245

With kind thanks to Paul Nash for creation of this index.