

Index

- A-B-C model of negative thinking 131
- abstinence from alcohol 21, 211
 - cognitive dysfunction 31, 67
 - cue exposure 231
 - pre-moderation 68, 159, 305
 - short-term targets 64, 69
- abstinence from drug use 69–70
 - see also* drug-free lifestyle goal
- abstinence-violation effect 228
- abstinence vs. moderation goals 66–69, 155, 156, 159, 162, 301, 304–305
 - see also* goal setting
- acamprosate 193, 200–201
- action plan 96, 98, 100, 225, 229, 239, 249
- active listening 125
- addiction *see* dependence
- adolescence 254, 265
 - see also* young people, working with
- advice to cut down 80, 84, 292
- advocacy 250
- affirming, counselling skill 9
- aggressive behaviour
 - vs. assertiveness 108, 111–114
 - responding to client's 171, 283
- agoraphobia 275
- alcohol
 - assessment of current use 21–22, 82–83, 257–258
 - damage to physical health 29, 66–67
 - dual diagnosis 270, 275, 276
 - effect on cognitive function 31–32, 67, 278
 - level of dependence 22–25
 - levels of risk in drinking 81, 94
 - pharmacotherapies for treating
 - dependence 198–202
 - treatment goals 66–69
 - withdrawal from 23
 - alcohol-free days 159, 164
 - see also* drinking guidelines
 - alcohol-related brain injury *see* cognitive dysfunction; memory loss
 - Alcoholics Anonymous 14, 68, 130, 293
 - twelve steps of 218
 - see also* self-help groups
 - ambivalence 46
 - about taking medication 196–197, 280
 - brief and early interventions 86
 - motivational interviewing 45, 46, 47, 53–54, 58, 304, 307
 - reflecting 7–8, 9
 - stages of change 32
 - amphetamines 22, 30, 276
 - see also* psychostimulants, withdrawal from
 - anger 116, 161, 226
 - see also* assertiveness
 - Antabuse *see* disulfiram
 - anxiety
 - and substance use 222
 - screening for 30, 258–259, 274–276, 278
 - test-related 30
 - treatment of 31, 129, 131, 144, 145, 226, 271, 279, 282–283, 288, 308, 310
 - withdrawal symptom 23
 - see also* post-traumatic stress disorder; relaxation training; social phobia

- apparently irrelevant decisions 227
- assertiveness 108
- assertiveness training 107–119, 161, 305
- goals 107
 - key concepts 108
 - recommended use 107
 - rights 108, 111, 119
- assessment 19–43
- brain injury and cognitive impairment 30–31
 - dependence 22–25, 85–86
 - estimating levels of consumption 21–22, 83, 257–258
 - involving concerned others 26, 172–173, 181, 230, 260–261
 - lifestyle and social stability 25–28
 - mental health disorders 30, 258–259, 271–278
 - motivation 20–21, 256–257
 - needs-based 247
 - pattern of drinking or drug use 21–22, 257–258
 - physical health problems 29–30, 82–83, 84, 85
 - risk-taking behaviour 27–28, 257
 - sexual abuse 27, 258–259, 275–276
 - sexual problems 27, 176, 278
 - social support 26, 230, 259–261
 - stages of change 31–33, 85
 - standardised questionnaires 20, 83, 224, 273
 - strengths-based 19, 25, 248–249
- assessment tools
- Addiction Severity Index (ASI) 25
 - Adolescent Relapse Coping Questionnaire 264
 - Alcohol Problems Questionnaire (APQ) 25
 - Alcohol Use Disorders Identification Test (AUDIT) 83–86, 92
 - Brief Drinker Profile (BDP) 20, 25
 - Brief Symptom Inventory (BSI) 273
 - Comprehensive Drinker Profile (CDP) 20
 - Drug Abuse Screening Test (DAST) 83, 93
 - General Health Questionnaire (GHQ) 273
 - Kessler 10 Symptom Scale (K10) 273
 - Maudsley Addiction Profile (MAP) 20
 - Mini-Mental State Examination 31
 - One-Week Retrospective Diary 83
 - Opiate Treatment Index (OTI) 20, 25
 - Severity of Alcohol Dependence Questionnaire (SADQ-C) 24, 56, 67, 38–39
 - Severity of Dependence Scale (SDS) 24, 43
 - Severity of Opiate Dependence Questionnaire (SODQ) 24, 41–42
 - Short form Alcohol Dependence Data Questionnaire (SADD) 24, 40, 67
 - Situational Confidence Questionnaire 224
 - Symptom Check List (SCL-90-R) 273
 - Time Followback Method (TLFB) 22
- attention-deficit/hyperactivity disorder (ADHD) 258
- automatic thoughts 130, 131–133
- behavioural self-management 71, 85, 155–168, 293, 305
- drinking limits 63–64, 158–160
 - goals 156
 - key concepts 156–157
 - rationale 157
 - recommended use 155–156
- beliefs 110, 132–133
- see also* automatic thoughts
- benzodiazepines 23, 24, 30, 83, 84, 279, 288, 309–310
- see also* pills, risk taking
- binge drinking 21, 158–159, 299–300
- see also* drinking guidelines
- bipolar disorder 273–274
- blackouts 31, 262
- blood alcohol concentration (BAC) 159–160, 167, 168
- boundaries 13, 262, 276, 283–284
- brainstorming 97–98
- examples of application 10, 57, 96, 100–102, 105, 132, 174, 177, 225, 228, 239, 265
- brief and early intervention 79–94, 295, 310–314
- for drug use 87
 - goals 80
 - in drug and alcohol settings 86–87, 292
 - in primary health care settings 84–86, 291–292

- key concepts 80–81
- recommended use 67, 79–80
- screening and detection 82–83
- buprenorphine 193, 202–204
- burnout 246

- Campral *see* acamprosate
- cannabis 22, 23, 24, 28, 30, 79, 83, 258, 276, 282, 310–314
- case conferences 247
- case management 65, 243–252, 281, 283, 293, 294, 310–314
 - assessment 247–249
 - goals 244
 - liaison with concerned others 250–251
 - other services 224–245, 249–250
 - recommended use 243–244
 - the role of case manager 13, 15, 245–246
- change talk 47, 53, 86
- child care 4, 25, 110, 244
- cocaine 22, 282
 - see also* psychostimulants, withdrawal from
- coercion 7, 21, 27, 31, 45, 69, 253, 255, 256–257, 310–314
 - see also* court mandated clients; involuntary admission
- cognitive therapy 71, 129–141, 195–196, 261–262, 282–283
 - goals 129–130
 - key concepts 130
 - rationale 130
 - recommended use 129
- cognitive dysfunction
 - abstinence as preferred option for 67
 - assessment of 30–31
 - treatment for clients with 129, 280, 283, 294
- communication skills 112, 121–127, 179, 182, 187, 230
 - goals 121
 - key concepts 122
 - rationale 122
 - recommended use 121
- compliments 10, 113, 115
- conduct disorder 258, 277

- confidence, building client's 3, 9, 19, 21, 54, 103, 104, 113, 130, 134, 213, 215, 222, 244, 255
 - see also* self-efficacy; self-esteem
- confidentiality 13, 15, 16, 20, 83, 181, 247, 251, 256, 271
- conflict 96, 111, 117, 124, 161, 173, 222
 - helping couples to resolve 170, 172, 173, 175, 178, 215
 - helping families to resolve 181, 183–184, 185–186, 187
 - in group therapy 11
 - see also* time-out
- confrontation, alternatives to 4, 7, 11, 46, 50–51, 263–264, 282
- conversation skills 123–124
- coordinated care 249–250, 281–282, 312–314
- coping skills 26, 222, 224–226, 261–262
- counselling skills 3–18, 47–48
 - goals 3
 - ideal therapist 4
- couples therapy 171–178, 302
 - assessment for 172–173
 - goals 171–172
 - responsibility for substance use 172, 177, 178
 - supporting medication 177
 - when partner presents alone 178–179
- court mandated clients 82
- cravings
 - ways of coping with 71, 213, 226
 - during therapy 105–106
 - high-risk situation 26, 222
 - pharmacotherapies to reduce 193, 199, 202, 204
 - self-monitoring 97, 166, 195, 207, 214, 224, 235, 261
 - self-talk as a trigger for 130, 131, 282
 - symptom of dependence 23
- criticism 112–115, 117
- cue exposure 231
- culture 5, 169, 254, 310–314

- day diary 69, 158, 160, 166, 224, 235
- decisional balance sheet (also known as decision matrix) 57, 61, 262
- dependence 23
 - assessment of 22–25, 83, 86, 292

- dependence level as a guide to
 - brief intervention 79–80, 86, 292, 295
 - case management 243
 - goal setting 67, 69–70, 193
 - inpatient/residential care 293
 - intensive treatment 85, 292, 295
 - pharmacotherapies 69–70, 193
 - self-help groups 211
- depression 273–274
 - and substance use 222, 270
 - screening for 30, 258–259, 274, 278
 - treatment of 129, 131, 133, 226, 271, 279, 282–283, 288, 302–303
 - withdrawal symptom 23
 - see also* suicide, risk of
- detoxification
 - dependence 22–23
 - intervention 64, 85, 247, 292, 293, 296
- developmental delay 258
- disease model 212, 215
- disulfiram 193, 201–202
- domestic or family violence 10, 25–26, 64, 169, 171, 173, 175, 179, 259, 260
- drink and drug refusal skills 26, 71, 103–106, 160, 225–226, 257, 262, 263–264, 293, 313
 - goals 103
 - rationale 103–104
 - recommended use 103
- drink driving 27, 28, 66, 179, 257
- drinking guidelines
 - BAC 167, 168
 - levels of risk 81, 82, 94
 - limit setting 85, 158–160, 173, 292, 293
 - sensible 80–81, 84, 155, 156, 158, 164, 291
- drug-free lifestyle goal 69, 212
- drug use
 - abstinence from 69
 - assessment of 22, 82–83, 257–258
 - cognitive dysfunction 30–31
 - dependence 22–25
 - health effects 29–30, 87, 193
 - see also* risk-taking behaviour
- DSM-IV 23–24
- dual diagnosis 258–259, 269–288
 - adjusted drug and alcohol counselling 282–284
 - basic guidelines 269
 - goals for working with 270
 - influence on goal setting 64, 67
 - key concepts 270
 - management flowchart 270–271, 272
 - screening for 27, 28, 30, 65, 258–259, 271–278, 293, 296
 - types of mental health treatment 278–282, 288
- dual relationships 14–16, 181, 245–246
- dysthymia 273, 301, 301, 302–303
 - see also* depression
- eating disorders 258, 278
- ecstasy *see* MDMA
- education 264–265
- elaborating, counselling skill 8
- emotions
 - dealing with 108–109, 132–133, 133–135, 161, 226
 - as a high-risk situation 109, 222, 228
- empathy 4, 5, 11, 19, 50, 56, 58, 87, 126, 174, 178, 246, 282, 291
 - defined as a skill 6–7, 46
 - defined as a value 5
- engaging your clients 3, 4–5, 19–20, 181, 255–256, 261, 282, 283
- ethical dilemmas 16
- ethical practice 12–16
- expectancies 255, 262–263
- extended care 4–5, 12, 32, 162, 243, 250, 237–239, 292, 293, 294
 - for young people 265
 - in response to lapses or relapses 230–231, 238–239
 - recommended use 237
 - re-entry phase, after inpatient care 294, 310
- family meetings 179–188, 312
 - goals 180
 - rationale 182
 - when the young person does not seek help 188
- family or concerned others
 - assessment of 25–26, 172–173, 181, 259–261, 291
 - case management involving 244, 247, 250–251

- involvement in goal setting 68, 169–170, 172, 182–183
- involvement in therapy 26, 169–192, 195, 201–202, 229, 230, 253, 260–261
- own drinking or drug problems 172–173, 179, 260, 306
- self-help groups 215–216, 283
- specialist therapy for 26, 169, 179
- see also* couples therapy; domestic or family violence; family meetings
- fire drill analogy *see* relapse drill
- foetal alcohol syndrome 29
- follow-up
 - after an intervention 16, 80, 84, 85, 246, 292
 - missed appointments 5, 11, 195, 283, 313
 - see also* extended care
- ‘FRAMES’ 58
- gambling 278
- gender xvii, 5, 10, 214
- generalised anxiety disorder 275
- goal setting 19, 21, 29, 31, 32, 63–75, 155
 - for alcohol 66–69
 - for opioids, other drugs 69–71
 - harm reduction 65–66, 70–71
 - lifestyle 64–65
 - negotiating 63–64, 249, 255, 257, 282
 - support of concerned others for 68, 169–170, 172, 182–183
 - see also* abstinence; abstinence vs. moderation; drug-free lifestyle; harm reduction; moderation; pharmacotherapies
- good and less good things 48–51
 - see also* motivational interviewing
- group therapy 10–12, 95, 103, 105–106, 107, 121, 125, 126, 129, 143, 156, 171, 221, 225, 239
- iatrogenic effects 11, 265
- motivational interviewing 57–58
- working with young people 265
- Guidelines for the Treatment of Alcohol Problems xiv, xv
- harm reduction 54–55, 70–71, 103, 155, 193, 214
 - brief and early interventions 82, 87
 - clients with dual diagnosis 282
 - goal setting 65–66, 70–71
 - with young people 253, 257, 268
- hepatitis 28, 29, 69, 70–71, 193, 199
- high-risk situation 222
 - identifying 26, 47, 86, 160, 177, 184–185, 223, 224, 292
 - planning for 87, 97, 160–161, 169–170, 186–187, 224–227
- HIV 28, 29, 56, 64, 69, 70–71, 193, 214
- inhalants 257–258
- injecting drug use
 - assessment of 22, 28, 257
 - risks associated with 28, 29–30, 31, 193
 - safer practices 65–66, 70, 257
 - see also* harm reduction
- inpatient/residential programmes xv, 229–230, 293–294
- interests and hobbies *see* pleasurable activities
- internet, use in therapy 256, 268
- intoxication during therapy 11, 21, 312
- I-statements 122
 - applications 112, 124, 171, 176, 179, 182, 187
- involuntary admission 279
- involving concerned others 169–192
 - basic guidelines 169
 - key concepts 170–171
 - see also* couples therapy; family or concerned others
- lapse 222
 - programme response to 229–230
 - reducing harm 66, 155
 - use in relapse prevention 173, 177, 183, 183–184, 186–187, 223, 229, 228–229
- legal problems 27
- length of treatment xvi–xvii, 291–292, 295–296
 - assertiveness skills training 107
 - behavioural self-management 156
 - brief interventions 79, 84, 86
 - cognitive therapy 129
 - pharmacotherapies 194, 198
 - relaxation training 145

- lifestyle treatment goals 64–65, 223, 230, 293, 293–294
- liver function 29
- MATCH, Project 59, 73, 99, 106, 117, 126–127, 135–136, 189, 215, 216, 217, 232, 291, 297
- MDMA (Ecstasy) 28, 30
- mean corpuscular volume (MCV) 29
- mental health services 272, 278–282, 296
- memory loss 30–31
see also blackouts; cognitive dysfunction
- menu or range of strategies 32, 54, 58, 71–72
- metaphors or stories, used in therapy 256, 261
- methadone maintenance 193, 194, 202, 203–204, 211, 307–308
- methamphetamine 258
see also psychostimulants
- moderation 67, 69, 80, 155, 165, 231
see also abstinence vs. moderation goals; behavioural self-management; brief and early interventions; goal setting
- Moderation Management, 211
- mood disorders 273–274
see also bipolar disorder; depression; dysthymia
- motivational interviewing 45–62
 counselling applications of 4, 21, 32, 63, 80, 86, 151, 152, 238, 249, 251, 282, 292, 292–293
 counselling with pharmacotherapies 195–197, 203
 counselling with young people 257, 259, 262
 goals 45–46
 key concepts 46–47
 recommended use 45
see also stages of change
- naltrexone 193, 198–200, 301–302
- Narcotics Anonymous 310
 twelve steps of 219
see also self-help groups
- negative thinking
 becoming aware of 131
 challenging 132–133, 139
- non-assertiveness 108, 109, 110–111
see also aggressive behaviour; passive under-assertion
- non-residential programmes *see* outpatient programmes
- non-verbal communication 104, 112, 116, 124, 125, 176
- notification 16, 20, 27, 256, 259, 272
- obsessive-compulsive disorder 275
- open-ended questions 8
 applications 4, 22, 47, 48–50, 123, 185, 197, 255, 260–261, 262, 263
- opioids 194
 assessment 20, 22
 brief intervention for 82–83, 87
 goal setting 69–71
 pharmacotherapies for 193–194, 202–204
 withdrawal from 23
- outpatient programmes xv, 292–293
- outreach 243, 245, 246
- overdose 27, 31, 70, 200, 202, 203, 278, 306, 308
- panic disorder 275
- partner or spouse *see* couples therapy; family or concerned others
- passive under-assertion 108, 110–111, 111–114
see also assertiveness training
- peer group 253, 255, 254, 259, 263–264, 265
see also social networks; young people
- personalising health effects 55–57
 assessment 19, 24–25, 29
 clinical applications 85, 86, 87, 157, 196, 292, 292–293
 with results in the normal range 57
- personality disorders 271, 276–277, 283
 antisocial personality disorder (ASPD) 277
 borderline personality disorder (BPD) 277–278
- pharmacotherapies 29, 71, 156, 193–208, 293, 296
 as a harm reduction strategy 66, 69–70, 193, 257
 goals 194

- helping your client stay on medication 177, 195–197
 - for treating alcohol dependence 197–202
 - for treating opioid dependence 202–204
 - key concepts 194
 - monitoring the medication 195, 207
 - rationale 195
 - recommended use 193
- pills, risk-taking 27, 28
- pleasurable activities 27, 123, 161–162, 230, 263, 264
- polydrug use 22, 69, 257, 257–258, 282
- positive interactions, promoting 170, 175–177, 192
- positive specific requests 122, 125, 175, 182
- post-traumatic stress disorder 258–259, 275–276, 309
- pregnancy 29, 80, 83
- primary health care settings 32, 45, 81, 84–86, 291–292
- problem-solving skills training 95–102
 - applications of 135, 151, 161, 175, 178, 185, 202, 224–225, 227, 228, 264, 283, 293
 - goals 95
 - rationale 95–96
 - recommended use 95
- protective factors 254–255, 259–265
- providing information 54–55
- psychiatric medication 279–280, 288
- psychoses 258, 271, 276, 283
- psychostimulants, withdrawal from 23

- Quality Assurance Project xiii–xiv, xv

- readiness for change
 - and resistance 7, 50–51
 - assessment of 85, 292
 - signs of 53
 - treatment responses to 3, 31, 45, 47–48, 80, 87, 253, 257, 281–282, 282
 - see also* stages of change
- referrals, making 85–86, 245, 271–272, 278–279
- reflective listening, counselling skill 6–7
 - applications 13–14, 47, 53, 56, 58, 125, 134, 182, 196–197, 248–249, 260–261
- reframing, counselling skill 8–9
 - in motivational interviewing 46, 51, 57, 58
 - involving concerned others 170–171, 174, 182–183
- relapse
 - beliefs about 196, 228
 - definition 222
 - part of a learning process 33, 183, 221, 229, 283
 - responding to 229–230, 230–231, 237, 238, 283, 293
 - triggers for 22, 96, 108, 121, 144, 269, 270, 276, 282
 - see also* relapse prevention training
- relapse prevention training 221–236
 - applications 4, 32, 72, 86, 103, 122, 133, 134, 174, 193, 198, 203, 204, 261, 263–264, 283, 292, 293, 296
 - assessment for 21, 26, 47, 256, 296
 - goals 221–222
 - group applications 10, 12, 211, 214, 221
 - harm-reduction and 66, 70, 155, 221
 - key concepts 222
 - recommended use 221
 - with concerned others 170, 172, 173, 174, 177–178, 180, 183, 186, 186–187, 188
 - with moderation goals 160–162
 - see also* high-risk situation; lapse; relapse
- relapse drill 177, 183, 228–229
- relaxation response 144
- relaxation training 143–154
 - applications 10, 71, 134, 161, 197, 204, 293
 - goals 143
 - key concepts 144
 - isometric 146–147, 150–151
 - meditative 146, 149–150
 - progressive muscle 146, 148–149
 - rationale 144–145
 - recommended use 143
- requests, dealing with 116
- residential programmes *see* inpatient/residential programmes
- resistance 4, 7, 46–47, 53, 186
 - rolling with 7–8, 50–51, 57–58, 86
 - see also* counselling skills; motivational interviewing
- resources list
 - assertiveness skills 117–118
 - assessment 34–37
 - behavioural self-management 162–163

- resources list (*cont.*)
- brief and early interventions 88–91
 - case management 251–252
 - cognitive therapy 135–136
 - communication skills training 126–127
 - drink and drug refusal skills 106
 - dual diagnosis 285–287
 - extended care 239
 - general counselling skills 16–18
 - goal setting 72–74
 - how to use this book xvii
 - involving concerned others 188–191
 - motivational interviewing 58–60
 - pharmacotherapies 206
 - problem solving skills 99
 - putting it all together 297–298
 - relapse-prevention training 231–234
 - relaxation training 151–152
 - self-help groups 216–217
 - working with young people 266–267
- Revia *see* naltrexone
- risk factors 254–255, 259–265
- risk-taking behaviour
- assessment of 22, 27–28, 82, 257
 - goal setting 64, 65, 70–71, 257
 - intervention for 54–55, 55–56, 79, 87, 193, 203
 - see also* harm reduction
- risk in drinking alcohol, levels of 94
- see also* drinking guidelines
- role-play 12
- assertiveness training 107, 111–112, 114, 117
 - communication skills training 121, 123, 124, 125
 - couples therapy 174, 176, 177
 - drink and drug refusal skills 103, 104, 105, 160, 225–226
 - problem solving 98–99, 100, 225
 - relapse prevention training 229
 - working with young people 256, 265
- rolling with resistance, counselling skill 7–8
- safe sex 71
- sensible drinking guidelines 164
- see also* drinking guidelines
- schizophrenia 276
- see also* psychoses
- screening
- for dual diagnosis 258–259, 271–278
 - for drug problems 83
 - for risky drinking 83
 - see also* brief and early interventions
- seizures, in response to withdrawal 23
- self-disclosure
- by counsellor to client 13, 14
 - by young person to family 186
- self-efficacy 47, 87, 224
- supporting 54, 58, 87, 248, 255, 282–283
- self-esteem 7, 46, 64, 52–53, 71, 109, 129
- see also* confidence, building client's
- self-help resources 85, 157
- assertiveness 118
 - behavioural self-management 90–91, 142
 - brief interventions 90–91, 142
 - families 191
 - harm reduction 73–74, 90–91
 - sexual intimacy 190
 - young people 68
- self-help groups 209–219
- goals 212
 - internet sites 217, 287
 - key concepts 212–213
 - linking with therapy 214–215
 - rationale 213–214
 - recommended use 211–212
 - social support and 68, 71, 211, 214, 229, 230, 283, 293
 - to assist family members 178, 215–216
 - see also* Alcoholics Anonymous; Moderation Management; Narcotics Anonymous; Smart Recovery
- self-monitoring
- behavioural self-management 156–157, 158, 159, 160, 162, 166
 - high-risk situations 160, 224
 - use of medication 195, 207
 - tension 146, 153
 - thoughts 131, 133, 135, 140–141
 - urges and cravings 224, 235
- self-motivational statements *see* change talk
- self-reward 155, 161
- self-talk 130, 131, 174, 177
- see also* automatic thoughts; beliefs

- sex differences in alcohol guidelines 158–159, 160, 164
- sexual abuse 10, 16, 27, 254–255, 258–259, 275–276, 309, 310
- sexual problems 27, 176, 278
- shared rewarding activities 170, 172, 175–176, 187
- side effects
 - helping your client to cope with 196–197, 280
 - ofacamprostate 200
 - of methadone and buprenorphine 203
 - of naltrexone 199
 - of psychiatric medications 288
- situational confidence questionnaires 26, 224, 233–234
- sleep disturbance 23, 273, 273–274, 275, 278
- Smart Recovery 211
 - see also* self-help groups
- splitting 13–14, 284
- social atom 260
 - applications 26, 230, 260–261, 263
- social networks
 - assessment of 25–26, 230, 259–261, 291
 - building 198, 211, 213–214, 214, 225–226, 230, 253, 256, 261, 263, 264, 283, 293, 313
 - influence on goal setting 68
 - support from other clients, 10, 12, 237–238, 239
- social phobia 275, 307, 308
- social pressure
 - high-risk situation 222, 265
 - interventions 103, 114, 160, 225–226, 263–264
 - see also* drink and drug refusal skills
- solution focused questions 9, 184, 248–249, 282–283
- sponsors, self-help groups 213, 214
- stages of change 31–33, 48, 84, 87
 - action 32
 - contemplative 32, 45, 54, 80
 - maintenance 32
 - precontemplative 31–32, 45, 50–51, 54–55, 80, 85, 28, 312
 - preparation 32
 - relapse 32, 33
 - see also* readiness for change
- standard drinks 80–81, 164
- estimating 21–22, 83, 166
- setting limits 63, 94, 156, 158, 158–159, 164, 167, 168
 - see also* drinking guidelines
- stress management *see* relaxation training
- substitution therapy 194
 - see also* pharmacotherapies, for treating opioid dependence
- Subutex *see* buprenorphine
- suicide, risk of 30, 259, 272, 274, 277, 279
 - see also* notification
- summarising, counselling skill 9
 - applications 47, 48, 50, 53, 53–54, 57, 57–58, 180, 182, 262
- supervision, professional 12, 14, 16, 246, 272, 274, 277–278
- tension 144, 145–146, 153
- therapeutic community xv, 69, 230
- thinking errors 131, 132, 137–139
 - see also* automatic thoughts; beliefs
- time-out
 - from drinking 68, 159
 - from interpersonal conflict 170, 174–175, 178, 184, 187
- tolerance 23, 57
- twelve
 - steps 212, 218, 219
 - traditions 212
 - see also* self-help groups
- validating
 - client's feelings 134
 - communication skill 125–126, 175, 182
 - vocation 25, 230, 264–265
- withdrawal symptoms
 - assessment 18
 - goal setting 67, 292, 293, 296
 - pharmacotherapies and 197, 199, 202, 202–203, 204
 - to distinguish from dual diagnosis 272, 275
- workplace interventions 81

X-Y-Z statements 124

young people, working with 253–268
 assessment 256–259
 basic guidelines 253

goals 253–254

key concepts 254–255

reducing risk and enhancing protection
 259–265

see also family meeting; family or
 concerned others