

---

# Index

---

- $\alpha$ -methylene- $\gamma$ -butyrolactones 190–2
- 1-acetonaphthone 100–2
- acetoxypalladation of alkynes 185–90
- 3-acetylbenzonitrile 102–3
- 4-acetylbiphenyl 134–6
- alcohols
  - diaryl methanols 161–3
  - homoallylic 156–61, 164, 166–8
  - stereoselective hydroformylation of allylic alcohols 226, 229–30
- aldehydes, regioselective and asymmetric 1,2-addition reactions
  - with 3-chloro-propenyl pivolate using Jacobsen's salen 164–8
  - with boronic acids using chiral ferrocenyl ligand 161–3
  - metal-mediated reaction with allyl bromides 156–61
- alders-ene synthesis of butyrolactones and cyclopentanones 190–2
- aldol reactions, asymmetric
  - $\alpha$ -hydroxyketones to *syn*-1,2-dihydroxyketones 202–8
  - catalysis by (*S,S,S*)-pyrrolidine-2-carboxylic acid (2'-hydroxyl-1',2'-diphenyl-ethyl)-amine 208–10
- syn*-alk-1-ene-3,4-diols 164–8
- alkenylation, of aromatic ketones using  $\text{RuH}_2(\text{CO})(\text{PPh}_3)_3$  14, 17–18
- alkylation of
  - allylic acetates using ferrocenyl-ligands with Pd 51, 55–6
  - allylic acetates using fluoros bisoxazoline with Pd 40–1, 45–6
  - amino acid derivatives using copper(II)chsalen 21, 24–7
  - aromatic ketones using  $\text{RuH}_2(\text{CO})(\text{PPh}_3)_3$  14, 16–17, 19–20
  - glycine imines using alkaloid derived quaternary ammonium salts 27, 31–3
- allylation of aldehydes 156–61
- allylic alcohols
  - homoallylic 156–61, 164, 166–8
  - stereoselective hydroformylation of 226, 229–30
- allylic substitutions
  - acetate group by amino derivatives 50
  - acetate group by benzylamine 36, 37–40
  - acetate group by malonate 36, 37–40, 50, 51, 55–6
  - using axially chiral diphosphine complexes 50
  - using chiral ferrocenyl-imino phosphines 51, 55–6
- amination, aryl
  - using *N*-heterocyclic carbene ligand 86, 88–9
  - using nickel-on-carbon catalyst 141–2
- $\alpha,\alpha$ -amino acids, asymmetric synthesis of 21, 24–7
- (1*S*,2*S*,4*S*,5*R*,1'*R*)-1-(anthracen-9-ylmethyl)-5-ethyl-2-[benzyloxy(quinolin-4-yl)methyl]-1-azoniabicyclo[2.2.2]-octane bromide 28–9, 33
- (1*S*,2*S*,4*S*,5*R*,1'*R*)-1-(anthracen-9-ylmethyl)-5-ethyl-2-[hydroxy(quinolin-4-yl)methyl]-1-azoniabicyclo[2.2.2]-octane bromide 28–9, 33
- aryl boronic acids, coupling reactions of 64, 65–7, 74–7, 147–9
- aryl boronic esters 151–2
- aryl halides
  - carbonylation 244, 245–7
  - in cross-coupling reactions 145–6
  - in Heck coupling reactions 100–4
  - reduction 144–5
  - Suzuki coupling reactions 81–2, 83–6

- 2-aryl oxazoline catalysed Suzuki couplings 74–7
- 2-aryl-2-methyl cyclopentanones 97–9
- arylation  
 of aromatic ketones using  $\text{RuH}_2(\text{CO})(\text{PPh}_3)_3$  18–19, 20  
 asymmetric phenylation of 2,3-dihydrofuran 110–12  
 Heck coupling of vinyl ethers 92–3, 95–9, 100–4  
 of ketones using *N*-heterocyclic carbene ligand 86–7
- 4-arylcoumarins 65–6, 67
- $\beta$ -benzo[*b*]thienyldehydroamino acid derivatives 60–4
- Bacillus megaterium* and carboxylation of pyrrole 247–50
- $\alpha$ -benzyl-alanine methyl ester 24–5
- benzyl(1,3-diphenylprop-2-enyl)amine 39
- (4-benzylidene-5-oxo-tetrahydrofuran-3-yl)-acetaldehyde 190–1
- biarylphosphine oxides 77–81
- BINAP and Intellectual Property Rights 6
- BINOL, (*S,S*-)linked 202, 211
- biphasic catalysis, non-polar 114–16
- 1-biphenyl-2-yl-2,2-dimethylpropan-1-one 18–19
- 1-biphenyl-4-yl-ethanone 115–16
- (2*R*,4*R*)-2,4-bis[(4*R*,6*R*)-4,6-dimethyl-1,3,2-dioxaphosphorinane-2-yloxy]-pentane 241–3
- 1,2-bis(dimethylaminophosphanyl)benzene 234–5
- o*-bis(dimethylamino)phosphinobromobenzene 233–4
- 9-borobicyclo[3.3.1]nonane 67–9
- 3-bromo-4-methylstyrene 120
- 3-bromo-4-methylacetophenone 117–18
- 1-(3-bromo-4-methylphenyl)-ethanol 118–20
- 4-bromo-hexylbenzene 130–1
- 1,3-butadiene as additive in Grignard reactions 128–33
- carbene ligands 86–90
- carbon dioxide, supercritical reaction system 248–9
- carbonylation of aryl halides using dimeric oximepalladacycle catalyst 244–7
- carbonyldihydrotris(triphenylphosphine)-ruthenium(II) 14–16
- carboxylation of pyrrole to pyrrole-2-carboxylate 247–8, 249–50
- catalysts and catalytic processes, overview of  
 fabrication costs 3–6  
 general considerations 3  
 industrial examples 9–12  
 pre-reaction/immobilisation 7–8  
 separation costs 3–6  
 separation techniques 8–9
- 3-chloro-propenyl pivolate 165–6
- chromium-catalysed 1,2-addition to aldehydes 164–8
- chsalen, synthesis of 22–3
- cinchona alkaloid derived quaternary ammonium salts 27, 31–3
- cobalt catalysed reactions 182–5
- coenzyme  $\text{Q}_{10}$  precursor 142
- copper-salen catalyst *see* copper(II)(chsalen)
- copper(II)(chsalen)  
 and alkylation of amino acid derivatives 21, 24–7  
 synthesis 23–4
- cross-coupling reactions  
 2-methylboronic acid with 2-(1-hexynyl)pyridine 148–9  
 alkyl halides with Grignard reagents in presence of 1,3-butadiene 128–33  
 aryl boron reagents with Rh catalysts in aqueous media 147–53  
 intramolecular cyclisation of norbornylene with an arylboronate ester 151–2  
 triorganoindium compounds with aryl and vinyl electrophiles 133–7  
 use of heterogeneous nickel-on-charcoal catalyst 138–47  
 vinyl zirconocenes and aryl halides 145–6
- cyclisation reactions  
 alder-ene synthesis of butyrolactones and cyclopentanones 190–2  
 insertion reactions of alkynes with aldehydes, ketones or nitriles 185–90  
 Pauson-Khand synthesis of cyclopentanones using molecular sieves 182–5  
 titanocene-catalysed cyclisation of epoxygermacrolides 196–200  
 [2 + 2 + 2] trimerisation of alkynes using water soluble Rh catalyst 193–7  
 (2*R*,3*S*)-3-cyclohexyl-2,3-dihydroxy-1-(2-methoxyphenyl)-1-propanone 205–7  
 1-cyclohexylpent-3-en-ol 156–60
- cyclopentanones  
 2-cyclopentanone and Michael reactions 219–22

- intramolecular alder-ene reaction 190, 191–2  
 Pauson-Khand reaction using molecular sieves 182–5  
 2-(1-cyclopentenyl)-1-decene 70–3  
 cyclopropyl arenes, alkynes and alkenes 67–9  
 4-cyclopropylbenzaldehyde 67–9  
  
 diaryl ethanals 95–7  
 diaryl ethanones 92  
 diaryl methanols 161–3  
*o*-diazobromobenzene 232  
 dibutyl-*p*-tolyl-amine 88–9  
*o*-(dichlorophosphine)bromobenzene 232–3  
 1,3-dienes *via* borylation-coupling 70–4  
 diethyl zinc/linked-BINOL catalyst 202–8, 210–16  
 2,3-dihydrofuran, asymmetric Heck reactions of 104–12  
 (+)-11 $\beta$ ,13-dihydroreynosin 197–8  
 (2*R*,3*S*)-2,3-dihydroxy-1-(2-methoxyphenyl)-5-phenyl-1-pentanone 203–5  
 dimeric oximepalladacycle catalyst  
   carbonylation of aryl iodides 244, 245–7  
   synthesis 244–5  
 (*S*)-3-di(methoxycarbonyl)methyl-1-cyclopentanone 219–22  
 dimethyl 3-acetylamino-4-butyrylcyclopent-3-ene-1,1-dicarboxylate 187–8  
 dimethyl malonate and Michael reactions 219–22  
 (4*R*,6*R*)-4,6-dimethyl-2-chloro-1,3,2-dioxaphosphorinane 240–1  
 (2,6-dimethyl-phenyl)-(1-phenyl-2,3,3*a*,8*a*-tetrahydro-1*H*-1-phospha-cyclopenta- $[\alpha]$ inden-8-ylidene)-amines 36–7  
 1,1-diphenyl-1-nonene 131–2  
 1,3-diphenyl-2-propen-1-one 136–7  
 1,2-diphenyl-ethanone 86–7  
*o*-diphenylphosphanylbenzoate group in stereoselective hydroformylation 226, 229–30  
 2-diphenylphosphino-2-methoxybiphenyl 79–80  
 (*S*<sub>FC</sub>)-1-diphenylphosphino-2-[(*o*-diphenylphosphino)phenyl]ferrocene 49–50  
 3-(diphenylphosphino)-4-methyl-styrene 121–2  
 2-diphenylphosphinyl-2-methoxybiphenyl 78–9  
  
 diphosphine complexes, axially chiral 47–51  
  
 enones, cyclic 217, 219–23  
 ESPHOS chiral ligand  
   hydroformylation of vinyl acetate 230, 236–7  
   synthesis and precursors 231–6  
 eudesmonolides 196–200  
  
 ferrocenyl ligands  
   aryl transfer to aldehydes 161–3  
   diphosphine complexes for asymmetric catalysis 47–50  
   ferrocenyl-imino complexes for allylic alkylations 51–6  
   for intermolecular Heck reaction 104–12  
 (*R*)-1-[(*S*)-2-ferrocenyldiene-ethyl-imine]ethyl-diphenylphosphine 54–5  
   precursors 52–4  
 Fibre-Cat<sup>®</sup> 7–8  
  
 glycine imines, alkylation of 27, 31–3  
 Grignard reagents, cross-coupling reactions of 128–33  
  
 Heck coupling reactions  
   asymmetric arylation 97–9  
   intermolecular phenylation of 2,3-dihydrofuran using ferrocenyl ligand 104–12  
   recyclable, soluble, palladacycle catalyst 116–24  
   terminal diarylation 95–7  
   triarylation 92–3  
   vinyl ethers with aryl halides 100–4  
 hydroformylation, stereoselective  
   of allylic alcohols using a catalyst-directing group 226, 229–30  
   asymmetric of styrene to give chiral aldehydes 238–40  
   of vinyl acetate using chiral ligand ESPHOS 230, 236–7  
 2-hydrox-1,5-diketones 210–16  
 (2*R*)-2-hydroxy-1-(2-methoxyphenyl)-1,5-hexanedione 211–12, 213–15  
 3*aS*\*,5*RS*\*-5-hydroxy-3,3*a*,4,5-tetrahydrocyclopenta[*a*]naphthalen-2-one 182–3  
 $\alpha$ -hydroxyketones 210–16

- $\beta$ -iminophosphine ligands  
 palladium catalysed allylic substitutions  
 36, 37–40  
 synthesis 36–7
- indium-mediated allylation 156–7, 160
- Intellectual Property Rights (IPR) 2, 3, 6
- (IPr)Pd(allyl)Cl catalyst 86–90
- 1-*RS*-( $\pm$ )-[(1-isopropyl-2-methyl)prop-2-enyl](2-diphenylphosphanyl)benzoate 227–8
- (1*R*\*,2*R*\*)-( $\pm$ )-[(1-isopropyl-4-oxo-2-methyl)butyl](2-diphenylphosphanyl)benzoate 228–9
- Jacobsen's salen and asymmetric *syn*-alk-1-ene-3,4-diols 164–8
- L-733,060 synthesis 178–80
- (*S,S*)-linked-BINOL 202, 211
- Lucite process 11–12
- metathesis reactions, ring-closure  
 and cross en-yne reactions using Ru  
 carbene complexes 169–73  
 synthesis of neurokinin receptor  
 antagonist 178–90  
 using recyclable polymer-supported Ru  
 catalyst 174–9
- 4-methoxybiphenyl 75–6, 89–90
- (*E*)-methyl 2-carbomethoxy-3,5-diphenylpent-4-enoate 45–6
- methyl (1*S*\*,4*R*\*,4*aS*\*,9*S*\*,9*aS*\*)-2,3,4,9,9a-hexahydro-1*H*-1,4, methano-fluoren-9-ylacetate 151–2
- methyl (2*EZ*)-3-[2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]acrylate 149–51
- methyl methacrylate, Lucite process for 11–12
- 8-(1-methyl-2-trimethylsilylvinyl)-3,4-dihydro-2*H*-naphthalen-1-one 17–18
- 1-[(4-methylphenyl)sulfonyl]-2,3,6,7-tetrahydro-1*H*-azepine 171–2
- Michael reactions, asymmetric  
 $\alpha$ -hydroxyketones to 2-hydroxy-1,5-diketones 210–16  
 enantioselective addition of malonates to  
 cyclic enones 217, 219–23
- molecular sieves  
 and diethyl zinc/linked-BINOL 202,  
 205–7, 211, 213–15  
 and Pauson-Khand reaction 182–5
- N*-(4-cyanophenyl)-morpholine 141–2
- N*-di-isopropylphosphino-*N*-methylpiperazine 82–3
- N*-heterocyclic carbene ligand 86–90
- N*-*tert*-butoxycarbonyl- $\beta$ -bromodehydroaminobutyric acid,  
 methyl ester of 61–2
- N*-*tert*-butoxycarbonyl-(*Z*)-[ $\beta$ -(2,3,7-trimethylbenzo[*b*]thien-6-yl)]dehydroaminobutyric acid, methyl ester of 62–3
- N*-tosyldiallylamine, cyclisation of 176–7
- neurokinin receptor antagonist, synthesis of 178–80
- nickel-on-carbon heterogeneous catalyst  
 aromatic aminations 141–2  
 and coenzyme Q<sub>10</sub> precursor 142  
 coupling of vinyl zirconocenes and aryl  
 halides 145–6  
 preparation 139–40  
 reductions of aryl chlorides 144–5  
 Suzuki couplings 140–1
- N,N*-dimethyl-2-[1,2,2-(triaryl)ethyleneoxy]ethanamines 92–4
- hydrolysis to triaryl ethanones 94
- N,N*-dimethyl-2-[2,2-diarylethyleneoxy]ethanamine 95–7
- hydrolysis to diary ethanals 96, 97
- 1,3,6,8,9,10,11,12,13-nonahydro-2,7-dioxacyclodeca[*e*]indene 194–5
- nonpolar biphasic catalysis 114–16
- nonylcyclopropane 128–9
- 1-octenyl-4-trifluoromethylbenzene 145–6
- olefin metathesis *see* metathesis reactions
- organoboron compounds, arylation with 18–19, 20
- Oxo process 9–11
- (3-oxo-2-pentylidene-cyclopentyl)-acetaldehyde 191–2
- palladacycle catalyst 117–24
- palladium catalysis  
 asymmetric allylic substitution 35–40  
 carbonylation of aryl iodides 245–7  
 cross-coupling reactions 133–8  
 cyclisation reactions 185–90  
 enantioselective allylic alkylations 51–6  
 and Heck coupling reactions 91–112  
 and Sonogashiro coupling reactions 113–25  
 and Suzuki coupling reactions 59–90
- Pauson-Khand reaction 182–5

- (9*H*,9'*H*,10*H*,10'*H*,11*H*,11'*H*,13*H*,13'*H*,14*H*,14'*H*,15*H*,15'*H*-perfluorotricosane-12, 12'-di-yl)bis[(4*S*)-4-phenyl-2-oxazoline], precursors and synthesis 41–5
- phase tags 114
- phenyl biphenyl-4-carboxylate 245–6
- 3-phenyl-3-hydroxy-4-(1-acetoxyhexylidene)tetrahydrofuran 186–7
- 2,2-(1,3-phenylene)bisoxazoline as catalyst 74–7
- 1-(4-phenylethynyl-phenyl)-ethanone 114–15
- 1-[4-(2-phenylethynyl)phenyl]ethan-1-one 124
- 2-phenylpropionic acid 243–4
- 3-phenylpropionic acid 243–4
- phosphine ligands *via* Suzuki coupling 77–81
- phosphino-amine ligands 81–6
- poly(4-methylstyrene) as phase tag 114
- polymer-supported catalysts 117–24, 174–6
- (*S,S,S*)-pyrrolidine-2-carboxylic acid (2-hydroxyl-1,2-diphenyl-ethyl)-amine 208–10
- pyrrole, carboxylation of 247–8, 249–50
- pyrrole-2-carboxylate 249–50
- Rh Oxo process 9–11
- rhodium catalysed reactions  
[2 + 2 + 2] cyclotrimerisation in aqueous media 193–6
- cross-coupling reactions in aqueous media 147–53
- intramolecular alder-ene reaction 190–2
- RuH<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub>
- akenylation of aromatic ketones 17–18
- alkylation of aromatic ketones 16–17, 19–20
- arylation of aromatic ketones 18–19, 20
- synthesis of 14–16
- ruthenium catalysts for metathesis carbene complexes 169–71
- polymer supported recyclable 174–6
- ruthenium-amido complexes 217–23
- for enantioselective Michael reaction 219–23
- synthesis of 217–19
- Sonogashira coupling reactions
- biphasic coupling of 4-bromoacetophenone with phenylacetylene 113–16
- reaction of 1-(4-bromophenyl)ethan-1-ol with phenylacetylene 124
- recyclable, soluble, palladacycle catalyst 116–24
- styrene, asymmetric hydroformylation of 238–40
- Suzuki coupling reactions
- 3-fluorobenzene with phenylboronic acid 84–5
- aryl bromides with aryl boronic acids catalysed by palladium-oxazolines 74–7
- aryl chlorides with phenylboronic acid using phosphino-amine ligands 81–2, 83–6
- biphasic reaction of 4-bromoacetophenone with phenylboronic acid 115–16
- in situ* borylated bromobenzo[*b*]thiophenes with  $\beta$ -bromodehydroamino acid derivatives 60–4
- in situ* synthesis of vinylboron compounds leading to unsymmetrical 1,3-dienes 70–3
- 4-tosylcoumarins with aryl boronic acids 64, 65–7
- OPPh<sub>2</sub>(*o*-C<sub>6</sub>H<sub>4</sub>Br) with aryl boronic acids 77–80
- phenylboronic acid and 2-chlorobenzonitrile 140–1
- and recyclable, soluble, palladacycle catalyst 116–24
- Suzuki-Miyaura coupling reactions
- 4-chloroanisole with phenylboronic acid using *N*-heterocyclic carbene ligand 86, 89–90
- cyclopropylation of aryl, alkynyl, and vinyl bromides 67–9
- (2*S*)-*tert*-butyl 2-amino-4-bromopent-4-enoate 31–2
- tert*-butyl *N*-(diphenylmethylene)glycinate, alkylation of 31–3
- 1-[4-(*S*)-*tert*-butyl-2-oxazoline-2-yl]-2(*S*)-(diphenylphosphino)ferrocene as ligand for Heck reaction 104, 110–12
- synthesis and precursors 106–10
- tin-mediated allylation 156, 157–8, 160
- titanocene-catalysed cyclisation of epoxygermacrolides 196–200
- (*S*)-4-tolyl-phenyl methanol 61–3

- (S<sub>FC</sub>)-1-[(S)-*p*-tolylsulfinyl]-2-[(*o*-diphenylphosphino)phenyl]ferrocene 47–9
- 4-tosylcoumarins 65, 67
- trans*-di(μ-acetato)-bis[3-(diphenylphosphino)-4-styryl]dipalladium(II) 122–3
- 1,2,3-triaryl ethanones 92–4
- 8-(2-triethoxysilanylethyl)-3,4-dihydro-2*H*-naphthalen-1-one 16–17
- triorganoindium compounds in cross-coupling reactions 133–7
- triphenylindium 134
- vinyl acetate, hydroformylation of 230, 236–7
- vinyl ethers, Heck coupling of 100–4
- zinc-mediated allylation 156, 159–60

*With thanks to W. Farrington for the creation of this index.*