

# Index

- Abbreviated Mental Test Score 69  
acamprosate 63  
access to care 23–7  
    ethnic minorities 84  
    homelessness 87, 88, 89  
    psychological treatments 103  
    routes to services 26  
accident & emergency departments 15–16  
acetaldehyde dehydrogenase 61  
acetylcholinesterase inhibitors 69–70  
acknowledgement of patient distress 9  
acute general hospitals 15–18, 20, 23  
acute stress reactions 28  
addiction 55–9, 60–63  
adherence to treatment 17, 101  
adjustment reactions 28  
admission to hospital 15–18, 20, 23, 50, 91–3  
adolescents 72–5, 93  
aftercare, violent incidents 21  
aggression 8, 9, 19–22, 109–10, 111–12  
    self-harm 1, 15–16, 39, 51, 77  
agnosia 69  
agoraphobia 29, 30–1, 34  
akathisia 99  
alcohol-related problems  
    addiction 60–3  
    diagnostic hierarchies 4  
    homelessness 86, 87, 88  
    physical illness-associated 16  
    prevalence 1  
Alzheimer-type dementia 68, 69, 71  
American Psychiatric Association 48  
amnesia 62, 68  
amok 83  
amphetamine misuse 56, 57, 58  
amylobarbitone 21, 22  
anorexia nervosa 1, 74  
anti-anxiety medication 99–100  
    *see also* benzodiazepines; serotonin reuptake inhibitors  
anticholinergic drugs 99  
antidepressant drugs  
    childbearing 38  
    cultural aspects 84  
    depression 96  
    maintenance treatment 37  
    personality disorders 51  
    psychiatric disorder in physically-ill patients 17  
    psychotherapy comparison 36  
    side effects 97  
antipsychotic drugs 97–9  
    atypical 98  
    depot 45, 98  
    personality disorders 51  
    schizophrenia 45–6, 97–9, 101  
    side effects 45, 101  
anxiety  
    cognitive models 28, 29  
    disorders 28–34  
    homelessness 88  
    intellectual disability 78, 79  
    management 29–30, 31, 32, 33  
    obsessive-compulsive disorder 29, 33–4  
    older people 65–6  
    physically-ill patients 17  
    physiological symptoms 65  
    prevalence 1  
    primary care 11–14  
    sexual dysfunction 53  
apathy 8  
aphasia 69  
appearance 3  
appropriate behaviour modelling 9  
approved clinicians 93  
apraxia 68  
Asperger syndrome 77  
assertive outreach teams 24  
assessment *see* diagnosis and assessment  
asylums 23  
asylum seekers 86, 87  
attention deficit hyperactivity disorder 73–4  
attorney, power of 93  
atypical antipsychotics 97, 98  
AUDIT 61  
autism 77  
behaviour  
    *see also* cognitive behavioural therapy  
    coping strategies 7–8  
    dementia 70  
    distressed patients 7–10  
    externalising disorders in children/  
        adolescents 73–4  
    intellectual disability 76–7  
    mental state examination 3  
    modelling appropriate 9  
    overshadowing 77  
    violence prediction 20  
benzodiazepines  
    alcohol detoxification 62  
    anxiety disorders 30  
    misuse 56  
    older people 66  
    prescribing 99, 100–101  
    rapid tranquillisation 21, 22  
    side effects 99  
    withdrawal 58  
bereavement 11  
beta-blockers 30  
bipolar affective disorder 1, 38, 40–43, 79, 100  
black and minority ethnic groups 87  
bodily distress mannerisms 9  
body image disorders 17  
brain injury 10  
bulimia nervosa 74  
buprenorphine 58  
buspirone 30  
CAGE questionnaire 61  
Caldicott Guardians 94, 95  
calm manner 9, 21  
cannabis 56, 57, 58  
capacity 3, 91, 92, 93, 94  
carbamazepine 100  
cardiovascular assessment 3  
Care Programme Approach 24, 80  
carer support workers 24  
challenging behaviour 7–10, 77  
    *see also* violence  
Child and Adolescent Mental Health Services 73,  
    74, 105  
childbearing 38, 40, 101  
children 72–5, 93, 105  
chlormethiazole 62  
chlorpromazine 97  
choice questions 2  
chronic conditions 12, 35, 37–8, 41–3  
classification  
    alcohol-related disorders 62  
    bipolar disorder 40  
    child/adolescent mental health problems 73  
    depression 37  
    intellectual disability 78  
    late-onset paranoid disorders 66  
    mania 47  
    personality disorders 48, 49  
    primary care problem typologies 12  
    sexual dysfunction 52–3  
clinical risk 108–13  
clinical tests 3, 61, 69, 109

- clinic layouts 19  
 clomipramine 34  
 clonazepam 100–101  
 closed/open questions 2, 3  
 clozapine 97, 98, 102  
 Clunis, Christopher 24  
 Cluster A/B/C (DSM-IV) disorders 48–50  
 cocaine 56, 57, 58  
 cognitive analytic therapy 105  
 cognitive behavioural therapy  
   anxiety disorders 28, 29, 30, 31, 33, 34  
   depression 36  
   overview 103–4  
   physical illness 17, 18  
   schizophrenia 46  
 cognitive models 28, 29  
 collaboration with patients 101  
 collateral patient history 3  
 combination antipsychotics 99  
 Committee on Safety of Medicines 71  
 Common Assessment Framework 72  
 common law duty of care 91, 92, 93, 101  
 communication difficulties 77, 84  
 community  
   care 23–4, 25, 26, 46  
   detoxification 62  
   mental health teams 23, 24, 25, 26, 27, 46  
   treatment orders 93  
 competence 93  
 compliance 17, 101  
 compulsory admission/treatment 90, 91  
 concordance with treatment 17, 101  
 conduct *see* behaviour  
 confidentiality 91, 94  
 confusion 4  
 consent to treatment 92–3, 101  
 coping patterns/strategies 7–8  
 cost of antidepressants 37  
 counselling 103, 104–5  
 County Asylums Act (1845) 24  
 crisis resolution/home treatment teams 24, 27  
 crisis responses 19–22, 24, 26–7, 41–2, 92  
 Crown Courts 93  
 Croydon Integrated Adult Mental Health  
   Service 25, 26  
 cultural aspects 81–5, 86–90  
 current problems assessment 2, 3  
 curses (obeah) 82
- Data Protection Act (1998) 94  
 decision-making 109  
 delirium 4, 66  
 delirium tremens 60, 61, 62  
 delusions  
   cultural aspects 82  
   definition 4, 45  
   medication 97  
   older people 66  
   violence prediction 20  
 dementia 68–71  
 denial 8  
 dependence 55–9, 60–63  
 depot antipsychotics 45, 98  
 depression 35–9  
   bipolar disorder 38, 41  
   childbearing 38  
   children/adolescents 74
- cultural aspects 81, 82  
 electroconvulsive therapy 38  
 forms/types 35–6, 37, 96–7  
 homelessness 87, 88  
 medication 36–7, 65, 96–7  
 mixed states 41  
 older people 64–5  
 physically-ill patients 17  
 prevalence 1  
 primary care 11–14  
 seasonal affective disorder 38  
 self-harm 15, 39  
 somatic syndrome 36  
 suicide 2, 13, 22, 39, 97, 111  
 detention, compulsory 90, 91  
 detoxification 61–2  
 developmental disorders 73, 74  
 dhat 83  
 diagnosis and assessment  
   bipolar disorder 40–41  
   dementia 69  
   depression 35, 36  
   drug misuse 58  
   generalised anxiety disorder 30  
   initial psychiatric 1–6  
   intellectual disability 77–9  
   panic disorder 31  
   personality disorders 48, 49–50  
   post-traumatic stress disorder 33  
   schizophrenia 45  
   self-harm, hospital situation 16  
   social phobias 32  
 Diagnostic Criteria for Learning Disabilities 78  
 diagnostic hierarchy of disorders 4  
 dialectical behaviour therapy 16, 50, 103, 104  
 diazepam 21  
 disability rates 5  
 distressed patients 7–10, 81–2  
 disulfiram 62–3  
 dosages 97, 98, 100  
 Down syndrome 76, 79  
 drive, fitness to 5  
 drug misuse 1, 4, 16, 55–9, 88  
 drug treatments *see* medication  
 dysfunctional coping 7–8  
 dysthymia 35, 37–8  
 dystonias 99
- early intervention teams 24  
 eating disorders 17, 74, 78, 79  
 eclectic approach 105  
 ecstasy 56, 57, 58  
 elderly people 64–7  
 electroconvulsive therapy 38  
 emergencies 15–16, 19–22, 41–2, 92  
 emotional distress 7–10, 81–2  
 empathy 9  
 engagement 12–13  
 environmental causes 72  
 epidemiology  
   alcohol-related problems 60–61  
   anxiety 1  
   childhood problems 72–3  
   dementia 68  
   depression in older people 64  
   homelessness 87–8  
   personality disorders 49
- psychiatric morbidity 1, 15  
 psychotic disorders in elderly people 66  
 self-harm/suicide 1, 2, 22, 111  
 violence 20, 109–10, 111–12  
 ethnic aspects 81–5  
 explanations of illness 82  
 extrapyramidal side effects 97, 98, 99  
 eye movement desensitisation reprocessing 33
- family 3, 72, 83–4, 93, 105  
 fixed coping behaviour 7–8  
 FP10 (MDA) prescriptions 58  
 fragile X syndrome 76  
 functional aspects 4, 7, 78
- gateway primary-secondary workers 25  
 gender problems 54  
 generalised anxiety disorder 28–30  
 general practitioners 11–14, 24–5, 72, 74  
 genetic predisposition 40, 44, 49  
 goals, achievable for patients 10  
 graduate primary care workers 25  
 grief and loss 11  
 group therapy 51, 105–6
- hallucinations 97  
   cultural aspects 82  
   definition 4  
   older people 66  
   schizophrenia 44–5  
   violence prediction 20  
 haloperidol 21, 22  
 HCR-20 scale 109  
 healers, traditional 83, 84  
*The Health of the Nation* 60  
 hepatic enzymes 100  
 heroin 56, 58  
 history of service provision 23–4  
 history-taking, initial 2  
 homelessness 86–90  
 hospital admission 15–18  
   compulsory 91  
   emergency 20  
   mental hospitals 23  
   mentally disordered offenders 93  
   personality disorders 50  
   somatisation 17, 18  
 housing 86–90  
 Human Rights Act (1998) 91, 92, 94  
 5-hydroxytryptamine *see* serotonin reuptake  
   inhibitors  
 hyperkinetic disorder 73  
 hypogonadism 53  
 hypomania 40, 41
- Improving Access to Psychological Therapies 103,  
   106  
 inflexible coping behaviour 7–8  
 information  
   patient-identifiable 94  
   sources  
   medication 101–2  
   psychological therapies 106  
   risk management 110  
   technology 14  
 initial assessment 1–6  
 initial crisis management plan 27

- injected drugs 21, 22, 45, 57, 98  
 insight 3  
 institutional care 15–18, 20, 23, 87–8, 91–3  
 integrated mental health care 23, 24, 26  
 integrative psychological therapy 105  
 intellect 3  
 intellectual disability 76–80  
 internalising disorders, children 73, 74  
 International Classification of Diseases 48, 49, 78  
 International Late Onset Schizophrenia Group 66  
 interview technique 2, 4, 10, 12  
 intramuscular route 21, 22, 45, 98  
 intravenous route 21, 22, 57
- jiryan 83
- laboratory tests 3, 69  
 lamotrigine 100  
 latah 83  
 late-onset paranoid disorders 66  
 law 91–5  
   County Asylums Act (1845) 24  
   Lunacy Act (1890) 24  
   Lunatics Act 24  
   Mental Capacity Act (2005) 3, 91, 92–3  
   Mental Health Act 16, 19, 20, 25, 27, 48, 82, 83  
     (1983) 48, 91–2  
     (2007) 91, 93  
   Mental Health (Patients in the community)  
   Act (1995) 93  
   Mental Treatment Act (1930) 24  
   National Assistance Act (1948) 94  
 learning disability 74  
 Lesch–Nyhan syndrome 76  
 lithium 41, 42, 100  
 London, homelessness 87  
 long-term prediction 20, 109  
 lorazepam 21, 22  
 Lunacy Act (1890) 24  
 Lunatics Act 24
- magistrates 92  
 major incidents 20–22  
 malignant social pathology 70  
 mania 40, 41, 47  
 mannerisms, distress 9  
 marijuana 56, 57, 58  
 marriage, arranged 84  
 MDMA 56, 57, 58  
 medical records 21, 25, 94, 110  
 medication 96–102  
   alcohol abstinence 62–3  
   anxiety disorders 29, 99–100  
   obsessive–compulsive disorder 34  
   older people 66  
   panic disorder 30, 31  
   post-traumatic stress disorder 33  
   bipolar affective disorder 38, 41  
   concordance with treatment 17, 101  
   dementia 69–70, 71  
   depression 36–7, 65, 96–7  
   dosage/duration 97, 98, 100  
   drug misuse 55–9  
   intellectual disability 80  
   interactions 57  
   mood-stabilising 41, 42, 100  
   personality disorders 51
- preliminary physical tests 3  
 primary care 13  
 psychiatric disorder in physically-ill  
   patients 17  
 rapid tranquillisation 20–22  
 schizophrenia 45–6, 97–9  
 sexual dysfunction cause 53  
 side effects 97, 99, 100  
 social phobia 32  
 medium-term risks 109  
 memory clinics 70  
 Mental Capacity Act (2005) 3, 91, 92–3  
 Mental Health Act 16, 19, 20, 25, 27, 82, 83  
   (1983) 48, 91–2  
   (2007) 91, 93  
 Mental Health (Patients in the community)  
   Act (1995) 93  
 Mental Health Review Tribunal 92, 93  
 mental hospitals 23  
 mental impairment 91  
 mentally disordered offenders/prisoners 93  
 mental state examination 3  
 Mental Treatment Act (1930) 24  
 methadone 58  
 3,4-methylenedioxymethamphetamine  
   56, 57, 58  
 midazolam 21, 22  
 migration 83  
 MIND 5  
 mindfulness techniques 103, 104  
 Mini-Mental State Examination 68, 69  
 mirroring behaviour 9  
 mirtazapine 96  
 mixed affective states 41  
 moclobemide 37  
 modelling appropriate behaviour 9  
 monoamine oxidase inhibitors 36, 96  
 mood  
   disorders  
     *see also* depression  
     bipolar 1, 38, 40–43, 79, 100  
     children/adolescents 73, 74  
     diagnostic hierarchies 4  
     intellectual disability 78, 79  
     physical illness-associated 16  
     stabilising drugs 42, 51, 100  
     mental state examination 3  
   morbidity 1, 11  
   mortality rates 5  
   multi-agency public protection  
   arrangements 94  
   multidisciplinary approach 23, 24, 78, 88  
   multiethnic aspects 81–5  
   multisystems therapy 73
- Nafsiyat Intercultural Therapy Centre 84  
 National Assistance Act (1948) 94  
 National Institute for Health and Clinical  
   Excellence 13  
   acetylcholinesterase inhibitors 69, 70  
   anxiety disorders 28, 29, 30, 65  
   attention deficit hyperactivity disorder 73, 74  
   atypical antipsychotics 98  
   depression 35  
   psychological treatments 103, 105, 106  
   self-harm guidelines 16  
 National Service Framework 24, 25, 64, 65, 74
- National Survey of Psychiatric Morbidity in  
   Great Britain 11  
 nearest relative 93  
 need 25, 26, 86  
 nefazodone 37  
 neuroleptic malignant syndrome 98  
 neurological assessment 3  
 neuroses  
   anxiety 11–14, 17, 28–34, 65–6, 78–9, 88  
   definition 4  
   NHS guidelines 10  
   NHS Plan 23, 24  
   no fixed address rotas 89  
   non-opioid drug withdrawal 58  
   non-statutory care provision 23, 62, 71  
   norms, cultural 81
- obeah (curses) 82  
 obsessive–compulsive disorder 29, 33–4,  
   74, 79  
 Office of National Statistics 28  
 olanzapine 21, 22, 41, 42, 97  
 older people 64–7  
 open/closed questions 2, 3  
 opioid misuse 56, 57, 58  
 oppositional defiant disorder 73  
 organic disease 4, 16, 20  
 outreach teams 24
- pain management model 8  
 panic 8, 29, 30, 31, 32  
 paraldehyde 21, 22  
 paranoia 3, 45, 66  
 paraphilias 54  
 parenting/family 72, 73, 83–4, 93, 105  
 parkinsonian syndrome 99, 101  
 patient-held records 25  
 patient history 20, 77  
 patient-identifiable information 94  
 patient needs 25, 26, 86  
 perceptions 3  
 persecutory beliefs 3, 45, 66  
 personal accounts 5  
   alcohol dependence 63  
   anxiety disorders 34  
   brain injury 10  
   children/adolescents 75  
   depression 39  
   ethnic aspects 85  
   hospital situation 18  
   intellectual disability 80  
   psychosexual problems 54  
   schizophrenia 47  
 personality disorders 48–51  
   aetiology 49  
   diagnosis and assessment 48, 49–50  
   epidemiology 49  
   homelessness 87, 88  
   hospital admission 50  
   intellectual disability 78, 79  
   medication 51  
   prevalence 1  
   psychological treatment 50–51  
   personality traits 4  
 pervasive developmental disorder 74  
 pharmacotherapy *see* medication  
 phobias 28, 29, 32, 74

- physical aspects  
*see also* violence  
 illness  
   general hospital situation 15–18  
   homelessness 88  
   mental disorder-associated 12, 16–17  
   older people 64–5  
 physical tests 3  
 sexual dysfunction 53  
 stress 83  
 place of safety 92  
 plans 10, 19–20  
 police 21–2, 92, 94  
 population-based studies 11  
 possession by spirits 81, 82  
 postpartum depression 38  
 post-traumatic stress disorder 28, 29, 32–3  
   children/adolescents 72  
   homelessness 87  
   intellectual disability 79  
 post-traumatic therapy 22  
 posture 9, 21  
 poverty 86–90  
 practical risk management 109–10  
 Prader–Willi syndrome 76  
 primary care  
   child/adolescent mental health 72, 74  
   counselling 104–5  
   depression 35  
   mental health problems in 11–14  
   NICE guidelines 25  
   reviews, proactive 14  
   safety netting 14  
   shared mental health care 24–5  
   system development 14  
   typologies of mental health problems 12  
 prison services 93, 94  
 private care 23  
 probation orders 94  
 professionals 21, 22, 108–9, 110  
   attitude to homelessness 88  
   carer support workers 24  
   community mental health teams 23, 24,  
     25, 26, 46  
   crisis resolution/home treatment teams 24, 27  
   gateway primary-secondary workers 25  
   general practitioners 11–14, 24–5, 72, 74  
   graduate primary care workers 25  
   legal requirements 91–5  
   multidisciplinary approach 23, 34, 78, 88, 94  
   psychological treatments 103  
   risk control 108–9, 110  
   second opinion approved doctors 92  
 promethazine 22  
 psychiatric assessment 1–6, 78  
 Psychiatric Assessment Schedule for Adults with  
   Developmental Disability 78  
 psychiatric emergencies 19–22, 41–2, 92  
 psychodynamic therapy 104, 105  
 psychologically-based physical syndromes 17, 18  
 psychological risk factors 20  
 psychological treatments 103–7  
   anxiety disorders 29, 30  
   basic hospital clinician skills 17  
   choice 106  
   depression in older people 65  
   obsessive–compulsive disorder 34  
   organisations 107  
   panic disorder 31–2  
   personality disorders 50–51  
   post-traumatic stress disorder 33  
   primary care 13  
   schizophrenia 46  
   sexual dysfunction 54  
   social phobias 32  
 psychopathic disorders 91  
 psychoses 4, 11, 87–8  
   *see also* psychotic disorders; schizophrenia  
 psychosexual disorders 52–4, 83  
 psychosocial impairment 73  
 psychosocial interventions 79, 80  
 psychotherapy 36  
 psychotic disorders 20, 37, 38, 55  
   *see also* schizophrenia  
 puerperal psychosis 40  
 rapid tranquillisation 20–22  
 recognition 8–9, 12–13  
 recordkeeping 21, 25, 94, 110  
 recurrent depression 12  
 referral to specialists  
   acute bipolar disorder states 41  
   criteria 25  
   dementia 70  
   depression 36, 65  
   intellectual disability 77  
   personality disorders 49  
   primary care function 13–14  
   report components 4–5  
 reflecting patient distress 9  
 refugees 87  
 relapse prevention 42–3, 46  
 relationship problems 52  
 religion 82, 83  
 residential services 26  
 responsible clinician 93  
 restraint 21  
 Rethink 5, 102  
 review processes 14  
 risk 108–13  
   Care Programme Approach 24  
   child/adolescent mental health 72–3  
   electroconvulsive therapy 38  
   initial assessment 2–3  
   management 108–10  
   mental health emergencies 19–21  
   primary care situation 13  
   self-harm, hospital situation 16  
   suicide 2, 15, 111  
   violence 20  
 risperidone 22  
 Royal College of General Practitioners 10  
 Royal College of Psychiatrists 5  
 safety *see* risk  
 safety netting 14  
 St John's wort 97  
 schizophrenia 44–7  
   antipsychotics 45–6, 97–9, 101  
   cultural aspects 82, 83  
   family intervention 105  
   homelessness 86, 87  
   intellectual disability 78–9  
   older people 66  
   prevalence 1  
   Rethink 5, 102  
   violence risk 20, 110  
 school 72, 73, 74  
 seasonal affective disorder 38  
 secondary care  
   *see also* referral to specialists  
   bipolar disorder 41–2  
   depression 36  
   mental health services 23, 24–5  
   NICE guidelines 25  
   role 25  
   teams 24–5, 26  
 second opinion approved doctors 92  
 sedation 97, 98  
 selective serotonin reuptake inhibitors 30, 96, 97  
   depression 36, 37  
   older people 65, 66  
   social phobias 32  
 self-harm 1, 15–16, 39, 51, 77  
 self-help 30, 31, 33, 42–3  
 serotonin-noradrenaline reuptake  
   inhibitors 36, 37  
 serotonin reuptake inhibitors 30, 96, 97  
   depression 36, 37  
   older people 65, 66  
   social phobias 32  
 service models 80, 88–9, 90  
 service provision 23–7, 84, 87, 88, 89, 103  
 severe mental impairment 91  
 sexual dysfunction 17, 52–4  
 shared care 24–5  
 short-term prediction 20, 109  
 side effects 97, 98  
   antidepressants 36, 37  
   antiepileptic drugs 100  
   antipsychotics 45, 101  
   benzodiazepines 99  
   lithium 100  
 simple schizophrenia 45  
 sleep disturbance 43  
 Smith–Magenis syndrome 76  
 social phobias 29, 32  
 social risk factors 20  
 social treatment 65, 70  
 sodium valproate 41, 42  
 somatic syndrome 36  
 somatisation 12, 17, 18, 82, 83  
 specialists *see* referral to specialists; secondary care  
 specific (isolated) phobias 32  
 speech 3  
 spirit possession 81, 82  
 staff *see* professionals  
 standardised mortality rate 5  
 stepped care model 13, 17  
 stereotyping 88  
 stigma 5, 11  
 stress 28, 83  
 substance misuse *see* drug misuse  
 suicide  
   antidepressant risk 97  
   depression 2, 13, 22, 39, 97, 111  
   emergency situation 22  
   lithium 100  
   prevalence 1  
   primary care questions 13  
   risk 2, 15, 111

- supervised discharge 93  
susto 83  
system development 14
- tardive dyskinesia 99  
team-based care 23, 24, 25, 26, 27, 46  
teratogenicity 101  
termination of interviews 10  
terminology, troublesome 4  
testamentary capacity 94  
tests  
    AUDIT 61  
    laboratory tests 3  
    risk assessment 109  
third-wave approaches 103, 104  
thoughts 3  
    *see also* delusions; paranoia  
trance-like states 82  
TREC Collaborative Group 21  
tricyclic antidepressants 30, 36, 37, 96  
Tuke, William 23
- two-phase surveys 11  
typologies of problems 12
- UK mental health service  
    provision 23–7
- valid consent 93  
valproate 100–101  
velocardiofacial syndrome 76  
venlafaxine 37, 96  
verbal expressions of distress 8, 9  
violence  
    emergencies 19–22  
    NHS guidelines 10  
    risk 20, 109–10, 111–12  
    self-harm 1, 15–16, 39, 51, 77  
vitamin B 62  
volatile substance misuse 57, 58  
voluntary organisations  
    alcohol detoxification 62  
    Alzheimer's disease 71
- ethnic 84  
    mental health service provision 23  
    role 5  
voting rights 94
- watchful waiting 12, 13, 33  
Williams syndrome 76  
witchcraft 81, 82  
withdrawal of non-opioid drugs 58  
withdrawal syndrome 55, 56, 62  
World Health Organization 1  
    Alcohol Use Disorders Identification  
        Test 61  
    depression 35  
    disability due to mental disorder 5  
    ICD-10 classifications 37, 48, 49, 78  
    study of mental disorder in general  
        healthcare 11
- Zito, Jonathan 24  
zuclopenthixol acetate 21