

INDEX

- Acrylates, ESR analysis:
 chain-transfer reaction, 111–119
 penultimate unit effect, dimeric model
 radicals, 124–128
- Acrylic polymers:
 nitroxide kinetics and HALS stabilization,
 272–276
 photoinitiation rate measurements, 267–269
- Acrylonitrile-butadiene-styrene (ABS) polymers,
 electron spin resonance imaging:
 FTIR *vs.*, 249–250
 photodegradation, 241–243
 spatially resolved degradation:
 aged heterophasic polymers, 238–243
 basic principles, 230–233
 experimental protocols, 233–235
 hindered amine stabilizers, 235–238
 thermal aging, 238–241
- Activation rate constants, ESR analysis,
 monomeric and dimeric alkyl
 bromides, 128–129
- Alkyl bromides, ESR analysis, penultimate unit
 effect:
 dimeric model radicals, 124–128
 monomeric model radicals, 122–124
- Alumina interactions, dendrimer structures, spin
 labeled nitroxides, 295–296
- Amino acids, dendrimer structures, spin labeled
 nitroxides, 295–296
- Amphiphilic probes, strategic applications,
 170–171
- Anisotropic properties:
 conductive polymers, high-field ESR, spin
 probe method, 332–333
 electron spin echo envelope modulation, 44–49
 electron spin resonance, basic principles, 5–9
 ESR spectrum, 54–55
 hyperfine interactions, *g*-tensor and, 10–12
 nitroxide magnetic tensors, 55–60
- Anisotropic viscosity model, spin-labeled
 polymer orientation, 68
- Arrhenius behavior:
 conductive polymers:
 saturation-transfer ESR, 329–330
 spin probe method, 332–333
 poly(2-hydroxyethyl methacrylate) segmental
 mobility, 158–161
 polymer dynamics, NMR and fluorescence
 techniques, 134–135
 polystyrene segmental mobility, 150–155
 segmental mobility of polymer radicals,
 143–147
- A*-tensor anisotropy, nitroxide rotational
 dynamics, 136–140

- Atom-transfer radical addition (ATRA):
 ESR analysis of radical polymerization,
 104–105
 metallo-dendrimer structures, 301
- Atom transfer radical polymerization (ATRP),
 ESR analysis:
 basic principles, 102–105
 chain length dependence, 105–111
 controlled chain length models, 106–111
 tert-butyl methacrylate, 105–106
 chain transfer reaction, 111–119
 chain length-dependent changes, 117–119
 controlled chain lengths, 113–117
 tert-butyl acrylate, 111–113
 penultimate unit effect, 119–129
 activation rate constants, 128–129
 dimeric model radicals, 124–128
 monomeric model radicals, 122–124
- Attenuated total reflectance (ATR) FTIR,
 acrylonitrile-butadiene-styrene
 polymers, 249–250
- Automotive paint technology, polymer coating
 stabilization, 256–258
- Azobisisobutyronitrile (AIBN), photoinitiation
 rate measurements, 265–269
- Basis sets, stochastic Liouville equation:
 calculation, 73–74
 pruning, 78–79
 symmetrization, 74
 truncation parameters, 74–78
- Best-fit values:
 poly(2-hydroxyethyl methacrylate) segmental
 mobility, 156–161
 polystyrene segmental mobility, 150–155
- Bioreduction rate, dendrimer structure, ESR
 analysis, nitroxide spin labeling,
 291–296
- Bjerrum length, polyelectrolyte-counterion
 interactions, 173–178
- Bohr magneton, double resonance ESR, dipole-
 dipole coupling, 28–29
- Boltzmann relation, spin-labeled polymer
 orientation, 65–68
- Burgers-Korteweg-de-Vries theory, high-field
 electron spin resonance, conductive
 polymers, charge carrier relaxation and
 dynamics, 320–326
- Catalyst-coated membranes (CCMs), *in situ* ESR
 experiments, fuel cell technology,
 222–224
- [2]Catenanes:
 coconformation and motional modes, 178–182
 mesoscopic interfaces and nanostructure,
 182–192
- Cationic nitroxide spin probes:
 dendrimer spin labeling, 291–296
 dendrimer structure, ESR analysis, 284–289
- Cetyl-trimethylammonium bromide (CTAB),
 dendrimer structure, spin probe
 analysis, 286–289
- Chain conformation, polyelectrolyte-counterion
 interactions, 174–178
- Chain-end radicals, Nafion and Dow ionomers,
 membrane-derived fluorinated radicals,
 217–221
- Chain length dependence:
 dendrimer structure, ESR analysis, spin probe
 techniques, 284–289
 ESR analysis of radical polymerization,
 105–111
 chain transfer reactions, 117–119
 controlled chain length models, 106–111
tert-butyl methacrylate, 105–106
- Chain transfer reaction:
 ESR analysis, radical polymerization,
 111–119
 chain length-dependent changes, 117–119
 controlled chain lengths, 113–117
tert-butyl acrylate, 111–113
 polymer coatings, photoinitiation rate
 measurements, 268–269
- Charge carriers, conductive polymers, high-field
 ESR:
 magnetic parameters, 310–317
 relaxation and dynamics, 317–326
- Charge transfer reactions, doped conductive
 polymers, 326–328
- Cobalt complexes, metallo-dendrimer structures,
 300–301
- Coherence:
 electron-electron double resonance, 32–35
 experimental techniques, 33–35
 electron spin echo envelope modulation,
 46–49
- Coil expansion, segmental mobility of polymer
 radicals, 146–147
- Conductive polymers, high-field electron spin
 resonance:
 basic principles, 307–310
 carrier relaxation and dynamics, 317–326
 charge carrier magnetic parameters, 310–317
 highly-doped charge transfer, 326–328
 saturation transfer, 328–330
 spin probes, 330–333
- Cone distributions, spin-labeled polymer
 orientation, 67–68

- Continuous-wave electron spin resonance (CW-ESR):
- anisotropic hyperfine interaction and *g*-tensor, 10–12
 - electron-electron double resonance, 32–33
 - macromolecular materials:
 - mesoscopic interfaces and nanostructure, 182–187
 - ionomers, 187–192
 - in solution, 171–182
 - [2]catenane coconformation and motional modes, 178–182
 - frozen solutions, 172–173
 - polyelectrolyte-counterion interactions, 173–178
 - solvent and temperature effects, 171–172
 - spin labeling:
 - research overview, 166–167
 - strategies for, 167–168
 - spin probing:
 - amphiphilic probes, 170–171
 - hydrogen-bonding probes, 169
 - hydrophobic probes, 169
 - ionic probes, 169–170
 - strategies for, 168–171
 - multifrequency and high-field electron spin resonance, 18
 - nitroxide line shape analysis, 138–140
 - nitroxide magnetic tensors, 55–60
 - paramagnetic quencher accessibility, 14–15
 - pulsed methods, 18–22
- Controlled chain length, ESR analysis of radical polymerization:
- model radicals, 106–111
 - tert*-butyl acrylate radicals, 113–119
- Copolymerization:
- ESR analysis, penultimate unit effect, dimeric model radicals, 128
 - mesoscopic ionomers, 188–192
 - polystyrene segmental mobility, 148–155
- Copper complexes:
- dendrimer structures, additions to, 296–299
 - membrane-derived fluorinated radicals in Nafion, 215–221
- Counterion-polyelectrolyte interactions, solution-based macromolecules, 173–178
- Counterions, Nafion and Dow ionomers, membrane-derived fluorinated radicals, 215–221
- Cross-linked coatings:
- nitroxide kinetics and HALS stabilization, 272–276
 - photoinitiation rate measurements:
 - nonweathered coatings, 267–269
 - weathered coatings, 270–272
- Crystalline polymers, microtoming sensitivity, 250–251
- Data analysis, electron spin echo envelope modulation, 47–49
- Davies electron-nuclear double resonance, experimental techniques, 41–42
- D-band electron spin resonance, conductive polymers:
 - charge carrier relaxation and dynamics, 317–326
 - magnetic parameters, 312–317
 - saturation transfer, 328–330
 - spin probe method, 330–333
- Degradation processes, electron spin resonance imaging of polymeric systems, 94–96
- Dendrimer structures, ESR analysis:
 - basic properties, 279–282
 - metallo-dendrimers, 299–301
 - nitroxide spin labeling, 289–296
 - paramagnetic metal ion additions, 296–299
 - spin probes, 283–289
- Density functional theory (DFT), membrane-derived fluorinated radicals in Nafion, 220–221
- Deuterated spin labels, nitroxide rotational dynamics, 138–139
- Diblock copolymers, mesoscopic ionomers, 188–192
- Dibutyl phthalate (DBP), segmental mobility, 149–155
- Dielectric permittivity, polyelectrolyte-counterion interactions, 174–178
- Diffusion-limited oxidation (DLO) regime:
 - acrylonitrile-butadiene-styrene polymers:
 - FTIR vs ESRI, 249–250
 - thermal aging, 239–241
 - electron spin resonance imaging:
 - polymeric systems, 95–96
 - spatially resolved degradation, 231–233
 - heterophasic propylene-ethylene copolymers, thermal aging, 243–244
- Diffusion models:
 - electron spin resonance imaging of polymeric systems, 93–96
 - slow-motion ESR spectra, 60–63
- Dilute solutions, nitroxide spin label dynamics, 140–141
- Dimeric radicals, ESR analysis:
 - activation rate constants, 128–129
 - chain length dependence, 117–119
 - penultimate unit effect, 124–128

- Dimyristoylphosphatidylcholine (DMPC),
dendrimer structure, spin probe
analysis, 286–289
- Diocetyl phthalate (DOP), segmental mobility,
149–155
- Dipole-dipole (DD) coupling:
conductive polymers, high-field ESR:
charge carrier relaxation and dynamics,
318–326
magnetic parameters, 313–317
spin probe method, 331–333
double resonance ESR methods, 27–29
electron spin resonance principles, 8–9
solution-based macromolecules, frozen
solutions, 172–173
- Direct computation techniques, electron-electron
double resonance, distance
distributions, 38–39
- Direct electron spin resonance, radical
intermediate detection, 204–206
- Direct methanol fuel cell, membrane stability,
201–204
- Director frame:
slow-motion ESR spectra, diffusion models,
61–63
spin-labeled polymer orientation, 63–68
- Dispersion mechanisms, mesoscopic interfaces
and nanostructure, 183–187
- Distance distributions, electron-electron double
resonance, direct computation, 38–39
- DMPO solution, *in situ* ESR experiments, fuel
cell technology, 224–226
- Dodecyl-trimethylammonium bromide (DTAB),
dendrimer structure, spin probe
analysis, 286–289
- Doped conductive polymers:
charge carrier relaxation and dynamics,
317–326, 321–326
charge transfer reactions, 326–328
- d* orbital spin density, double resonance ESR, 31
- Double electron-electron resonance (DEER):
[2]catenane coconformation and motional
modes, 179–182
experimental techniques, 33–35
mesoscopic ionomers, 189–192
polyelectrolyte-counterion interactions, 176–178
- Double resonance ESR methods:
basic principles, 26–27
electron-electron double resonance, 31–39
basic principles, 31–33
distance distribution computation, 38–39
experimental techniques, 33–35
scattering, form factor and structure factor,
35–38
- electron-nuclear double resonance, 39–43
basic principles, 39–40
experimental techniques, 40–42
high-field ENDOR, 42–43
- electron spin echo envelope modulation, 44–49
basic principles, 44
data analysis, 47–49
experimental techniques, 44–47
- spin-spin couplings, 27–31
dipole-dipole coupling, 27–29
exchange coupling, 30
Fermi contact interaction, 30–31
spin density, *p* and *d* orbitals, 31
- Dow ionomers, fluorinated radicals, 215–221
- DOXYL nitroxides:
amphiphilic probes, 170–171
mesoscopic interfaces and nanostructure,
polymer lattices, 185–187
spin labeling and probing, 168
- Dynamic electrostatic attachment,
polyelectrolyte-counterion interactions,
175–178
- Dynamic light scattering (DLS), poly(2-
hydroxyethyl methacrylate) segmental
mobility, 159–161
- Dysonian ESR spectrum, doped conductive
polymers, charge transfer reactions,
326–328
- Echo-detected electron spin resonance:
anisotropic hyperfine interaction and *g*-tensor,
10–12
mesoscopic ionomers, 192
- Electron-electron double resonance (ELDOR),
31–39
basic principles, 31–33
dipole-dipole coupling, 27–29
distance distribution computation, 38–39
experimental techniques, 33–35
pulsed ESR methods, 19–22
scattering, form factor and structure factor,
35–38
weak coupling separation, 26
- Electron-nuclear double resonance (ENDOR),
39–43
basic principles, 39–40
conductive polymers, 310
dipole-dipole coupling, 27–29
experimental techniques, 40–42
high-field ENDOR, 42–43
isotropic hyperfine analysis, 12
metallo-dendrimer structures, 300
multifrequency and high-field electron spin
resonance, 17–18

- pulsed ESR methods, 19–22
- weak coupling separation, 26
- Electron spin echo envelope modulation (ESEEM), 44–49
 - basic principles, 44
 - conductive polymers, 310
 - data analysis, 47–49
 - dipole-dipole coupling, 27–29
 - double-resonance techniques, 47
 - experimental techniques, 44–47
 - multifrequency and high-field electron spin resonance, 17–18
 - polyelectrolyte-counterion interactions, 175–178
 - pulsed ESR methods, 19–22
- Electron spin resonance (ESR):
 - anisotropic hyperfine interactions and *g*-tensor, 10–12
 - basic principles, 3–9
 - environmental effects, 12–13
 - imaging studies:
 - basic principles, 85–87
 - polymeric systems, 93–96
 - spatially resolved degradation, 230–251
 - aged heterophasic polymers, 238–251
 - experimental protocols, 233–235
 - hindered amine stabilizers, 235–238
 - system components, 87–93
 - hardware requirements, 88–90
 - magnetic field gradients, 87–88
 - one-dimensional intensity profiling, 90–92
 - two-dimensional line shape profiling, 92–93
 - isotropic hyperfine analysis, 12
 - line shape analysis, tumbling nitroxide radicals, 15–16
 - multifrequency and high-field methods, 16–18
 - paramagnetic quencher accessibility, 13–15
 - pulsed methods, 18–22
- Electron Zeeman (EZ) interaction, basic principles, 4–9
- Elliot mechanism, conductive polymers, high-field ESR, magnetic parameters, 314–317
- Elongated spot distribution, spin-labeled polymer orientation, 67–68
- Energy potential, spin-labeled polymer orientation, 65–68
- Environmental effects, *g*-tensor and hyperfine interaction, 12–13
- EPRLL program:
 - available packages, 81–82
 - microscopic order-macroscopic disorder model, 68–70, 69–70
 - nonlinear least-squares analysis, 80–81
 - slowly relaxing local structure model (SRLS) model, 70–73
 - slow-motional ESR analysis, 55
 - diffusion models, 62–63
 - nitroxide magnetic tensors, 59–60
 - spin-labeled polymer orientation, 63–68
 - stochastic Liouville equation basis sets, 73–78
 - pruning, 78–79
- Ether bridges, fuel cell membranes, 207–210
- Euler angles:
 - nitroxide diffusion models, 62–63
 - spin-labeled polymer orientation, 64–68
- Exchange coupling, dipole-dipole interactions, 30
- Far-infrared (FIR) electron spin resonance (ESR), polystyrene segmental mobility, 147–155
- Fast internal motion (FIM) model, slow-motion ESR spectra, 71–73
- Fast-motional spectra, nitroxide line shape analysis, 138–139
- Fenton media, fuel cell membrane stability, 202–204
 - direct ESR and spin trapping detection, radical intermediates, 204–206
- Fermi contact interaction, double resonance ESR, 30–31
- Fermi velocity, conductive polymers, high-field ESR, charge carrier relaxation and dynamics, 324–326
- Ferromagnetic exchange coupling, electron spin resonance principles, 8–9
- Fluorescence analysis, polymer dynamics, 134–136
- Fluorinated alkyl radicals:
 - Nafion membrane in Fenton media, 214–215
 - UV-irradiated Nafion and Dow monomers, 215–221
- Fourier transform (FT):
 - electron-electron double resonance:
 - distance distributions, 38–39
 - form and structure scattering, 35–38
 - one-dimensional electron spin resonance imaging, intensity profiling, 91–92
- Fourier transform infrared (FTIR) spectroscopy:
 - fuel cell membrane stability, 203–204
 - spatially resolved degradation, electron spin resonance imaging, 233, 248–250
- Free induction decay (FID), pulsed ESR methods, 20–22
- Free ions, polyelectrolyte-counterion interactions, 173–178

- Free-radical reactions, polymer coating
 - photooxidation and stabilization, 257–260
 - nitroxide decay experiments, 263–265
- Fremy's salt dianion, polyelectrolyte-counterion interactions, 174–178
- Frozen solutions, solution-based macromolecules, 172–173
- Fuel cells:
 - basic elements, 198–201
 - membrane stability, 201–204
 - future research issues, 224–226
 - Nafion membranes in Fenton media, 210–215
 - fluorinated alkyl radicals, 214–215
 - oxygen-centered radicals, 211–213
 - oxygen radical reactions, 206–210
 - in situ* ESR experiments, 222–224
 - UV-irradiated Nafion and Dow ionomers, 215–221
 - fluorinated radical fragments, 217–221
 - radical intermediate detection, direct ESR and spin trapping, 204–206
- Gas voltaic battery, fuel cell technology, 198–201
- Gaussian line shape analysis, nitroxide magnetic tensors, 59–60
- Gauss-Newton method, nonlinear least-squares analysis, 80–81
- Genetic algorithm, one-dimensional electron spin resonance imaging, intensity profiling, 92
- Gibbs free energy, fuel cell membrane stability, 203–204
- Glass transition temperature:
 - mesoscopic interfaces and nanostructure, polymer lattices, 183–187
 - solution-based macromolecules, frozen solutions, 172–173
 - spin labeling and, 167
- Gradient coils, electron spin resonance imaging, 89–90
- g*-tensor:
 - anisotropic hyperfine interactions, 10–12
 - conductive polymers, high-field ESR, magnetic parameters, 311–317
 - electron spin resonance, basic principles, 4–9
 - environmental effects, 12–13
 - nitroxide magnetic tensors, 55–60
 - nitroxide rotational dynamics, 136–140
- Hahn echo experiments, pulsed ESR methods, 20–22
- Hamiltonian equations, electron spin resonance principles, 4–9
- Hardware requirements, electron spin resonance imaging, 88–90
- Heterophasic polymers, electron spin resonance imaging, spatially resolved degradation, 230–251
 - aged heterophasic polymers, 238–251
 - poly(acrylonitrile-butadiene-styrene) systems, 238–243
 - propylene-ethylene copolymers, 243–248
 - crystalline polymers microtome sensitivity, 250–251
 - experimental protocols, 233–235
 - FTIR comparison, 248–250
 - hindered amine stabilizers, 235–238
 - nitroxides derived from hindered amine stabilizers, 235–238
- Heterophasic propylene-ethylene copolymers (HPEC):
 - photodegradation, 244–248
 - spatially resolved degradation:
 - basic properties, 233–235
 - ESRI vs. FTIR, 249–250
 - thermal aging, 243–244
- High-field electron-nuclear double resonance (ENDOR), basic principles, 42–43
- High-field electron spin resonance:
 - basic principles, 16–18
 - conductive polymers:
 - basic principles, 307–310
 - carrier relaxation and dynamics, 317–326
 - charge carrier magnetic parameters, 310–317
 - highly-doped charge transfer, 326–328
 - saturation transfer, 328–330
 - spin probes, 330–333
- High-spin polycarbenes, dendrimer spin labeling, 290–296
- Hindered amine stabilizers (HAS):
 - electron spin resonance imaging:
 - nitroxide derivation, 235–238
 - polymeric formation from HAS, 94–96
 - spatially resolved degradation, 230–233
 - heterophasic propylene-ethylene copolymers, thermal aging, 243–244
 - polymer coating photooxidation and stabilization:
 - basic properties, 257–258
 - ESR-based nitroxide concentration, 260–262
 - free-radical reactions, 259–260
 - nitroxide kinetics, 272–276
 - nitroxide measurements, 264
 - photoinitiation rate measurements:
 - nonweathered coatings, 269
 - weathered coatings, 272

- Houzé-Nechtschein model, conductive polymers, high-field ESR, magnetic parameters, 313–317
- Hydrogen bonding:
local polarity, 13
solution-based macromolecules, solvent and temperature effects, 172
- Hydrogen peroxide:
fuel cell membrane stability, 202–204
direct ESR and spin trapping detection, 204–206
Nafion membranes in Fenton media, 210–215
toluene sulfonic acid reaction, 207–210
UV-irradiated Nafion and Dow ionomers, 215–221
- Hydroperoxyl radicals (HOO•):
fuel cell membrane stability, 202–204
Nafion membranes in Fenton media, 210–215
fluorinated alkyl radicals, 214–215
polymer coating photooxidation and stabilization, 258–260
toluene sulfonic acid reaction, 207–210
- Hydrophobic probes, strategic applications, 169
- 2-Hydroxyethyl methacrylate (HEMA), segmental mobility, 155–161
- Hydroxyl radicals (HO•):
fuel cell membrane stability, 202–204
direct ESR and spin trapping detection, 204–206
toluene sulfonic acid reaction, 206–210
- Hyperfine interactions:
anisotropic parameters and g -tensor, 10–12
dendrimer spin labeling, 289–296
double resonance ESR, 26–27
electron spin echo envelope modulation, 44–49
electron spin resonance, basic principles, 5–9
ESR analysis, radical polymerization:
chain length dependence, 117–119
penultimate unit effect, 123–124
isotropic analysis, 12
Nafion and Dow ionomers, membrane-derived fluorinated radicals, 217–221
nitroxide magnetic tensors, 55–60
pulsed ESR methods, 18–22
- Hyperfine sublevel correlation (HYSCORE) experiments:
double-resonance techniques, 27
electron spin echo envelope modulation, 47–49
metallo-dendrimer structures, 300–301
pulsed ESR methods, 19–22
- Imaging studies with ESR:
basic principles, 85–87
polymeric systems, 93–96
system components, 87–93
hardware requirements, 88–90
magnetic field gradients, 87–88
one-dimensional intensity profiling, 90–92
two-dimensional line shape profiling, 92–93
- “Immobilized” nitroxides, polymer coating photooxidation and stabilization, 261–262
- Infrared spectroscopy, photoinitiation rate measurements, weathered coatings, 271–272
- In situ* electron spin resonance, fuel cell membrane stability, 222–224
- Intensity profiling, one-dimensional electron spin resonance imaging, 90–92
- Intercluster distance, mesoscopic ionomers, 188–192
- Interpulse delay, electron spin echo envelope modulation, 45–49
- Ionic probes:
polyelectrolyte-counterion interactions, 174–178
- Ionomers:
Dow ionomers, membrane-derived fluorinated radicals, 215–221
mesoscopic interfaces and nanostructure, 187–192
- Iron complexes:
metallo-dendrimer structures, 300–301
Nafion and Dow ionomers, membrane-derived fluorinated radicals, 215–221
- Isotropic values:
electron spin resonance:
basic principles, 5–9
hyperfine analysis, 12
hyperfine analysis, 12
membrane-derived fluorinated radicals in Nafion, 218–221
- J couplings, dipole-dipole interactions, 30
- Ketone formation, polymer coating photooxidation and stabilization, photoinitiation rate measurements, 269
- Kinetic models, polymer coatings, nitroxide kinetics and HALS stabilization, 272–276
- Kivelson theory, high-field electron spin resonance, conductive polymers, charge carrier relaxation and dynamics, 320–326
- Kramers’ theory:
polymer dynamics, NMR and fluorescence techniques, 134–135
polystyrene segmental mobility, 149–155

- K* values, stochastic Liouville equation truncation parameters, 76–78
- Least-squares analysis, nitroxide magnetic tensors, 57–60
- Levenberg-Marquardt algorithm, nonlinear least-squares analysis, 80–81
- Ligand-to-metal charge transfer (LMCT), Nafion membrane radicals in Fenton media, 210–215
- Linear charge density, polyelectrolyte-counterion interactions, 174–178
- Line shape analysis:
 - [2]catenanes, 181–182
 - conductive polymers, high-field ESR, magnetic parameters, 311–317
 - ESR spectrum calculations, 55
 - nitroxide magnetic tensors, 59–60
 - nitroxide rotational dynamics, 136–140
 - fast-motional spectra, 138–139
 - slow-motional spectra and ordering, 139–140
 - polystyrene segmental mobility, 150–155
 - tumbling nitroxide radicals, 15–16
 - two-dimensional spectral-spatial ESRI, 92–93
- Liquid crystalline structures in dendrimers, 299
- Longitudinal relaxation:
 - paramagnetic quencher accessibility, 14–15
 - pulsed ESR methods, 20–22
- Lorentzian line shapes:
 - [2]catenanes, 181–182
 - conductive polymers, magnetic parameters, 315–317
 - nitroxide rotational dynamics, 138–139
 - polystyrene segmental mobility, 149–155
- L* values, stochastic Liouville equation truncation parameters, 76–78
- Macrocycle motion, [2]catenane coconformation and motional modes, 179–182
- Macromolecular materials:
 - dendrimer structures, ESR analysis:
 - basic properties, 279–282
 - metallo-dendrimers, 299–301
 - nitroxide spin labeling, 289–296
 - paramagnetic metal ion additions, 296–299
 - spin probes, 283–289
 - mesoscopic interfaces and nanostructure, 182–187
 - ionomers, 187–192
 - solution-based ESR, 171–182
 - [2]catenane coconformation and motional modes, 178–182
 - frozen solutions, 172–173
 - polyelectrolyte-counterion interactions, 173–178
 - solvent and temperature effects, 171–172
- spin labeling:
 - research overview, 166–167
 - strategies for, 167–168
- spin probing:
 - amphiphilic probes, 170–171
 - hydrogen-bonding probes, 169
 - hydrophobic probes, 169
 - ionic probes, 169–170
 - strategies for, 168–171
- Magnetic frame, nitroxide spin labels, 61–62
- Magnetic parameters:
 - conductive polymers, high-field ESR:
 - charge carriers, 310–317
 - spin probe method, 332–333
 - dendrimer structures, 302–303
 - electron spin resonance imaging, 87–88
 - membrane-derived fluorinated radicals in Nafion, 218–221
 - one-dimensional electron spin resonance imaging, intensity profiling, 90–92
- Magnetic tensors, nitroxide labels, 55–60
- Magnetic tilt angles, nitroxide magnetic tensors, 56–60
- Manganese complexes, dendrimer structure, ESR analysis, paramagnetic metal ion additions, 297–299
- Manning parameter, polyelectrolyte-counterion interactions, 173–178
- Maximum dipolar time evolution, electron-electron double resonance, 32–33
- Membrane electrode assembly (MEA), fuel cell technology, 198–201
- Membrane stability, fuel cell technology, 201–204
 - future research issues, 224–226
 - Nafion membranes in Fenton media, 210–215
 - fluorinated alkyl radicals, 214–215
 - oxygen-centered radicals, 211–213
 - oxygen radical reactions, 206–210
 - in situ* ESR experiments, 222–224
 - UV-irradiated Nafion and Dow ionomers, 215–221
 - fluorinated radical fragments, 217–221
- Mesososcopic interfaces and nanostructure, macromolecular materials, 182–192
 - ionomers, 187–192
 - polymer lattices, 182–187
- Metallo-dendrimers, ESR analysis, 299–301
- Methacrylates, ESR analysis:
 - chain-transfer reaction, 111–119
 - penultimate unit effect, dimeric model radicals, 124–128

- Micelle concentrations, dendrimer structure, spin probe analysis, 285–289
- Microscopic order-macroscopic disorder (MOMD) model:
 nitroxide rotational dynamics, slow-motional spectra, 140
 poly(2-hydroxyethyl methacrylate) segmental mobility, 156–161
 polystyrene segmental mobility, 147–155
 slowly relaxing local structure model, 71–73
 slow-motion ESR spectra, 68–70
 stochastic Liouville equation truncation parameters, 75–78
- Microtoming sensitivity, crystalline polymers, 250–251
- Mims electron-nuclear double resonance, experimental techniques, 41–42
- Minimum truncation set (MTS), stochastic Liouville equation, 75–78
 basis set pruning, 78–79
- Molybdenum complexes, metallo-dendrimer structures, 301
- Monomeric radicals, ESR analysis:
 activation rate constants, 128–129
 chain length dependence, 117–119
 penultimate unit effect, 119–124
- Monte Carlo procedures, one-dimensional electron spin resonance imaging, intensity profiling, 91–92
- Motional modes:
 [2]catenanes, 178–182
 Nafion membrane radicals in Fenton media, 211–215
- Multifrequency electron spin resonance, basic principles, 16–18
- Nafion:
 fluorinated radicals in UV-irradiated compounds, 215–221
 fuel cell technology, 198–201
 radical intermediate detection, 205–206
 radical species in Fenton media, 210–215
- Nanomaterials:
 [2]catenane coconformation and motional modes, 178–182
 mesoscopic interfaces and nanostructure:
 ionomers, 187–192
 polymer lattices, 182–187
- Nelder-Mead search, nonlinear least-squares analysis, 80–81
- Nickel catalysts, metallo-dendrimer structures, 301
- Nitroxide compounds:
 anisotropic hyperfine interaction and *g*-tensor, 10–12
 electron spin resonance imaging, 87–88
 hindered amine stabilizers, 235–238
 spatially resolved degradation, 231–233
 environmental effects on *g*-tensor and hyperfine interactions, 12–13
 heterophasic propylene-ethylene copolymers:
 photodegradation, 244–248
 thermal aging, 243–244
 line shape analysis:
 rotational effects, 136–140
 fast-motional spectra, 138–139
 slow-motional spectra and ordering, 139–140
 tumbling radicals, 15–16
 microscopic order-macroscopic disorder model, 69–70
 polymer coating photooxidation and stabilization:
 decay experiments, 263–264
 ESR-based concentration, 260–262
 free-radical decay kinetics and photoinitiation rates, 264–265
 HALS-stabilized sample preparation, 264
 kinetics, 272–276
 photoinitiation rate measurements:
 nonweathered coatings, 265–269
 weathered coatings, 269–272
 UV exposures, 262–263
 radical intermediate detection, spin trapping techniques, 205–206
 segmental dynamics, 140–141
 spin labeling and probing:
 TEMPO and PROXYL labels, 168–169
- Non-Brownian diffusion models, slow-motional ESR analysis, 62–63
- Nonlinear least-squares analysis (NLLS):
 poly(2-hydroxyethyl methacrylate) segmental mobility, 156–161
 polystyrene segmental mobility, 148–155
 slow-motional ESR, 79–81
- Nonweathered coatings, photoinitiation rate measurements, 265–269
- Nuclear frequency spectrum, electron spin echo envelope modulation, 45–49
- Nuclear Overhauser effect (NOE) enhancements, polymer dynamics, 134
- Nuclear Zeeman (NZ) interaction, basic principles, 6–9
- Odijk-Skolnick-Fixman theory, polyelectrolyte-counterion interactions, 174–178
- Oligomeric radicals, ESR analysis, controlled chain lengths, 109–111

- One-dimensional electron spin echo envelope modulation, data analysis, 48–49
- One-dimensional electron spin resonance imaging:
 basic components, 87–88
 intensity profiling, 90–91
 polymeric systems studied, 94–96
 spatially resolved degradation, 232–233
 aged heterophasic polymers, 238–251
- Optimization techniques, one-dimensional electron spin resonance imaging, intensity profiling, 90–92
- Orientalional ordering:
 nitroxide rotational dynamics, 139–140
 poly(2-hydroxyethyl methacrylate) segmental mobility, 156–161
 polystyrene segmental mobility, 151–155
 spin-labeled polymer orientation, 63–68
- Oxygen centered radicals:
 ESR analysis, controlled chain lengths, 116–119
 Nafion membrane radicals in Fenton media, 211–215
- Pair distance distribution function, electron-electron double resonance, 35–38
- Pake pattern, double resonance ESR, dipole-dipole coupling, 29
- PANI conductive polymer:
 charge carrier relaxation and dynamics, 321–326
 charge transfer reactions, 327–328
 magnetic parameters, 314–317
 saturation-transfer ESR, 329–330
- Paramagnetic parameters:
 conductive polymers, high-field ESR, 310
 dendrimer structures, ion additions, 296–299
 electron spin resonance principles, 8–9
 ESR imaging studies, 85–87
 quencher accessibility, 13–15
- Pauli susceptibility, high-field electron spin resonance, conductive polymers, magnetic parameters, 315–317
- Penultimate unit effect, ESR analysis, radical polymerization, 119–129
 activation rate constants, 128–129
 dimeric model radicals, 124–128
 monomeric model radicals, 122–124
- Persistence length, polyelectrolyte-counterion interactions, 174–178
- Phenoxyl radicals, toluene sulfonic acid reaction, 207–210
- Photoacoustic (PA) FTIR spectroscopy, acrylonitrile-butadiene-styrene polymers, 249–250
- Photodegradation:
 acrylonitrile-butadiene-styrene polymers, 241–243
 heterophasic propylene-ethylene copolymers, 244–248
 FTIR spectroscopy, 249–250
- Photoinitiation rate (PIR), polymer coating photooxidation and stabilization:
 free-radical reactions, 258–260
 nonweathered coatings, free-radical decay kinetics, 265–269
 weathered coatings, 269–272
- Photooxidation, polymer coating stabilization:
 basic principles, 256–258
 free-radical processes, 258–260
 nitroxide decay, 264–272
 nitroxide concentration by ESR, 260–262
 nitroxide decay sample preparation, 263–264
 nitroxide kinetics and HALS stabilization, 272–277
 ultraviolet exposures, 262–263
- π -conjugated polymers, high-field electron spin resonance, 308–310
- π radicals, isotropic hyperfine analysis, 12
- Platinum catalysts, *in situ* electron spin resonance experiments, fuel cell membrane stability, 222–224
- Poisson-Boltzmann equation:
 electron-electron double resonance, form factor and structure factor in scattering, 36–38
 polyelectrolyte-counterion interactions, 173–178
- Polar angles, spin-labeled polymer orientation, 64–68
- Polaron motion:
 conductive polymers, high-field ESR:
 magnetic parameters, 312–317
 spin probe method, 331–333
 high-field electron spin resonance, conductive polymers, charge carrier relaxation and dynamics, 321–326
- Polyacetylene (PA), high-field electron spin resonance, 308–310
- Polyacrylates, ESR analysis, controlled chain lengths, 116–119
- Poly(acrylic acid) (PAA), segmental mobility, 141–147
- Polyamidoamine (PAMAM), dendrimer structure, ESR analysis, 280–282
 nitroxide spin labeling, 291–296
 paramagnetic metal ion additions, 296–299
 spin probe techniques, 284–289

- Polybutadiene (PB), electron spin resonance imaging, spatially resolved degradation, 233–235
- Poly(butylacrylate) (P(BA)), mesoscopic interfaces and nanostructure, polymer lattices, 183–187
- Poly(butylmethacrylate) (PBA), mesoscopic interfaces and nanostructure, 183–187
- Poly(diallyldimethylammonium chloride) (PDADMAC), polyelectrolyte-counterion interactions, 174–178
- Polyelectrolytes:
counterion interactions, solution-based macromolecules, 173–178
segmental mobility, 141–147
- Poly(2-hydroxyethyl methacrylate) (PHEMA), segmental mobility, 141, 155–161
- Polymer coatings, photooxidation and stabilization:
basic principles, 256–258
free-radical processes, 258–260
nitroxide decay, 264–272
nitroxide concentration by ESR, 260–262
nitroxide decay sample preparation, 263–264
nitroxide kinetics and HALS stabilization, 272–277
ultraviolet exposures, 262–263
- Polymeric systems, electron spin resonance imaging, 93–96
- Polymer lattices, mesoscopic interfaces and nanostructure, 182–187
- Poly(methacrylic acid) (PMA), segmental mobility, 141–147
- Poly(methyl methacrylate) (polyMMA) radicals:
ESR analysis, controlled chain lengths, 109–111
segmental mobility, 141–142
- Polynucleotides, dendrimer structures, spin labeled nitroxides, 293–296
- Poly(propyleneimine) (PPI) dendrimer:
paramagnetic metal ion additions, 298–299
spin labeling with nitroxides, 289–296
spin probe analysis, 283–289
- Polypropylenepolyamine dendrimer, spin probe analysis, 283–289
- Polypropylene (PP), spatially resolved degradation, basic properties, 233–235
- Polystyrene (PS), segmental mobility, 141, 147–155
- Polythiophene, conductive polymers, high-field ESR, magnetic parameters, 312–317
- Population distributions, spin-labeled polymer orientation, 65–68
- p* orbital spin density, double resonance ESR, 31
- Probe ordering, spin-labeled polymer orientation, 63–68
- Projection slice algorithm, two-dimensional electron spin resonance imaging, line shape profiling, 92–93
- Proton exchange membrane (PEM):
direct ESR and spin trapping detection, radical intermediates, 205–206
fuel cell technology, 198–201
future research issues, 224–226
oxygen radical reactions, 210–215
stability in, 201–204
- PROXYL nitroxides, spin labeling and probing, 168–169
dendrimer structures, 289–296
- Pruning, stochastic Liouville equation basis sets, 78–79
- Pseudo-spherical tensor components, nitroxide magnetic tensors, 59–60
- Pulsed electron-electron double resonance:
experimental techniques, 33–35
form factor and structure factor in scattering, 35–38
mesoscopic ionomers, 189–192
- Pulsed electron spin resonance:
basic principles, 18–22
dendrimer spin labeling, 290–296
metallo-dendrimer structures, 300–301
electron-electron double resonance, 33–35
DEER sequences, 33–35
- Pump pulse, electron-electron double resonance, 31–33
- bis[2-(2-Pyridyl)ethyl]-amine (PY2) ligands, dendrimer structures, paramagnetic metal ion additions, 298–299
- α -(4-Pyridyl-1-oxide)-*N-tert* butylnitron (POBN), *in situ* ESR experiments, fuel cell technology, 223–225
- Quadruple interaction, basic principles, 6–9
- Quartet ESR spectra, Nafion and Dow ionomers, membrane-derived fluorinated radicals, 216–217
- Quasi-one-dimensional (Q1D) systems, high-field electron spin resonance, conductive polymers, 308–310
charge carrier relaxation and dynamics, 319–326
magnetic parameters, 315–317
- Quenchers, paramagnetic, accessibility to, 13–15
- Quintet ESR spectra, Nafion and Dow ionomers, membrane-derived fluorinated radicals, 215–221

- Radial distribution function, polyelectrolyte-counterion interactions, 176–178
- Radical intermediates:
 direct ESR and spin trapping detection, 204–206
 ESR imaging, 230–233
- Radical polymerization. *See also* Free-radical formation
- ESR analysis:
 basic principles, 102–105
 chain length dependence, 105–111
 controlled chain length models, 106–111
tert-butyl methacrylate, 105–106
 chain transfer reaction, 111–119
 chain length-dependent changes, 117–119
 controlled chain lengths, 113–117
tert-butyl acrylate, 111–113
 penultimate unit effect, 119–129
 activation rate constants, 128–129
 dimeric model radicals, 124–128
 monomeric model radicals, 122–124
- Radical-radical collisions, dipole-dipole interactions, 30
- Relative anisotropy, line shape analysis, tumbling nitroxide radicals, 15–16
- Relaxation times:
 dendrimer structures, 302–303
 electron-electron double resonance, 32–33
 high-field electron spin resonance, conductive polymers, charge carrier relaxation and dynamics, 321–326
 mesoscopic ionomers, 191–192
- Resonant radio frequency irradiation, electron-nuclear double resonance, 40–42
- Rhombic distortion:
 poly(2-hydroxyethyl methacrylate) segmental mobility, 156–161
 polystyrene segmental mobility, 151–155
- Rotational correlation time (τ):
 dendrimer structures:
 paramagnetic metal ion additions, 298–299
 spin probe analysis, 284–289
 electron-electron double resonance, 33–35
 electron spin echo envelope modulation, 45–49
 line shape analysis, tumbling nitroxide radicals, 15–16
 nitroxide line shape analysis, 136–140
 nitroxide spin label dynamics, 140–141
 polyelectrolyte-counterion interactions, 175–178
 poly(2-hydroxyethyl methacrylate) segmental mobility, 157–161
 polystyrene segmental mobility, 152–155
 segmental mobility of polymer radicals, 145–147
 solution-based macromolecules, solvent and temperature effects, 171–172
- Rotational diffusion frame:
 nitroxide spin labels, 61–62
 polyelectrolyte-counterion interactions, 174–178
 slowly relaxing local structure model, 71–73
- Ruthenium complexes, dendrimer structure, spin probe analysis, 288–289
- Saturation recovery, mesoscopic ionomers, 191–192
- Saturation-transfer ESR (ST-ESR), conductive polymers, 328–330
- Scanning electron microscopy (SEM), heterophasic propylene-ethylene copolymers, 234–235
- Schneider-Freed programs, nitroxide rotational dynamics, slow-motional spectra, 139–140
- Segmental mobility, solution-based polymer dynamics:
 NMR and fluorescence techniques, 134–135
 spin-label electron spin resonance, 140–162
 nitroxide spin labels, 140–141
 poly(2-hydroxyethyl methacrylate), 155–161
 poly(methacrylic acid) and poly(acrylic acid), 142–147
 poly(methyl methacrylate), 142
 polystyrene, 147–155
- Semiquinone radicals, hydroxyl radicals, 207–210
- Single crystal samples:
 anisotropic hyperfine interaction and *g*-tensor, 10–12
 nitroxide magnetic tensors, 56–60
- Site-bound ions, polyelectrolyte-counterion interactions, 173–178
- Slowly relaxing local structure model (SRLS) model, slow-motion ESR spectra, 70–73
- Slow-motion ESR spectra:
 nitroxide parameters, 55–73
 diffusion models, 60–63
 magnetic tensors, 55–60
 microscopic order-macroscopic disorder model, 69–70
 slowly relaxing local structure model, 70–73
 spin-labeled polymer orientation, 63–68
 nitroxide rotational dynamics, 139–140
 nonlinear least-squares analysis (NLLS), 79–81

- program availability, 81–82
- stochastic Liouville equation basis sets, 73–79
 - calculation, 73–74
 - pruning, 78–79
 - symmetrization, 74
 - truncation parameters, 74–78
- Small angle neutron scattering (SANS), electron-electron double resonance, 35–38
- Small angle X-ray scattering (SAXS):
 - electron-electron double resonance, 35–38
 - mesoscopic interfaces and nanostructure, ionomers, 187–192
 - poly(2-hydroxyethyl methacrylate) segmental mobility, 159–161
- Solution-based macromolecules, 171–182
 - [2]catenane coconformation and motional modes, 178–182
 - frozen solutions, 172–173
 - polyelectrolyte-counterion interactions, 173–178
 - solvent and temperature effects, 171–172
- Solution-based polymer dynamics, spin-labeled ESR analysis:
 - basic principles, 134
 - nitroxide rotational effects, ESR line shapes, 136–140
 - fast-motional spectra, 138–139
 - slow-motional spectra and ordering, 139–140
 - NMR and fluorescence analysis, 134–136
 - segmental mobility, 134–135
 - segmental mobility, 140–162
 - nitroxide spin labels, 140–141
 - poly(2-hydroxyethyl methacrylate), 155–161
 - poly(methacrylic acid) and poly(acrylic acid), 142–147
 - poly(methyl methacrylate), 142
 - polystyrene, 147–155
- Solvent effects:
 - polymer coating stabilization, 256–258
 - polystyrene segmental mobility, 151–155
- Spatially resolved degradation, electron spin resonance imaging, 230–251
 - aged heterophasic polymers, 238–251
 - poly(acrylonitrile-butadiene-styrene) systems, 238–243
 - propylene-ethylene copolymers, 243–248
 - crystalline polymers microtome sensitivity, 250–251
 - experimental protocols, 233–235
 - FTIR comparisons, 248–250
 - hindered amine stabilizers, 235–238
- Spatial resolution, electron spin resonance imaging, 87–88
- Spectral diffusion, pulsed ESR methods, 21–22
- Spin density, double resonance ESR, 31
- Spin diffusion, conductive polymers, charge carrier relaxation and dynamics, 318–326
- Spin labeling:
 - dendrimer structures, 289–296
 - macromolecular materials:
 - research overview, 166–167
 - strategies for, 167–168
 - orientational ordering, 63–68
 - solution-based polymer dynamics:
 - basic principles, 134
 - nitroxide rotational effects, ESR line shapes, 136–140
 - fast-motional spectra, 138–139
 - slow-motional spectra and ordering, 139–140
 - NMR and fluorescence analysis, 134–136
 - segmental mobility, 134–135
 - segmental mobility, 140–162
 - nitroxide spin labels, 140–141
 - poly(2-hydroxyethyl methacrylate), 155–161
 - poly(methacrylic acid) and poly(acrylic acid), 142–147
 - poly(methyl methacrylate), 142
 - polystyrene, 147–155
- Spin-orbit coupling, environmental effects on g -tensor and hyperfine interactions, 13
- Spin probing:
 - conductive polymers, high-field ESR, 330–333
 - dendrimer structures, ESR analysis, 283–289
 - macromolecular materials:
 - amphiphilic probes, 170–171
 - hydrogen-bonding probes, 169
 - hydrophobic probes, 169
 - ionic probes, 169–170
 - strategies for, 168–171
 - mesoscopic interfaces and nanostructure, polymer lattices, 183–187
- Spin-spin couplings:
 - conductive polymers, high-field ESR:
 - magnetic parameters, 314–317
 - spin probe method, 331–333
 - double resonance ESR methods, 27–31
 - dipole-dipole coupling, 27–29
 - exchange coupling, 30
 - Fermi contact interaction, 30–31
 - spin density, p and d orbitals, 31
- Spin trapping:
 - radical intermediate detection, 204–206
 - in situ* ESR experiments, fuel cell technology, 224
- Stabilization kinetics, polymer coatings:

- basic principles, 256–258
- free-radical processes, 258–260
 - nitroxide decay, 264–272
- nitroxide concentration by ESR, 260–262
- nitroxide decay sample preparation, 263–264
- nitroxide kinetics and HALS stabilization, 272–277
- ultraviolet exposures, 262–263
- Starburst dendrimers, ESR analysis, 280–282
- Steric hindrance, dendrimer structures, ESR analysis, 280–282
- Stochastic Liouville equation (SLE):
 - basis set:
 - calculation, 73–74
 - pruning, 78–79
 - symmetrization, 74
 - truncation parameters, 74–78
 - ESR spectrum calculations, 54–55
 - nitroxide rotational dynamics, slow-motional spectra, 139–140
 - nonlinear least-squares analysis, 80–81
 - slow-motion ESR spectra, diffusion models, 60–63
- Structure factor scattering, electron-electron double resonance, 35–38
- Styrene-acrylonitrile (SAN):
 - ESRI vs. FTIR, 249–250
 - spatially resolved degradation, 233–235
 - thermal aging, 241
- Sulfonic acid groups:
 - fuel cell membrane stability, 202–204
 - oxygen radical reactions, 206–210
- Superoxide radical anions, toluene sulfonic reaction, 207–210
- Supramolecular dendrimer complexes, spin probe analysis, 284–289
- Swept-field calculations, stochastic Liouville equation basis set pruning, 78–79
- Symmetrization:
 - poly(2-hydroxyethyl methacrylate) segmental mobility, 156–161
 - polystyrene segmental mobility, 150–155
 - stochastic Liouville equation basis sets, 74
- Teflon surfaces:
 - ESR imaging, spatially resolved degradation studies, 230–233
 - in situ* ESR experiments, fuel cell technology, 223–224
- Temperature dependence:
 - conductive polymers, high-field ESR:
 - charge carrier relaxation and dynamics, 319–326
 - magnetic parameters, 312–317
 - ESR analysis of radical polymerization, controlled chain lengths, 108–111
 - poly(2-hydroxyethyl methacrylate) segmental mobility, 156–161
 - polystyrene segmental mobility, 151–155
 - solution-based macromolecules, 171–172
- TEMPO-4-carboxylate spin probe, strategic applications, 170
- TEMPOL probes, hydrogen-bond acceptors, 169
- TEMPO nitroxides:
 - dendrimer structures:
 - spin labeling, 291–296
 - spin probe analysis, 284–289
 - hydrophobic probes, 169
 - mesoscopic interfaces and nanostructure:
 - ionomers, 187–192
 - polymer lattices, 184–187
 - polymer coating photooxidation and stabilization, decay experiments, 263–264
 - spin labeling and probing, 168–169
- TEMPO-4-phosphonoxyolate spin probe, strategic applications, 170
- Territorially bound ions, polyelectrolyte-counterion interactions, 173–178
- tert*-Butyl acrylate radicals, ESR analysis:
 - chain length dependence, 117–119
 - controlled chain lengths, 113–119
 - penultimate unit effect, 119–129
- tert*-Butyl methacrylate (tBMA), ESR analysis:
 - chain length dependence, 105–106
 - controlled chain lengths, 106–111
- 7,7,8,8-Tetracyanoquinodimethane (TCNQ), spin probe analysis, 283–289
- Tetrathiafulvalene (TTF), dendrimer spin labeling, 291–296
- Thermal aging:
 - acrylonitrile-butadiene-styrene polymers, 238–241
 - heterophasic propylene-ethylene copolymers, 243–244
- Tikhonov regularization, electron-electron double resonance, distance distribution computation, 38–39
- TIN 440/770/292 compounds, nitroxide kinetics and HALS stabilization, 273–276
- Titanium complexes, Nafion membrane radicals in Fenton media, 210–215
- fluorinated alkyl radicals, 214–215
- Toluene, segmental mobility of polymers in, 149–155
- Toluene sulfonic acid, oxygen radical reactions, 206–210

- Trajectory method, ESR spectrum calculations, 54–55
- Transition metal ions:
dendrimer structures, additions to, 296–299
metallo-dendrimer structures, 299–301
electron spin resonance principles, 7–9
fuel cell membrane stability, 202–204
Nafion and Dow ionomers, membrane-derived fluorinated radicals, 215–221
- Triarylmethyl (TAM) trianion radical,
polyelectrolyte-counterion interactions, 178
- Truncation parameters, stochastic Liouville equation basis sets, 74–78
- Turning points, nitroxide magnetic tensors, 57–60
- Twisted boat configuration, nitroxide magnetic tensors, 57–60
- Two-dimensional electron spin resonance imaging:
line shape profiling, 92–93
spatially resolved degradation, 232–233
aged heterophasic polymers, 238–251
- Two-dimensional electron-spin transient nutation (2D-ESTN), dendrimer spin labeling, 290–296
- Two-pulse echo decay, mesoscopic ionomers, 191–192
- Ultraviolet (UV) radiation:
electron spin resonance imaging, spatially resolved degradation, 232–233
Nafion, membrane-derived fluorinated radicals, 215–221
polymer coating stabilization:
basic principles, 256–258
experimental protocols, 262–263
free-radical reactions, 259–260
- Urethane coatings, photoinitiation rate measurements, 271–272
- Vanadium catalysts, Nafion membrane radicals in Fenton media, 211
- Vanadyl complexes, dendrimer structures, paramagnetic metal ion additions, 298–299
- Viscosity dependence, polystyrene segmental mobility, 151–155
- Weathered polymer coatings, photoinitiation rate measurements, 269–272
- Weighting factors, stochastic Liouville equation basis set pruning, 78–79
- Wigner rotation matrix, polystyrene segmental mobility, 148–155
- X-ray photoelectron spectroscopy (XPS), Nafion membrane stability, 201
- X-ray powder diffraction (XRD), Nafion membrane stability, 201
- Zeeman interaction:
basic principles, 4–9
dipole-dipole coupling, 27–29
electron-nuclear double resonance, 40
electron spin echo envelope modulation, 44–49
electron spin resonance imaging, 89–90
high-field electron-nuclear double resonance, 43
- Zero-field splitting (ZFS):
dendrimer structure, ESR analysis:
paramagnetic metal ion additions, 297–299
electron-electron double resonance, 31–33
electron spin resonance principles, 8–9
multifrequency and high-field electron spin resonance, 17–18
- Zwitterionic diblock copolymers, mesoscopic ionomers, 189–192

